arXiv:math-ph/0411075v3 22 Jun 2005

# UNIVERSALITY IN RANDOM MATRIX THEORY FOR ORTHOGONAL AND SYMPLECTIC ENSEMBLES

PERCY DEIFT AND DIMITRI GIOEV

Abstract. We give a proof of universality in the bulk for orthogonal ($\beta = 1$) and symplectic ($\beta = 4$) ensembles of random matrices in the scaling limit for a class of weights $w(x) = e^{-V(x)}$ where $V$ is a polynomial, $V(x) = \kappa_{2m} x^{2m} + \cdots$, $\kappa_{2m} > 0$. For such weights the associated equilibrium measure is supported on a single interval. The precise statement of our results is given in Theorem 1.1 below. For a proof of universality in the bulk for unitary ensembles ($\beta = 2$), for the same class of weights, see [DKMVZ2].

Our starting point is Widom's representation [W] of the orthogonal and symplectic correlation kernels in terms of the kernel arising in the unitary case ($\beta = 2$) plus a correction term which is constructed out of derivatives and integrals of orthonormal polynomials (OP's) $\{p_j\}_{j\geq 0}$ with respect to the weight $w(x)$. The calculations in [W] in turn depend on the earlier work of Tracy and Widom [TW2]. It turns out (see [W] and also Theorems 2.1 and 2.2 below) that only the OP's in the range $j = N + O(1)$, $N \to \infty$, contribute to the correction term. In controlling this correction term, and hence proving universality for $\beta = 1$ and 4, the uniform Plancherel–Rotach type asymptotics for the OP's found in [DKMVZ2] play an important role, but there are significant new analytical difficulties that must be overcome which are not present in the case $\beta = 2$. We note that we do not use skew orthogonal polynomials.

## 1. Introduction

In this paper we will be concerned with ensembles of matrices $\{M\}$ with probability distributions

$$\mathcal{P}_{N,\beta}(M)\, dM = \frac{1}{\mathcal{Z}_{N,\beta}}\, e^{-\operatorname{tr} V_\beta(M)}\, dM, \tag{1.1}$$

for $\beta = 1$, 2 and 4, the so-called Orthogonal, Unitary and Symplectic ensembles, respectively (see [M]). For $\beta = 1$, 2, 4, the ensemble consists of $N \times N$ real symmetric matrices, $N \times N$ Hermitian matrices, and $2N \times 2N$ Hermitian self-dual matrices, respectively. In general the potential $V_\beta(x)$ is a real-valued function growing sufficiently rapidly as $|x| \to \infty$, but we will restrict our attention henceforth to $V_\beta$'s which are polynomials,

$$V_\beta(x) = \kappa_{2m,\beta} x^{2m} + \cdots, \qquad \kappa_{2m,\beta} > 0. \tag{1.2}$$

In (1.1), $dM$ denotes Lebesgue measure on the algebraically independent entries of $M$, and $\mathcal{Z}_{N,\beta}$ is a normalization constant. The above terminology for $\beta = 1$, 2 and 4 reflects the fact that (1.1) is invariant under conjugation of $M$, $M \mapsto UMU^{-1}$, by orthogonal, unitary and unitary-symplectic matrices $U$. It follows from (1.1) that the distribution of the eigenvalues $x_1, \cdots, x_N$ of $M$ is given (see [M]) by

$$P_{N,\beta}(x_1, \cdots, x_N) = \frac{1}{Z_{N,\beta}} \prod_{1 \le j < k \le N} |x_j - x_k|^\beta \prod_{j=1}^{N} w_\beta(x_j) \tag{1.3}$$

where again $Z_{N,\beta}$ is a normalization constant (partition function). Here

$$w_\beta(x) = \begin{cases} e^{-V_\beta(x)}, & \beta = 1, 2 \\ e^{-2V_\beta(x)}, & \beta = 4. \end{cases} \tag{1.4}$$

(The factor 2 in $w_{\beta=4}$ reflects the fact that the eigenvalues of self-dual Hermitian matrices come in pairs.) Let $\{p_j\}_{j \ge 0}$ be the normalized orthogonal polynomials (OP's) on $\mathbb{R}$ with respect to the weight $w \equiv w_{\beta=2}$, and define $\phi_j \equiv p_j w^{1/2}$. Note that $(\phi_j, \phi_k) = \delta_{jk}$ where $(\cdot, \cdot)$ denotes the standard inner product in $L^2(\mathbb{R})$.

For the unitary matrix ensembles ($\beta = 2$) an important role is played by the kernel

$$K_N(x, y) \equiv K_{N,2}(x, y) = \sum_{k=0}^{N-1} \phi_k(x)\, \phi_k(y). \tag{1.5}$$

In particular the probability density (1.3), the $l$-point correlation function $R_{N,l,2}$ and also the gap probability $E_2(0; J)$ that a set $J$ contains no eigenvalues, can all be expressed in terms of $K_N$, see e.g. [M]. For example

$$R_{N,l,2}(x_1, \cdots, x_l) = \det(K_N(x_j, x_k))_{1 \le j,k \le l}. \tag{1.6}$$

The Universality Conjecture, in our situation, states that the limiting statistical behavior of the eigenvalues $x_1, \cdots, x_N$ distributed according to the law (1.3), in the appropriate scale as $N \to \infty$, should be independent of the weight $w_\beta$, and should depend only on the invariance properties of $\mathcal{P}_{N,\beta}$, $\beta = 1$, 2 or 4, mentioned above. Universality has been considered extensively in the physics literature, see e.g. [BrZ, Be, HWe, SeV].

The kernel $K_N(x, y)$ can also be expressed via the Christoffel–Darboux formula

$$K_N(x, y) = b_{N-1}\, \frac{\phi_N(x)\phi_{N-1}(y) - \phi_{N-1}(x)\phi_N(y)}{x - y}, \tag{1.7}$$

where $b_{N-1}$ is a coefficient in the three-term recurrence relation for OP's (see (2.1) below). In view of the preceding remarks it follows that in the case $\beta = 2$, the study of the large $N$ behavior of $P_{N,2}$, and in particular the proof of universality, reduces to the asymptotic analysis of $b_{N-1}$ and the OP's $p_{N+j}$ with $j = 0$ or $-1$. By a fundamental observation of Fokas, Its and Kitaev [FoIKi] the OP's solve a Riemann–Hilbert problem (RHP) of a type that is amenable to the steepest descent method introduced by Deift and Zhou in [DZ] and further developed in [DVZ]. In [DKMVZ1, DKMVZ2] the authors analyzed the asymptotics of OP's for very general classes of weights. In particular they proved the Universality Conjecture in the case $\beta = 2$ for weights $w(x) = e^{-V(x)}$ where $V(x)$ is a polynomial as above, and also for $w(x) = e^{-NV(x)}$ where $V(x)$ is real analytic and $V(x)/\log|x| \to +\infty$, as $|x| \to \infty$. The (bulk) scaling limit $N \to \infty$ is described in terms of the so-called *sine kernel* $K_\infty(x - y)$ where

$$K_\infty(t) \equiv \frac{\sin \pi t}{\pi t}. \tag{1.8}$$

For example [DKMVZ2, Theorem 1.4], for $w(x) = e^{-V(x)}$, $V(x)$ polynomial, and for any $l = 2, 3, \cdots$ and $r, y_1, \cdots, y_l$ in a compact set, one has as $N \to \infty$

$$\frac{1}{(K_N(0,0))^l} R_{N,l,2}\Big(r + \frac{y_1}{K_N(0,0)}, \cdots, r + \frac{y_l}{K_N(0,0)}\Big) \to \det(K_\infty(y_j - y_k))_{1 \le j,k \le l}. \tag{1.9}$$

The scale $x = y/K_N(0,0)$ is chosen so that the expected number of eigenvalues per unit $y$-interval is one: This scaling is standard in Random Matrix Theory. Indeed for any Borel set $B \subset \mathbb{R}$,

$$\int_B R_{N,l=1,2}(x)\, dx = \mathbb{E}\{ \text{ number of eigenvalues in } B \}. \tag{1.10}$$

Thus by (1.6) $K_N(0,0) = R_{N,1,2}(0)$ gives the density of the expected number of eigenvalues near zero. In other words, in the appropriate scale, the large $N$ behavior of the eigenvalues is *universal* (i.e. independent of $V$). Pioneering mathematical work on the Universality Conjecture was done in [PS] and for the case of quartic two-interval potential $V(x) = N(x^4 - tx^2)$, $t > 0$ (sufficiently) large, in [BI]. We note again that all these results apply only in the case $\beta = 2$.

In the case $\beta = 1$ and 4 the situation is more complicated. In place of (1.5) one must use $2 \times 2$ matrix kernels (see e.g. [M, TW2])

$$K_{N,1}(x,y) = \begin{pmatrix} S_{N,1}(x,y) & (S_{N,1}D)(x,y) \\ (\epsilon S_{N,1})(x,y) - \frac{1}{2}\operatorname{sgn}(x-y) & S_{N,1}(y,x) \end{pmatrix}, \quad N \text{ even}, \tag{1.11}$$

and

$$K_{N,4}(x,y) = \frac{1}{2} \begin{pmatrix} S_{N,4}(x,y) & (S_{N,4}D)(x,y) \\ (\epsilon S_{N,4})(x,y) & S_{N,4}(y,x) \end{pmatrix}. \tag{1.12}$$

Here $S_{N,\beta}(x,y)$, $\beta = 1, 4$, are certain scalar kernels (see (1.17), (1.18) below), $D$ denotes the differentiation operator, and $\epsilon$ is the operator with kernel $\epsilon(x,y) = \frac{1}{2}\operatorname{sgn}(x-y)$[1]. Such matrix kernels were first introduced by Dyson [Dy2] in the context of circular ensembles with a view to computing correlation functions. Dyson's approach was extended to Hermitian ensembles, first by Mehta [M2] for $V(x) = x^2$, and then for more general weights by Mahoux and Mehta in [MaM]. A more direct and unifying approach to the results of Dyson–Mahoux–Mehta was given by Tracy and Widom in [TW2], where formulae (1.17), (1.18) below were derived. We see that once the kernels $S_{N,\beta}(x,y)$ are known, then so are the other kernels in $K_{N,\beta}$. As in the case $\beta = 2$, the kernels $K_{N,\beta}$ give rise to explicit formulae for $R_{N,l,\beta}$ and $E_\beta(0; J)$. For example

$$R_{N,1,\beta}(x) \equiv R_{1,\beta}(x) = \frac{1}{2} \operatorname{tr} K_{N,\beta}(x,x) \tag{1.13}$$

and

$$R_{N,2,\beta}(x,y) = \frac{1}{4}\big(\operatorname{tr} K_{N,\beta}(x,x)\big)\big(\operatorname{tr} K_{N,\beta}(y,y)\big) - \frac{1}{2}\operatorname{tr}\big(K_{N,\beta}(x,y)K_{N,\beta}(y,x)\big),$$

and so on, see [TW2]. As indicated above, formula (1.11) only applies to the case when $N$ is even. When $N$ is odd, there is a similar, but slightly more complicated, formula (see [AFNvM]). Throughout this paper, for $\beta = 1$, we will restrict our attention *to the case when $N$ is even.* We expect that the methods in this paper also extend to the case $\beta = 1$, $N$ odd, and we plan to consider this situation in a later publication. Of course, in situations where the asymptotics of (1.11) has been analyzed (e.g. $V(x) = x^2$) for all $N$ as $N \to \infty$, the limiting behavior of $R_{N,l,\beta=1}$ is indeed seen to be independent of the parity of $N$ (see e.g. [M, NW]).

---

[1] We use the standard notation $\operatorname{sgn} x = 1, 0, -1$ for $x > 0$, $x = 0$, $x < 0$, respectively.

Let $\{q_j(x)\}_{j\geq 0}$ be any sequence of polynomials of exact degree $j$, $q_j(x) = q_{j,j}x^j + \cdots$, $q_{j,j} \neq 0$. For $j = 0, 1, 2, \cdots$, set

$$\psi_{j,\beta}(x) = \begin{cases} q_j(x)w_1(x), & \beta = 1 \\ q_j(x)(w_4(x))^{1/2}, & \beta = 4. \end{cases} \tag{1.14}$$

Let $M_{N,1}$ denote the $N \times N$ matrix with entries

$$(M_{N,1})_{jk} = (\psi_{j,1}, \epsilon\psi_{k,1}), \qquad 0 \leq j,k \leq N-1, \tag{1.15}$$

and let $M_{N,4}$ denote the $2N \times 2N$ matrix with entries

$$(M_{N,4})_{jk} = (\psi_{j,4}, D\psi_{k,4}), \qquad 0 \leq j,k \leq 2N-1, \tag{1.16}$$

where again $(\cdot,\cdot)$ denotes the standard real inner product on $\mathbb{R}$. The matrices $M_{N,1}$ and $M_{N,4}$ are invertible (see e.g. [AvM, (4.17), (4.20)]). Let $\mu_{N,1}$, $\mu_{N,4}$ denote the inverses of $M_{N,1}$, $M_{N,4}$ respectively. With these notations we have [TW2] the following formulae for $S_{N,\beta}$ in (1.11), (1.12)

$$S_{N,1}(x,y) = -\sum_{j,k=0}^{N-1} \psi_{j,1}(x)\,(\mu_{N,1})_{jk}\,(\epsilon\psi_{k,1})(y) \tag{1.17}$$

$$S_{N,4}(x,y) = \sum_{j,k=0}^{2N-1} \psi'_{j,4}(x)\,(\mu_{N,4})_{jk}\,\psi_{k,4}(y). \tag{1.18}$$

An essential feature of the above formulae is that the polynomials $\{q_j\}$ are arbitrary and we are free to choose them conveniently to facilitate the asymptotic analysis of (1.11), (1.12) as $N \to \infty$. In view of the unitary case $\beta = 2$ described above, specific issues that must be addressed by the choice of the $q_j$'s, are the following:

(a) We need an analog of the Christoffel–Darboux formula to convert $S_{N,\beta}$ into a form similar to (1.7), which depends *only* on polynomials of high order as $N \to \infty$.

(b) By (a), the large $N$ behavior of (1.11), (1.12) becomes *purely* a question of the asymptotics of polynomials of high order. Thus we need to choose polynomials whose asymptotic behavior can be analyzed.

(c) Finally the inverses of the matrices $M_{N,1}$ and $M_{N,4}$ must be controlled as $N \to \infty$.

Property (c) is new in the cases $\beta = 1$ and 4, and does not arise in the unitary case $\beta = 2$. Following [W], we will choose the $q_j$'s to be orthogonal polynomials with respect to an appropriate weight. It turns out that this choice addresses all three issues (a), (b), (c) simultaneously.

Previously the study of the orthogonal and symplectic ensembles was often carried out via so-called *skew orthogonal polynomials (SOP's)* (see [M] for classical references and, e.g., [NW, AFNvM, AvM, FNH] for more recent work). Skew orthogonal polynomials are characterized by the property that (in our situation) $M_{N,\beta}$ has the form of a direct sum of scalar multiples of the matrix $\begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$. For many classical weights $w_\beta$, e.g., $w_\beta(x) = e^{-x^2}$, the corresponding SOP's can be computed explicitly [AFNvM, F]. The proof of universality, however, involves arbitrary weights $w_\beta$ for which no such explicit formulae for the SOP's are available. As noted above, the crucial ingredient in the proof of universality for the $\beta = 2$ ensemble is the asymptotic analysis of the OP's corresponding to an arbitrary weight $w_2$. For $\beta = 1, 4$, the problem of computing the asymptotics of SOP's raises challenging technical difficulties which have not yet been fully overcome (however see the work [E]).

In order to state our main result we need more notation. For any $m \in \mathbb{N}$ let $V(x)$ be a polynomial of degree $2m$

$$V(x) = \kappa_{2m} x^{2m} + \cdots , \qquad \kappa_{2m} > 0 \tag{1.19}$$

and let $w(x) \equiv w_{\beta=2}(x) = e^{-V(x)}$ as before. Let $p_j(x)$, $j \geq 0$, denote the OP's with respect to $w$, and set $\phi_j(x) \equiv p_j(x)(w(x))^{1/2}$, $j \geq 0$, as above. For $\beta = 1, 4$ set

$$V_\beta(x) \equiv \frac{1}{2} V(x) \tag{1.20}$$

and let $N$ be even. Then by (1.4), $w_4 = e^{-2V_4} = e^{-V}$ and $w_1 = e^{-V_1} = e^{-V/2}$. This ensures that for the choice $q_j = p_j$ in (1.14)

$$\psi_{j,\beta=1}(x) = \psi_{j,\beta=4}(x) = \phi_j(x), \tag{1.21}$$

which enables us in turn to handle $S_{N,1}$ and $S_{N/2,4}$ in (1.17), (1.18) simultaneously (see Remark 1.3 below). Henceforth and throughout the paper, $K_N$ denotes the Christoffel–Darboux kernel (1.5), (1.7) constructed out of these functions $\phi_j$.

The main result we state here (Theorem 1.1 below) concerns universality for the kernels $K_{N,\beta}$ in the bulk scaling limit. Universality for other standard statistical quantities also holds, for example for the $l$-point cluster functions and for the gap probability, see Corollaries 1.2 and 1.3 below.

As in the case $\beta = 2$ above, we scale so that the expected number of eigenvalues per unit interval is one. Note that formula (1.10) also holds for $\beta = 1$ and 4 and so $R_{N,l=1,\beta}(0)$ gives the density of the expected number of (simple) eigenvalues near zero. For $\beta = 1, 4$, in view of (1.13) and (1.11), (1.12)

$$\begin{aligned} R_{N,1,1}(r) &= S_{N,1}(r,r) \\ R_{N/2,1,4}(r) &= \frac{1}{2} S_{N/2,4}(r,r). \end{aligned} \tag{1.22}$$

It follows from (4.154), (4.155) below that as (even) $N \to \infty$ for $r$ in a compact set

$$\begin{aligned} S_{N,1}(r,r) &= K_N(0,0)(1 + O(N^{-1/2})) \\ S_{N/2,4}(r,r) &= K_N(0,0)(1 + O(N^{-1/2})) \end{aligned} \tag{1.23}$$

and hence

$$\begin{aligned} R_{N,1,1}(r) &= K_N(0,0)(1 + O(N^{-1/2})) \quad \text{and} \\ R_{N/2,1,4}(r) &= \frac{1}{2} K_N(0,0)(1 + O(N^{-1/2})) \end{aligned} \tag{1.24}$$

give the scaling in Theorem 1.1 below.

*Remark* 1.1. The asymptotics in (1.23), (1.24) and in many related situations (see e.g. Remark 1.2 below) remains true for $r = o(N^{1/(2m)})$, but we do not need this fact in what follows.

It turns out that the off-diagonal elements in $K_{N,\beta}$ scale differently as $N \to \infty$. On the other hand, the statistics of the ensembles are clearly invariant under the conjugation

$$K_{N,\beta} \mapsto K_{N,\beta}^{(\lambda)} \equiv \begin{pmatrix} \lambda^{-1} & 0 \\ 0 & \lambda \end{pmatrix} \cdot K_{N,\beta} \cdot \begin{pmatrix} \lambda & 0 \\ 0 & \lambda^{-1} \end{pmatrix} = \begin{pmatrix} (K_{N,\beta})_{11} & \lambda^{-2}(K_{N,\beta})_{12} \\ \lambda^{2}(K_{N,\beta})_{21} & (K_{N,\beta})_{22} \end{pmatrix}$$

for any scalar $\lambda = \lambda_{N,\beta}$. For example, this is obviously true for the cluster functions $T_{N,l,\beta}$ which have the form

(1.25)
$$T_{N,l,\beta}(y_1, \cdots, y_l) = \frac{1}{2l} \sum_{\sigma} \operatorname{tr}\Big( K_{N,\beta}(y_{\sigma_1}, y_{\sigma_2}) K_{N,\beta}(y_{\sigma_2}, y_{\sigma_3}) \cdots K_{N,\beta}(y_{\sigma_l}, y_{\sigma_1}) \Big)$$

where the sum is taken over all permutations of $\{1, \cdots, l\}$ (see [TW2, p. 816]), etc.

Set

$$\lambda_{N,1} = (R_{N,1,1}(r))^{1/2}, \qquad \lambda_{N,4} = (R_{N/2,1,4}(r))^{1/2}.$$

**Theorem 1.1.** *Let $\beta = 1$ or 4. For any $V(x)$ of degree $2m$ as in (1.19) define $V_\beta(x)$ and $w_\beta(x)$ as in (1.20), (1.4). Then for $r$, $\xi$, $\eta$ in a compact set, as (even) $N \to \infty$*

$$\begin{aligned} \mathcal{E}_{N,1} &\equiv \frac{1}{\lambda_{N,1}^2} K_{N,1}^{(\lambda_{N,1})}\left( r + \frac{\xi}{\lambda_{N,1}^2}, r + \frac{\eta}{\lambda_{N,1}^2} \right) - K^{(1)}(\xi,\eta) \to 0 \\ \mathcal{E}_{N,4} &\equiv \frac{1}{\lambda_{N,4}^2} K_{N/2,4}^{(\lambda_{N,4})}\left( r + \frac{\xi}{\lambda_{N,4}^2}, r + \frac{\eta}{\lambda_{N,4}^2} \right) - K^{(4)}(\xi,\eta) \to 0 \end{aligned} \tag{1.26}$$

*where*

$$\begin{aligned} K^{(1)}(\xi,\eta) &\equiv \begin{pmatrix} K_\infty(\xi-\eta) & \frac{\partial}{\partial\xi}\big(K_\infty(\xi-\eta)\big) \\ \int_0^{\xi-\eta} K_\infty(t)\,dt - \frac{1}{2}\operatorname{sgn}(\xi-\eta) & K_\infty(\eta-\xi) \end{pmatrix} \\ K^{(4)}(\xi,\eta) &\equiv \begin{pmatrix} K_\infty\big(2(\xi-\eta)\big) & \frac{\partial}{\partial\xi}\big(K_\infty(2(\xi-\eta))\big) \\ \int_0^{\xi-\eta} K_\infty(2t)\,dt & K_\infty\big(2(\eta-\xi)\big) \end{pmatrix}. \end{aligned} \tag{1.27}$$

*For $\beta = 1, 4$ and $r$, $\xi, \eta$ in a compact set, we have as $N \to \infty$*

$$\mathcal{E}_{N,\beta} = \begin{pmatrix} O\big(N^{-1/2}\big) & O\big(N^{-1/2}\big) \\ O\big(N^{-1/4}\big) & O\big(N^{-1/2}\big) \end{pmatrix}. \tag{1.28}$$

*Remark* 1.2. In the above theorem, we do not need to restrict $r$ to a compact set. The theorem remains true for $r = o(N^{1/(2m)})$.

*Remark* 1.3. The choice in (1.20) implies that $V_1(x) = V_4(x)$. Other authors make different choices. For example, in the Gaussian case [M], Mehta considers $V_4(x) = 2V_1(x)$. Our choice insures that we can use the same auxiliary OP's in (1.21) for $\beta = 1$ and $\beta = 4$. On the other hand, Mehta's choice ensures that the density of states for $P_{N,\beta}(x_1, \cdots, x_N)$ in (1.3) is the same for $\beta = 1$ and $\beta = 4$. Nevertheless we obtain precisely the same answer (1.26) in Theorem 1.1 as computed in [M] for $\beta = 1$ and $\beta = 4$ in the Gaussian case. Of course, this is due to the fact that in the end we use the same scaling as in [M], viz., the expected number of eigenvalues per unit interval is one.

From the formula for the cluster functions (1.25) we immediately have the following result.

**Corollary 1.2.** *Let $\beta = 1$ or $4$. Let $V$ be a polynomial of degree $2m$ and let $K^{(\beta)}$, $\beta = 1, 4$ be as in Theorem 1.1. Then for $\beta = 1$ and $l = 2, 3, \cdots$ we have, uniformly for $r$ and $\xi_1, \cdots, \xi_l$ in a compact set*

$$\begin{aligned}\lim_{N\to\infty} \frac{1}{(\lambda_{N,1}^2)^l} T_{N,l,1}\left(r + \frac{\xi_1}{\lambda_{N,1}^2}, \cdots, r + \frac{\xi_l}{\lambda_{N,1}^2}\right) \\ = \frac{1}{2l} \sum_{\sigma} \operatorname{tr}\left(K^{(1)}(\xi_{\sigma_1}, \xi_{\sigma_2}) K^{(1)}(\xi_{\sigma_2}, \xi_{\sigma_3}) \cdots K^{(1)}(\xi_{\sigma_l}, \xi_{\sigma_1})\right).\end{aligned} \tag{1.29}$$

*For $\beta = 4$, the same result is true provided we replace $T_{N,l,1} \to T_{N/2,l,4}$, $\lambda_{N,1} \to \lambda_{N,4}$, and $K^{(1)} \to K^{(4)}$.*

Together with some additional estimates (see Section 3), Theorem 1.1 also yields the following universality result for the gap probability. Recall that for a $2 \times 2$ block operator $A = (A_{ij})_{i,j=1,2}$ with $A_{11}, A_{22}$ in trace class and $A_{12}, A_{21}$ Hilbert–Schmidt, the regularized 2-determinant (see e.g. [Si]) is defined by $\det_2(I + A) \equiv \det((I + A)e^{-A})\, e^{\operatorname{tr}(A_{11}+A_{22})}$.

**Corollary 1.3.** *Let $\beta = 1$ or $4$. Let $V$ be a polynomial of degree $2m$ and let $K^{(\beta)}$, $\beta = 1, 4$ be as in Theorem 1.1. Fix $\theta > 0$ and $r$. Then*

$$\lim_{N\to\infty} \operatorname{Prob}\left\{\textit{no eigenvalues in } \left(r - \frac{\theta}{\lambda_{N,\beta}^2}, r + \frac{\theta}{\lambda_{N,\beta}^2}\right)\right\} = \sqrt{\det{}^{(\beta)}(I - K_\theta^{(\beta)})} \tag{1.30}$$

*where $K_\theta^{(\beta)}$ denotes the operator with kernel $K^{(\beta)}(\xi, \eta)$ acting in $L^2(-\theta, \theta)$. Here $\det^{(4)}(I - K_\theta^{(4)})$ is the regular determinant of the trace class operator $K_\theta^{(4)}$, but $\det^{(1)}(I - K_\theta^{(1)})$ is the regularized 2-determinant $\det_2(I - K_\theta^{(1)})$ defined above.*

*Remark* 1.4. The regularized 2-determinant is needed for $\beta = 1$ because the operator with kernel $\frac{1}{2}\operatorname{sgn}(\xi - \eta)$ is Hilbert–Schmidt but not trace class on $L^2(-\theta, \theta)$ (see proof of Corollary 1.3 in Section 3).

We complete this introduction with a description of Widom's result [W] which is basic for our approach in this paper. Widom's method applies to general weights $w_\beta$ with the property that $w'_\beta/w_\beta$ is a rational function. This property certainly holds for our weights as in (1.4), (1.2). Introduce the semi-infinite matrices

$$D_\infty \equiv ((D\phi_j, \phi_k))_{j,k\geq 0}, \qquad \epsilon_\infty \equiv ((\epsilon\phi_j, \phi_k))_{j,k\geq 0}. \tag{1.31}$$

It follows from [TW1, Section 6] that the matrix $D_\infty$ is banded with bandwidth $2n + 1$ where

$$n \equiv 2m - 1. \tag{1.32}$$

Thus $(D_\infty)_{jk} = 0$ if $|j - k| > n$. Next, let $N$ be greater than $n$, and introduce the following $N$-dependent $n$-column vectors

$$\begin{aligned}\Phi_1(x) &\equiv (\phi_{N-n}(x), \cdots, \phi_{N-1}(x))^T \\ \Phi_2(x) &\equiv (\phi_N(x), \cdots, \phi_{N+n-1}(x))^T \\ \epsilon\Phi_1(x) &\equiv (\epsilon\phi_{N-n}(x), \cdots, \epsilon\phi_{N-1}(x))^T \\ \epsilon\Phi_2(x) &\equiv (\epsilon\phi_N(x), \cdots, \epsilon\phi_{N+n-1}(x))^T\end{aligned} \tag{1.33}$$

and the following $2n \times 2n$ matrices consisting of four $n \times n$ blocks

$$B \equiv \begin{pmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{pmatrix} = ((\epsilon\phi_j, \phi_k))_{N-n \le j,k \le N+n-1}. \tag{1.34}$$

and

$$A \equiv \begin{pmatrix} 0 & A_{12} \\ A_{21} & 0 \end{pmatrix} = \begin{pmatrix} 0 & D_{12} \\ -D_{21} & 0 \end{pmatrix} \tag{1.35}$$

where $\begin{pmatrix} D_{11} & D_{12} \\ D_{21} & D_{22} \end{pmatrix} \equiv ((D\phi_j, \phi_k))_{N-n \le j,k \le N+n-1}$. Finally, set

$$C = \begin{pmatrix} C_{11} & C_{12} \\ C_{21} & C_{22} \end{pmatrix} \equiv \begin{pmatrix} I_n + (BA)_{11} & (BA)_{12} \\ (BA)_{21} & (BA)_{22} \end{pmatrix}.$$

Note that

$$C_{11} = I_n + B_{12}A_{21} = I_n - B_{12}D_{21}.$$

The main result in [W] is the following pair of formulae for $S_{N,1}$ and $S_{N/2,4}$

$$\begin{aligned} S_{N,1}(x,y) = K_N(x,y) - (\Phi_1(x)^T, 0^T) \cdot (AC(I_{2n} - BAC)^{-1})^T \\ \cdot (\epsilon\Phi_1(y)^T, \epsilon\Phi_2(y)^T)^T \end{aligned} \tag{1.36}$$

and

$$\begin{aligned} S_{N/2,4}(x,y) = K_N(x,y) + \Phi_2(x)^T \cdot D_{21} \cdot \epsilon\Phi_1(y) \\ + \Phi_2(x)^T \cdot D_{21} C_{11}^{-1} B_{11} D_{12} \cdot \epsilon\Phi_2(y). \end{aligned} \tag{1.37}$$

Observe that $S_{N,1}$ and $S_{N/2,4}$ are sums of the $\beta = 2$ kernel $K_N(x,y)$ together with correction terms that depend only on $\phi_{N+j}$ for $j \in \{-n, \cdots, n-1\}$. Together with the Christoffel–Darboux formula (1.7) we see immediately that the representations (1.36), (1.37) address issue (a) above. Moreover, the $\phi_{N+j}$'s are proportional to orthonormal polynomials $p_{N+j}$, and hence can be evaluated asymptotically as $N \to \infty$ by the methods in [DKMVZ2]. This addresses issue (b). Finally we see that issue (c), the question of the control of the inverses of the (large) matrices $M_{N,1}$, $M_{N,4}$, reduces, via (1.36), (1.37) to a question of controlling the inverses of matrices of *fixed* size $2n \times 2n$ and $n \times n$, respectively. It turns out that these fixed size matrices converge as $N \to \infty$. The proof that in both cases the limiting matrices are invertible for any $V$ of type (1.19), is not obvious, and constitutes a significant part of the proof of Theorem 1.1 in this paper.

In Section 2, we state a variety of auxiliary results which we use in the proof of Theorem 1.1. The results, some of which are of independent interest, are of two types:

(i) asymptotics of $(D\phi_{N+j}, \phi_{N+k})$ and $(\epsilon\phi_{N+j}, \phi_{N+k})$ for fixed $j,k$ as $N \to \infty$;

(ii) the equality of four specific determinants that arise in the analysis: This reduces the proof of the invertibility of the limiting matrices mentioned above to the proof of the invertibility of a single matrix $T_{m-1}$ (see (2.17) below).

In Section 3, we prove Theorem 1.1 (and hence Corollary 1.2) and Corollary 1.3 assuming the validity of the results in Section 2 (and 4). In Section 4, we prove the results of type (i) above. In Section 5, we prove the equality of the determinants in (ii). Finally, in Section 6 we prove that for any $V(x) = \kappa_{2m}x^{2m} + \cdots$, the determinant of $T_{m-1}$ is nonzero. In the course of the proof in Section 6, we need to estimate certain explicit integrals: A rigorous analysis of the error bounds in these estimates is given in an Appendix.

*Remark* 1.5. After the second version of this paper was posted on the arXiv, A. Boutet de Monvel informed us of the work of A. Stojanovic ([St1, St2, St3]) in which universality (in the bulk and also at the edge) is proved for $\beta = 1$ and $\beta = 4$ in the even, quartic (two-interval) case considered previously by Bleher and Its [BI] for $\beta = 2$. The method uses a variant of the formulae in [W] together with the asymptotics for OP's in [BI].

In [St2, Remark 2.4], an interesting connection is noted between the problem of controlling $\lim_{N\to\infty} \det C_{11}$ (in our notation) and the problem of estimating partition functions $Z_{N,\beta}$ (see (1.3)) as $N \to \infty$. Indeed from the formula $C_{11} = I_n - B_{12}D_{21}$ above and (2.18) below, we see that $\det C_{11} = \det \epsilon_N \det D_N$, where $\epsilon_N, D_N$ are the leading $N \times N$ sections of the matrices $\epsilon_\infty, D_\infty$, respectively. But $\det \epsilon_N$ and $\det D_N$ can be expressed in terms of partition functions (see [AvM]) and this leads directly to a formula for $\det C_{11}$ as a ratio of $Z_{N,\beta}$'s, $\beta = 1, 2, 4$. Thus one may try to use the methods of statistical mechanics to control $\lim_{N\to\infty} \det C_{11}$. The estimates in [J] show that the partition functions $Z_{N,\beta}$ have leading order asymptotics of the form $e^{a_\beta N^2(1+o(1))}$ as $N \to \infty$, and moreover, their combined contributions to $\det C_{11}$ cancel to this order. Thus, in order to show that $\lim_{N\to\infty} \det C_{11} \neq 0$, one needs higher order asymptotics for the $Z_{N,\beta}$'s. In one of our early attempts to prove universality, for $\beta = 1$ and 4 we indeed tried to derive such higher order asymptotics, but we were not successful. The statistical mechanical approach to prove $\lim_{N\to\infty} \det C_{11} \neq 0$, remains an unresolved, and intriguing, possibility.

**Notational remark:** Throughout this paper $c, C, C(m), c_1, c_2, \cdots$ refer to constants independent of $N, P$. More specifically, the symbols $c, C$ refer to generic constants, whose precise value may change from one inequality to another. The symbol $c_N$ however always refers to the $N$-dependent constant (2.3) below.

**Acknowledgments.** The authors would like to thank Harold Widom for alerting one of us (P.D.) to his paper [W], and for suggesting that formulae in this paper, together with the asymptotics for OP's in [DKMVZ2], might be used to prove universality for $\beta = 1$ and 4. We would also like to thank Alexei Borodin for suggesting a short proof of Lemma 4.1. The work was supported in part by an NSF grant DMS–0296084. In 2002–03 the second author was supported in full by a postdoctoral scholarship from the Swedish Foundation for International Cooperation in Research and Higher Education (STINT), Dnr. PD2001–128. The second author would like to thank the University of Pennsylvania where he had a postdoctoral appointment in 2001–02 and 2003–04. The second author would also like to thank the Courant Institute, New York University, where he has spent the year 2002–03 and parts of 2003–04, for hospitality and financial support.

## 2. Auxiliary results

In order to evaluate the asymptotic behavior of the correction terms in (1.36) and (1.37), one first needs to determine the asymptotics of $(D\phi_{N+j}, \phi_{N+k})$ and $(\epsilon\phi_{N+j}, \phi_{N+k})$ for fixed $j, k \in \{-n, \cdots, n-1\}$, as $N \to \infty$. Such asymptotics are given by the next two theorems which are proved in Section 4 below. Let $b_j$

be the coefficients in the three term recurrence relation [Sz] satisfied by the OP's $p_j$ corresponding to the weight $w(x) = e^{-V(x)}$ (and hence also by the functions $\phi_j = p_j w^{1/2}$)

$$xp_j(x) = b_{j-1}p_{j-1}(x) + a_j p_j(x) + b_j p_{j+1}(x), \qquad j \geq 0, \tag{2.1}$$

$(b_{-1} \equiv 0)$.

**Theorem 2.1.** *Let* $V(x) = \kappa_{2m}x^{2m} + \cdots$, $\kappa_{2m} > 0$, *and set* $n \equiv 2m-1$ *as before. Then the matrix* $D_\infty$ *has* $2n+1$ *bands and is asymptotic to the product of a diagonal matrix and a Toeplitz matrix. More precisely, for any fixed* $j, k \in \mathbb{Z}$, *as* $N \to \infty$

$$\begin{aligned}&(D\phi_{N+j}, \phi_{N+k})\\ &= m\kappa_{2m}b_{N+j}^n \cdot \begin{cases} 0, & j-k=0 \text{ or } |j-k| \geq n+1 \\ \operatorname{sgn}(j-k)\binom{n}{(n-|j-k|)/2} + o(1), & |j-k| = 1, 3, \cdots, n \\ o(1), & |j-k| = 2, 4, \cdots, n-1. \end{cases}\end{aligned} \tag{2.2}$$

To formulate the second theorem we need information on the equilibrium measure $d\mu_N^{(\mathrm{eq})}(x)$ (see e.g. [SaTo]) for OP's corresponding to the rescaled weight $e^{-NV_N(x)}$, $V_N = \frac{1}{N}V(c_N x + d_N)$, where $c_N$, $d_N$ are the so-called Mhaskar–Rakhmanov–Saff (MRS) numbers (see [MhSa, Ra]). For $V(x) = \kappa_{2m}x^{2m} + \kappa_{2m-1}x^{2m-1} + \cdots$ as in (1.19), we have [DKMVZ2] to any order $q$ as $N \to \infty$

$$c_N = \left(\frac{1}{\kappa_{2m}}\frac{(2m)!!}{m(2m-1)!!}\right)^{1/(2m)} N^{1/(2m)} + \sum_{j=0}^{q} c_{(j)}N^{-j/(2m)} + O(N^{-(q+1)/(2m)}) \tag{2.3}$$

and

$$d_N = -\frac{\kappa_{2m-1}}{2m\kappa_{2m}} + \sum_{j=1}^{q} d_{(j)}N^{-j/(2m)} + O(N^{-(q+1)/(2m)}). \tag{2.4}$$

As $N \to \infty$, the equilibrium measure is absolutely continuous with respect to Lebesgue measure, $d\mu_N^{(\mathrm{eq})}(x) = \psi_N^{(\mathrm{eq})}(x)\,dx$, and is supported on the (single) interval $[-1, 1]$,

$$\psi_N^{(\mathrm{eq})}(x) \equiv \psi_N(x) = \frac{1}{2\pi}|1-x^2|^{1/2}\chi_{[-1,1]}(x)\,h_N(x) \tag{2.5}$$

(see [DKMVZ2]) where $h_N(x)$ is a real polynomial of degree $2m-2$

$$h_N(x) = \sum_{k=0}^{2m-2} h_{N,k}x^k \tag{2.6}$$

and the coefficients $h_{N,k}$ can be expanded to any order in powers of $N^{-1/(2m)}$ as above. In particular, to any order $q = 1, 2, \cdots$, as $N \to \infty$, we find uniformly for $x$ in compact sets

$$h_N(x) = h(x) + \sum_{j=1}^{q} N^{-j/(2m)}\,h_{(j)}(x) + O(N^{-(q+1)/(2m)}) \tag{2.7}$$

where (see [DKMVZ2, (2.7)])

$$h(x) = \sum_{k=0}^{m-1} \beta_k x^{2k}, \qquad \beta_k \equiv 2\frac{(2m)(2(m-1))\cdots(2(m-k))}{(2m-1)(2(m-1)-1)\cdots(2(m-k)-1)}. \tag{2.8}$$

Also by [DKMVZ2, p. 1501, Remark 3]

$$h(1) = 4m, \qquad h(0) = \frac{4m}{2m-1}. \tag{2.9}$$

Note that $h(x)$ depends only on the degree $2m$ and is independent of the coefficients of $V$. Note also that $h(x) \geq h(0) > 0$ for all $x \in \mathbb{R}$.

**Theorem 2.2.** *Let $V(x) = \kappa_{2m}x^{2m} + \cdots$, $\kappa_{2m} > 0$ as before. Then, for any fixed $j, k \in \mathbb{Z}$, as $N \to \infty$*

$$\begin{aligned}&(\epsilon\phi_{N+j}, \phi_{N+k}) \\ &= c_{N+j}(N+j)^{-1} \cdot \begin{cases} \frac{(-1)^{N+j}}{2m} - I(j-k) + o(1), & j-k \text{ odd} \\ o(1), & j-k \text{ even}, j-k \neq 0 \\ 0, & j-k = 0 \end{cases}\end{aligned} \tag{2.10}$$

*where for $q = \pm 1, \pm 3, \cdots$ we define*

$$I(q) \equiv \frac{2}{\pi} \sin\frac{q\pi}{2} \int_{-1}^{1} \frac{\cos(q \arcsin x)}{h(x)} \frac{dx}{1-x^2}. \tag{2.11}$$

In the proof of (2.10) we compute a double integral over all the different asymptotic regions in $\mathbb{R}$ for the OP's $\{p_j\}$ obtained in [DKMVZ2] (see Subsection 4.2 et seq. below).

We see from the above, that in a fixed size neighborhood of the diagonal the matrices $D_\infty$ and $\epsilon_\infty$ do not depend asymptotically on lower order terms in $V(x)$. Next note that the coefficient in (2.1) satisfies [DKMVZ2, (2.11)]

$$b_N = \left(\frac{1}{2} + o(1)\right) c_N, \qquad a_N = d_N + o(1) = O(1), \qquad N \to \infty. \tag{2.12}$$

Combining our two theorems we can find an explicit expression for $C_{11}$, and also for all other matrices involved in the $\beta = 1$ and 4 correction terms in (1.36), (1.37). In particular, as (even) $N \to \infty$

(2.13)

$$\begin{aligned} B_{12} = \frac{c_N}{N} &\left[ \frac{1}{2m} \begin{pmatrix} -1 & 0 & -1 & \cdots & -1 \\ 0 & 1 & 0 & \cdots & 0 \\ \cdots & & & & \\ -1 & 0 & -1 & \cdots & -1 \end{pmatrix} \right. \\ &\left. + \begin{pmatrix} I(n) & 0 & I(n+2) & \cdots & I(2n-1) \\ 0 & I(n) & 0 & \cdots & 0 \\ \cdots & & & & \\ I(1) & 0 & I(3) & \cdots & I(n) \end{pmatrix} + o(1) \right] \\ A_{21} = -D_{21} = -\frac{m\kappa_{2m}}{2^{2m-1}} c_N^{2m-1} &\left[ \begin{pmatrix} \binom{n}{0} & 0 & \binom{n}{1} & \cdots & \binom{n}{(n-1)/2} \\ 0 & 1 & 0 & \cdots & 0 \\ \cdots & & & & \\ 0 & 0 & 0 & \cdots & 1 \end{pmatrix} + o(1) \right]. \end{aligned}$$

The second and the third matrices are Toeplitz.

By (2.3) the asymptotic orders of $B_{12}$, $D_{21}$ are exactly opposite. The same is true for all the blocks $B_{ij}$ and $D_{ij}$, $i, j = 1, 2$, and so whenever we have a product of elements from $B$ and $A$ in (1.36), (1.37), the product is asymptotically constant. Also

$$\begin{aligned}\lim_{N\to\infty}\frac{c_N}{N}\frac{m\kappa_{2m}}{2^{2m-1}}\,c_N^{2m-1} &= \frac{(2m)!!}{2^{2m-1}(2m-1)!!}\\ &= \frac{2^m m!}{2^{2m-1}1\cdot 3\cdot\ldots\cdot(2m-1)}\,\frac{1\cdot 2\cdot\ldots\cdot m}{1\cdot 2\cdot\ldots\cdot m}=\frac{2(m!)^2}{(2m)!}\end{aligned}\tag{2.14}$$

so that in $BA$ the coefficient $\kappa_{2m}$ cancels out, and as $N\to\infty$, $BA$ converges to a constant matrix that depends only on $m$.

Recall that in each of the cases $\beta=1$ and 4, in order to compute the correction terms in (1.36), (1.37) we need to invert a certain matrix. Now we know that the $\beta=4$ matrix to be inverted, $I_n+B_{12}A_{21}$, converges as $N\to\infty$. In the $\beta=1$ case the matrix to be inverted is $I_{2n}+BAC$, which also converges as $N\to\infty$. It is not a priori clear whether the two limiting matrices are related. However our structure Theorems 2.1 and 2.2, and also Theorem 2.5 below, imply the following. Let $R\equiv R_n$ denote the $n\times n$ matrix with all zero entries apart from ones on the anti-diagonal (thus $R_{i,j}=1$ if $j=n-i+1$, $1\le i\le n$, and $R_{i,j}=0$ otherwise).

**Theorem 2.3.** *As $N\to\infty$, $N$ even, we have $(BA)_{22}=-R(BA)_{11}R+o(1)$. Also, quite remarkably,*

$$BAC=\begin{pmatrix}0 & 0\\ (BA)_{21}+o(1) & (BA)_{22}+o(1)\end{pmatrix},\tag{2.15}$$

*so that in particular $BAC$ and $BA$ asymptotically have the same two lower blocks.*

It is interesting that the 11 and the 21 entries of $BAC$ are identically zero, not just $o(1)$.

Now we see that inverting $I_{2n}+BAC$ boils down to inverting

$$I_n-(-R(BA)_{11}R)+o(1)=R(I_n+B_{12}A_{21})R+o(1)$$

so that the determinants of the limiting matrices in the $\beta=1$ and 4 cases that are to be inverted are the same. Thus in both cases we have to check that $\det(I_n+B_{12}A_{21})=\text{const}+o(1)$, $\text{const}\neq 0$.

Next, we observe that the zero pattern in $B_{12}$ and $A_{21}$ in (2.13) implies that

$$\det(I+B_{12}A_{21})=(\det T'_m+o(1))\cdot(\det T_{m-1}+o(1))$$

as $N\to\infty$, $N$ even, where

$$\begin{aligned}T'_m\equiv I_m-\frac{2(m!)^2}{(2m)!}&\left[\frac{1}{2m}\begin{pmatrix}-1 & -1 & \cdots & -1\\ -1 & -1 & \cdots & -1\\ \cdots & & & \\ -1 & -1 & \cdots & -1\end{pmatrix}\right.\\ &\left.\quad+\begin{pmatrix}I(n) & I(n+2) & \cdots & I(2n-1)\\ I(n-2) & I(n) & \cdots & I(2n-3)\\ \cdots & & & \\ I(1) & I(3) & \cdots & I(n)\end{pmatrix}\right]\\ &\times\begin{pmatrix}\binom{n}{0} & \binom{n}{1} & \cdots & \binom{n}{(n-1)/2}\\ 0 & \binom{n}{0} & \cdots & \binom{n}{(n-3)/2}\\ \cdots & & & \\ 0 & 0 & \cdots & 1\end{pmatrix}\end{aligned}\tag{2.16}$$

and

(2.17)

$$T_{m-1} \equiv I_{m-1} - \frac{2(m!)^2}{(2m)!}\left[\frac{1}{2m}\begin{pmatrix} 1 & 1 & \cdots & 1 \\ 1 & 1 & \cdots & 1 \\ \cdots & & & \\ 1 & 1 & \cdots & 1 \end{pmatrix} + \begin{pmatrix} I(n) & I(n+2) & \cdots & I(2n-3) \\ I(n-2) & I(n) & \cdots & I(2n-5) \\ \cdots & & & \\ I(3) & I(5) & \cdots & I(n) \end{pmatrix}\right] \times \begin{pmatrix} \binom{n}{0} & \binom{n}{1} & \cdots & \binom{n}{(n-3)/2} \\ 0 & \binom{n}{0} & \cdots & \binom{n}{(n-5)/2} \\ \cdots & & & \\ 0 & 0 & \cdots & 1 \end{pmatrix}.$$

Quite remarkably, we also have

**Theorem 2.4.** *For any $m \in \mathbb{N}$, $m \geq 2$,*

$$\det T'_m = \det T_{m-1}.$$

Thus it suffices to show that the *single* determinant, $\det T_{m-1}$, is nonzero (see Theorem 2.6 below). The proofs of Theorem 2.3 and 2.4 rely on the following identities. Let $D_N$, $\epsilon_N$ denote the $N \times N$ matrices formed by the first $N$ rows and columns of $D_\infty$, $\epsilon_\infty$, respectively. Recall that by (1.32) $D_\infty$ has bandwidth $2n+1$. Let $I_\infty$ denote the semi-infinite identity matrix.

**Theorem 2.5.** *(i) The semi-infinite matrices satisfy*

$$D_\infty \epsilon_\infty = \epsilon_\infty D_\infty = I_\infty$$

*(the products are well-defined since $D_\infty$ is banded);*

*(ii) the sections for $N > n$ satisfy*

$$\epsilon_N D_N = \begin{pmatrix} I_{N-n} & -\epsilon^*_{N-n,n} D_{21} \\ & \\ 0 & I_n - B_{12} D_{21} \end{pmatrix} \tag{2.18}$$

*where $\epsilon^*_{N-n,n}$ is the $(N-n) \times n$ matrix formed by the rows $0, \cdots, N-n-1$ and columns $N, \cdots, N+n-1$ of the matrix $\epsilon_\infty$.*

*Remark* 2.1. Assuming $N$ is even and denoting $W_N \equiv \epsilon_N D_N$ we see from (2.18) that

$$\epsilon_N^{-1} = D_N W_N^{-1}, \qquad D_N^{-1} = W_N^{-1} \epsilon_N. \tag{2.19}$$

Thus the question of the invertibility of both $\epsilon_N$ and $D_N$, reduces to checking that

$$\det(I_n - B_{12} D_{21}) = \det(I + B_{12} A_{21}) \neq 0.$$

On the other hand, one can show that the corrections in (1.36), (1.37) can be rewritten in terms of the lower-right $n \times n$ corner of the matrices $\epsilon_N^{-1}$ and $D_N^{-1}$, and so (2.18) and (2.19) explain why the same determinant should be checked to be nonzero for both the $\beta = 1$ and 4 cases. Also using the fact that

$$W_N^{-1} = \begin{pmatrix} I_{N-n} & * \\ & \\ 0 & (I_n - B_{12}D_{21})^{-1} \end{pmatrix}$$

along with (2.19), the skew symmetricity of $\epsilon_N$, and (2.18), we can show that $\epsilon_N^{-1}$ coincides with $D_N$ everywhere apart from the above mentioned lower-right $n \times n$ corner.

Next we state our main technical result that ensures the invertibility of the matrices in (1.36), (1.37) in both the $\beta = 1$ and 4 cases.

**Theorem 2.6.** *For any $m \in \mathbb{N}$, $m \geq 2$,*

$$\det T_{m-1} \neq 0.$$

*Remark* 2.2. From numerical computations for $m = 2, \cdots, 14$, it seems that $\det T_{m-1}$ approaches the value $\frac{1}{\sqrt{2}}$ as $m$ grows, but we have not been able to use our methods to establish such a result.

*Remark* 2.3. The basic idea in proving Theorem 2.6 is to show that for an appropriate norm $\|\cdot\|$, $\|T_{m-1} - I_{m-1}\| < 1$. Rather than analyzing the quantities $I(q)$ in (2.11), (2.17) directly, we consider

$$\tilde{I}(q) \equiv mI(q) - \frac{1}{2} = \frac{2}{\pi} \sin\frac{q\pi}{2} \int_{-\pi/2}^{\pi/2} \cos(q\theta)\, y_m(\theta)\, d\theta, \qquad q = 3, 5, \cdots 4m-5 \tag{2.20}$$

where

$$y_m(\theta) \equiv \frac{m}{\cos\theta}\left(\frac{1}{h(\sin\theta)} - \frac{1}{4m} - \frac{\cos^2\theta}{2}\right) \tag{2.21}$$

(cf. (6.3) et seq. below). It turns out that $h(x)$ is (essentially) a ${}_2F_1$ hypergeometric function (see (6.11)) and satisfies a *first* order differential equation (see (6.8)). This in turn implies that $y_m$ satisfies a Riccati equation

$$y_m' = \frac{4}{\sin\theta}\left(y_m + \frac{2m+1}{4}\cos\theta\right)\left(y_m + \frac{1}{2\cos\theta}\right), \tag{2.22}$$

where we see that the right-hand side conveniently splits into a product with explicit factors. Equation (2.22) plays a key role in proving that the magnitude of $y_m$ is $O(\sqrt{m})$, a crucial fact in the proof of Theorem 2.6. In estimating the magnitude of the quantities $\tilde{I}(q)$, it is clear from (2.20) that we are in a double-scaling situation: if $q$ is large compared to $m$, then $\tilde{I}(q)$ is small by virtue of the usual decay of Fourier coefficients of real analytic functions. On the other hand if $q$ is small with respect to $m$, the asymptotic behavior of $y_m$ dominates. Both regimes must be analyzed separately. These two effects are comparable when $q \sim \sqrt{m}$.

## 3. Proofs of Theorem 1.1 and Corollary 1.3

We only consider the case $\beta = 1$. The proofs for $\beta = 4$ are similar and are left to the reader.

We need the following result which is based on the Plancherel–Rotach type asymptotics for OP's in [DKMVZ2] (see Section 4 for the proof).

**Theorem 3.1.** *Let* $V(x) = \kappa_{2m}x^{2m} + \cdots$ *as before. Let* $a = o(c_N)$ *as* $N \to \infty$. *Then, as* $N \to \infty$

$$\|\phi_N\|_{L^\infty((-a,a))} = O(c_N^{-1/2}), \qquad \|\epsilon\phi_N\|_{L^\infty(\mathbb{R})} = O(c_N^{1/2}\cdot N^{-1/2}). \tag{3.1}$$

*Also* $\|\phi_N\|_{L^\infty(\mathbb{R})} = O(c_N^{-1/2}N^{1/6})$.

3.1. **Proof of Theorem 1.1.** Set $q_N \equiv \lambda_{N,1}^2 = R_{N,1,1}(r)$. Observe from (1.24) and (4.152), (4.154) that $q_N$ satisfies (4.138) with $\alpha = 1/2$. Thus $\alpha^*$ in (4.168) is 1/4. In view of (1.11), (1.36), we have

(3.2)

$$\begin{aligned}
&\frac{1}{q_N}K_{N,1}^{(\lambda_{N,1})}\left(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\right)\\
&= \begin{pmatrix} \frac{1}{q_N}S_{N,1}\big(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\big) & \frac{1}{q_N^2}(S_{N,1}D)\big(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\big) \\ (\epsilon S_{N,1})\big(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\big) - \frac{1}{2}\operatorname{sgn}(\xi-\eta) & \frac{1}{q_N}S_{N,1}\big(r+\frac{\eta}{q_N}, r+\frac{\xi}{q_N}\big) \end{pmatrix}
\end{aligned}$$

where

(3.3)

$$\begin{aligned}
\frac{1}{q_N}S_{N,1}\left(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\right) =& \frac{1}{q_N}K_N\left(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\right)\\
&- \frac{1}{q_N}(\Phi_1(r+\xi/q_N)^T, 0^T)\cdot(AC(I_{2n}-BAC)^{-1})^T\\
&\cdot(\epsilon\Phi_1(r+\eta/q_N)^T, \epsilon\Phi_2(r+\eta/q_N)^T)^T
\end{aligned}$$

The convergence of the derivatives and integrals of the Christoffel–Darboux kernel $\frac{1}{q_N}K_N\big(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\big)$ with uniform error estimates for $\xi, \eta, r \in [-2L_0, 2L_0]$, for a fixed $L_0 < \infty$, is established in Subsection 4.3, see (4.150), (4.157), (4.158) and (4.171) below.

We now consider the correction terms. By the results of Section 2 (see, in particular, (2.13) et seq.), $BA$ and $C$ converge as $N \to \infty$. Moreover $I_{2n} - \lim_{N\to\infty} BAC$ is invertible and hence $(I_{2n} - BAC)^{-1} = O(1)$ as $N \to \infty$. From (2.2) and (2.3), $A \sim b_N^n \sim c_N^n = O(N^{1-1/(2m)})$ and from (3.1)

$$\max_{|\xi|,|r|\le 2L_0}|\Phi_1(r+\xi/q_N)| = O(c_N^{-1/2}), \qquad \|\epsilon\Phi_2\|_{L^\infty(\mathbb{R})} = O(c_N^{1/2}N^{-1/2}). \tag{3.4}$$

Thus the correction term for the 11 and 22 entries in (3.2) is bounded by

$$\text{const}\cdot\frac{1}{N^{1-1/(2m)}}\cdot c_N^{-1/2}\cdot N^{1-1/(2m)}\cdot(c_N^{1/2}N^{-1/2}) = O\left(\frac{1}{\sqrt{N}}\right) \tag{3.5}$$

as $N \to \infty$, uniformly for $\xi, \eta, r \in [-2L_0, 2L_0]$.

The correction term in the 12 entry of (3.2) has the form

$$\begin{aligned}
&\frac{1}{q_N^2}(\Phi_1(r+\xi/q_N)^T, 0^T)\cdot(AC(I_{2n}-BAC)^{-1})^T\\
&\qquad\qquad\cdot(\Phi_1(r+\eta/q_N)^T, \Phi_2(r+\eta/q_N)^T)^T
\end{aligned}$$

and is bounded by

$$\text{const}\cdot\frac{1}{(N^{1-1/(2m)})^2}\cdot c_N^{-1/2}\cdot N^{1-1/(2m)}\cdot c_N^{-1/2} = O\left(\frac{1}{N}\right) \tag{3.6}$$

as $N \to \infty$, uniformly for $\xi, \eta, r \in [-2L_0, 2L_0]$.

Finally, to analyze the 21 entry of (3.2) (see (1.36)), we use the following observation. By (1.17), $(\epsilon S_{N,1})(x,y)$ is skew symmetric. Thus

$$(\epsilon S_{N,1})(x,y) = (\epsilon S_{N,1})(x,y) - (\epsilon S_{N,1})(y,y) = -\int_x^y S_{N,1}(t,y)\,dt. \tag{3.7}$$

After setting $x = r + \xi/q_N$, $y = r + \eta/q_N$ we see that the correction term has the form

$$\begin{aligned} -\Big(\Big(-\int_{r+\xi/q_N}^{r+\eta/q_N} \Phi_1(t)\,dt\Big)^T, 0^T\Big) &\cdot (AC(I_{2n} - BAC)^{-1})^T \\ &\cdot \big(\epsilon\Phi_1(r+\eta/q_N)^T, \epsilon\Phi_2(r+\eta/q_N)^T\big)^T. \end{aligned} \tag{3.8}$$

The integration is over a subset of the region $I_3$ (see (4.6) below). Hence as in (3.5), but now using (4.18) below, (3.8) is bounded by

$$\text{const} \cdot c_N^{1/2} N^{-1} \cdot N^{1-1/(2m)} \cdot c_N^{1/2} N^{-1/2} = O(N^{-1/2}) \tag{3.9}$$

as $N \to \infty$. We compute finally the contribution of the Christoffel–Darboux to (3.7). We have

$$-\int_x^y K_N(t,y)\,dt = (\epsilon K_N)(x,y) - (\epsilon K_N)(y,y) \tag{3.10}$$

where $x = r + \xi/q_N$, $y = r + \eta/q_N$. By (4.159), (4.171) below with $\alpha = 1/2, \alpha^* = 1/4$, (3.10) equals as $N \to \infty$

$$\Big(\int_0^{\xi-\eta} K_\infty(t)\,dt - \int_0^{\eta-\eta} K_\infty(t)\,dt\Big) + O(N^{-1/4}) = \int_0^{\xi-\eta} K_\infty(t)\,dt + O(N^{-1/4})$$

where the estimates are uniform for $r, \xi, \eta \in [-2L_0, 2L_0]$.

Assembling the above estimates we see that we have proved Theorem 1.1 in the case $\beta = 1$ with error term

$$\mathcal{E}_{N,1} = \begin{pmatrix} O(N^{-1/2}) & O(N^{-1/2}) \\ O(N^{-1/4}) & O(N^{-1/2}) \end{pmatrix} \tag{3.11}$$

as $N \to \infty$, uniformly for $\xi, \eta, r \in [-2L_0, 2L_0]$.

### 3.2. **Proof of Corollary 1.3.**

In [TW2] the authors prove that if $\chi_B$ denotes the characteristic function of a bounded Borel set $B$ then for finite $N$ in the case $\beta = 1$

$$\text{Prob}\{\text{ no eigenvalues in B }\} = \sqrt{\det\left((1 - K_{N,1}\chi_B)\begin{pmatrix} 1 & 0 \\ -\epsilon\chi_B & 1 \end{pmatrix}\right)}$$

where again $\epsilon$ denotes the operator with kernel $\frac{1}{2}\operatorname{sgn}(x-y)$. Now the product of the operators above is of the type identity plus trace class, but the individual operators are not. This is because $\epsilon$ is Hilbert–Schmidt but not trace class. However, as indicated in Section 1, we can use regularized 2-determinants. One easily sees, moreover, that $\det_2 \begin{pmatrix} 1 & 0 \\ -\epsilon\chi_B & 1 \end{pmatrix} = 1$. Thus

$$\det\Big((1-K_{N,1}\chi_B)\begin{pmatrix} 1 & 0 \\ -\epsilon\chi_B & 1 \end{pmatrix}\Big) = \det_2\Big((1-K_{N,1}\chi_B)\begin{pmatrix} 1 & 0 \\ -\epsilon\chi_B & 1 \end{pmatrix}\Big)$$

$$= \det_2(1-K_{N,1}\chi_B)\det_2\begin{pmatrix} 1 & 0 \\ -\epsilon\chi_B & 1 \end{pmatrix} = \det_2(1-K_{N,1}\chi_B).$$

For $\beta = 4$ such regularization issues do not arise. For $\beta = 1$ (cf. (1.30))

$$\begin{aligned} &\operatorname{Prob}\Big\{\text{ no eigenvalues in } \Big(r - \frac{\theta}{\lambda_{N,1}^2}, r + \frac{\theta}{\lambda_{N,1}^2}\Big)\Big\} \\ &\qquad = \sqrt{\det{}^{(1)}\Big(1 - K_{N,1}\chi_{\left(r-\frac{\theta}{\lambda_{N,1}^2}, r+\frac{\theta}{\lambda_{N,1}^2}\right)}\Big)}. \end{aligned} \tag{3.12}$$

For $\beta = 4$, the same formula is true provided we replace $\lambda_{N,1} \to \lambda_{N,4}$, $\det^{(1)} \to \det^{(4)}$, $K_{N,1} \to K_{N/2,4}$ (cf. (1.29)). In order to prove Corollary 1.3 for $\beta = 1$ it is clearly sufficient to prove that the entries of $\frac{1}{q_N}K_{N,1}^{(\lambda_{N,1})}\big(r + \frac{\xi}{q_N}, r + \frac{\eta}{q_N}\big)$, $q_N \equiv \lambda_{N,1}^2$, apart from the term $\frac{1}{2}\operatorname{sgn}(\xi-\eta)$, converge to the corresponding entries in $K^{(1)}(\xi,\eta)$ (cf. (3.2), (3.3), (1.27)) in trace norm for $L^2(-\theta,\theta)$. The case $\beta = 4$ is similar, mutatis mutandis, and is left to the reader.

We consider $r, \xi, \eta$ in the compact set $[-2L_0, 2L_0]$ with $L_0 \equiv \theta$. We have

$$\frac{1}{q_N}S_{N,1}\big(r + \frac{\xi}{q_N}, r + \frac{\eta}{q_N}\big) = \frac{1}{q_N}K_N\big(r + \frac{\xi}{q_N}, r + \frac{\eta}{q_N}\big) + \text{ correction term.}$$

By (3.3), the correction term corresponds to a finite rank operator, and hence its trace norm is bounded by the $L^2(-\theta,\theta)$ norms of the vectors $(\Phi_1(r+\xi/q_N)^T, 0^T)$, $(\epsilon\Phi_1(r+\eta/q_N)^T, \epsilon\Phi_2(r+\eta/q_N)^T)^T$ which in turn may be estimated by the bounds in (3.4). We conclude that the trace norm of the correction term decays as $N^{-1/2}$ uniformly for $|r| \leq 2\theta$.

The fact that $\frac{1}{q_N}K_N\big(r + \frac{\xi}{q_N}, r + \frac{\eta}{q_N}\big) \to K_\infty(\xi - \eta)$ in trace norm was proved in the case $V(x) = x^{2m}$ in [D]. We now give a proof for general $V(x) = \kappa_{2m}x^{2m} + \cdots$, which in addition also applies to the other entries of $\frac{1}{q_N}K_{N,1}^{(\lambda_{N,1})}$. Note first that it is sufficient to prove that $\chi\frac{1}{q_N}K_N\big(r + \frac{\cdot}{q_N}, r + \frac{\cdot\cdot}{q_N}\big)\chi$ converges to $\chi K_\infty(\cdot - \cdot\cdot)\chi$ in trace norm for any $C_0^\infty(-2\theta, 2\theta)$ function $\chi$ with $\chi(\xi) = 1$ for $\xi \in (-\theta,\theta)$. Let $P \equiv iD \equiv i\frac{\partial}{\partial\xi}$ denote the self-adjoint operator of differentiation with periodic boundary conditions on $[-2\theta, 2\theta]$. Then $D + I = \frac{1}{i}P + I$ is an invertible operator with eigenvalues $\gamma_k = \frac{\pi k}{i2\theta} + 1$, $k \in \mathbb{Z}$. In particular $\frac{1}{D+I}$ is Hilbert–Schmidt on $L^2(-2\theta, 2\theta)$. Clearly $\chi\frac{1}{q_N}K_N\chi$ maps $L^2(-2\theta, 2\theta)$ into the domain of $P$, $\operatorname{Dom} P = \{f \in L^2(-2\theta,2\theta) : f' \in L^2(-2\theta, 2\theta),\ f(-2\theta) = f(2\theta)\}$ and so

$$\begin{aligned} \chi\frac{1}{q_N}K_N\chi &= \frac{1}{D+I}(D+I)\chi\frac{1}{q_N}K_N\chi \\ &= \frac{1}{D+I}\chi\frac{1}{q_N}K_N\chi + \frac{1}{D+I}\chi'\frac{1}{q_N}K_N\chi + \frac{1}{D+I}\chi\Big(\frac{\partial}{\partial\xi}\frac{1}{q_N}K_N\Big)\chi \end{aligned} \tag{3.13}$$

But by (4.150) below, $\frac{1}{q_N}K_N\big(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\big)$ converges to $K_\infty(\xi-\eta)$ uniformly for $|r|, |\xi|, |\eta| \le 2\theta$ and hence $\chi\frac{1}{q_N}K_N\chi \to \chi K_\infty\chi$ in Hilbert–Schmidt norm. But then $\frac{1}{D+I}\chi\frac{1}{q_N}K_N\chi \to \frac{1}{D+I}\chi K_\infty\chi$ in trace norm. The same is clearly true for the second term in (3.13), and also for the third term (take $j=1$, $k=0$ in (4.150) below). This shows that $\frac{1}{q_N}K_N \to K_\infty$ in trace norm and completes the proof that the 11 entry of $\frac{1}{q_N}K_N\big(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\big) \to K_\infty(\xi-\eta)$ converges in trace norm to $(K^{(1)})_{11}$, as desired. Clearly the same is true for the 22 entry. Similar considerations using (4.157) in place of (4.150) prove the corresponding result for the 12 entry. Finally, for the 21 entry we observe that $Y_N(\xi,\eta)$ in (4.171) is a sum of two terms $A_N(\xi,\eta)+B_N(\eta)$. The first term can be controlled in trace norm using the operator $P=iD$ as above, together with the estimate (4.171)(a). On the other hand $B_N(\eta)$ corresponds to a rank one operator in $L^2(-\theta,\theta)$ and hence its trace norm goes to zero as $N\to\infty$ by (4.171)(b). Finally, the correction term in the 21 entry can be controlled as above. This completes the proof of Corollary 1.3.

## 4. ASYMPTOTICS OF INTEGRATED OP'S, PROOFS OF THEOREM 2.1 AND THEOREMS 2.2, 3.1

**Notational remark:** Throughout this Section, the notation $N, P \dot{\to} \infty$ means that $N, P\to\infty$, $N \ge P$, $N-P$ is fixed (and finite).

4.1. **Proof of Theorem 2.1.** Let $V(x) = \kappa_{2m}x^{2m}+\cdots$, $\{p_j\}_{j\ge0}$, $\{\phi_j\}_{j\ge0}$ be as before. The following observation is due to A. Borodin.

**Lemma 4.1.** *For any* $j,k\ge 0$

$$(D\phi_j,\phi_k) = \frac{1}{2}\operatorname{sgn}(j-k)\,(V'\phi_j,\phi_k). \tag{4.1}$$

*Proof.* For $j\le k$ we have

$$\begin{aligned}(D\phi_j,\phi_k) &= \int \big(p_j(x)e^{-V(x)/2}\big)' p_k e^{-V(x)/2}\,dx \\ &= \int p_j'(x)p_k(x)e^{-V(x)}\,dx - \frac{1}{2}(V'\phi_j,\phi_k)\end{aligned}$$

and the last integral vanishes since $p_k$ is orthogonal to all polynomials of degree $< k$, and we have assumed that $j\le k$. Observe that $(V'\phi_k,\phi_k)=0$ as it should be. The case $j>k$ follows by skew symmetry. □

*Remark* 4.1. Clearly Lemma 4.1 is true for any differentiable $V(x)$ such that

$$\int |x|^q\,(1+|V'(x)|)\,e^{-V(x)}\,dx<\infty, \qquad q=0,1,2,\cdots.$$

As $V'(x)$ is a polynomial of degree $n=2m-1$, it follows immediately from (4.1) that $((D\phi_j,\phi_k))_{j,k\ge0}$ is a banded batrix, $(D\phi_j,\phi_k)=0$ for $|j-k|>n$.

For the proof of Theorem 2.1 we need to evaluate the leading asymptotics of $(V'\phi_{N+j},\phi_{N+k})$, $j,k$ fixed, as $N\to\infty$. This clearly reduces to computing the large $N$ asymptotics of $(x^q\phi_{N+j},\phi_{N+k})$, $j,k$ fixed, $0\le q\le n$. In order to compute such asymptotics we will use the following relations for the recurrence coefficients in (2.1)

$$b_N/b_{N-1}\to 1, \qquad \text{and} \qquad a_N = o(b_N), \qquad \text{as } N\to\infty. \tag{4.2}$$

These relations certainly hold for our potential $V(x)=\kappa_{2m}x^{2m}+\cdots$, see (2.12), (2.3), (2.4).

**Lemma 4.2.** *For any fixed $q\in\mathbb{N}$, as $N\to\infty$*

$$\begin{aligned} x^q\phi_N(x)=&b_N^q\cdot\Bigg[\sum_{l=0}^{q}\binom{q}{l}(1+o(1))\cdot\phi_{N-q+2l}(x)\\ &+\sum_{l=1}^{q}o(1)\cdot\phi_{N-q+(2l-1)}(x)\Bigg]\end{aligned}\tag{4.3}$$

*where the notation $o(1)$ indicates terms which are independent of $x$ and uniform for $0\le l\le q$. (The $o(1)$ terms above can be replaced with $O(N^{-1/(2m)})$.)*

*Proof.* By (4.2) the three term recurrence relation (2.1) takes the form as $N\to\infty$,

$$\begin{aligned} x\phi_N(x)=b_N&\left[\phi_{N+1}(x)+(1+o(1))\,\phi_{N-1}(x)\right.\\ &\left.+\,o(1)\,\phi_N(x)\right]\end{aligned}\tag{4.4}$$

which coincides with (4.3) for $q=1$. A simple induction on $q$ using (4.4) and Pascal's triangle, now gives the result. □

We are now in a position to prove Theorem 2.1. Write $V(x)=\sum_{q=0}^{2m}v_qx^q$, $v_{2m}=\kappa_{2m}$. Then by Lemma 4.1, Lemma 4.2 and the orthogonality of the $\phi_j$'s, (4.5)

$$\begin{aligned}(D\phi_{N+j},\phi_{N+k})=&\frac{1}{2}\operatorname{sgn}(j-k)\sum_{q=1}^{2m}qv_q(x^{q-1}\phi_{N+j},\phi_{N+k})\\ =&\frac{1}{2}\operatorname{sgn}(j-k)\sum_{q=1}^{2m}qv_qb_N^{q-1}\Bigg[\sum_{l=0}^{q-1}\binom{q-1}{l}(1+o(1))\,\delta_{2l,k-j+q-1}\\ &+\sum_{l=1}^{q-1}o(1)\,\delta_{2l,k-j+q}\Bigg],\end{aligned}$$

as $N\to\infty$. Observe that for $|j-k|\le n$ and $q=2m=n+1$, $k-j+q-1=k-j+n$ lies between 0 and $2n$. Hence if $|j-k|\le n$ and $j-k$ is odd, we see that the square bracket in (4.5) for $q=2m$ gives rise to a leading contribution $\binom{n}{(n+k-j)/2}(1+o(1))$. However if $|j-k|\le n$ and $j-k$ is even, the contribution is $o(1)$. But by (2.12), (2.3), $b_N\to\infty$, and hence the terms corresponding to $q=1,2,\cdots,2m-1$ in (4.5) contribute to lower order. This completes the proof of Theorem 2.1.

## 4.2. **Integrating the Plancherel–Rotach type asymptotics for OP's and the proof of Theorems 3.1 and 2.2.**

4.2.1. *Auxiliary estimates on the integrated Plancherel–Rotach asymptotics.* For the convenience of the reader, we recall relevant results from [DKMVZ2]. Assume that $V(x)=\kappa_{2m}x^{2m}+\cdots$, $m\in\mathbb{N}$, $\kappa_{2m}>0$. Let $p_N$ be the $N$th OP on $\mathbb{R}$ with respect to the weight $e^{-V(x)}$ and set $\phi_N(x)\equiv p_N(x)e^{-V(x)/2}$ as before. In [DKMVZ2, Theorem 2.2] an asymptotic expansion, as $N\to\infty$, is derived for $\phi_N$ in the whole complex plane, and the leading term in the expansion in each region (see below) is computed explicitly. Here we are only interested in real values of the argument. Fix $\delta_0 > 0$ and sufficiently small (see [DKMVZ2]). For $0 < \delta \le \delta_0$ define

$$\begin{aligned} &I_5 \equiv (-\infty, -1-\delta), \qquad I_4 \equiv (-1-\delta, -1+\delta) \\ &I_3 \equiv (-1+\delta, 1-\delta), \qquad I_2 \equiv (1-\delta, 1+\delta) \\ &I_1 \equiv (1+\delta, +\infty) \end{aligned} \tag{4.6}$$

and let $\psi_N(x)$, $h_N(x)$, $h(x)$ be as in (2.5), (2.7) and (2.8), respectively. In [DKMVZ2], each of the regions $I_2$ and $I_4$ is split further into two regions, $I_2 = C_{1,\delta} \cup C_{2,\delta}$, $I_4 = D_{1,\delta} \cup D_{2,\delta}$, the estimates in $C_{2,\delta}$ and $D_{2,\delta}$ being finer than those in $C_{1,\delta}$ and $D_{1,\delta}$, respectively. For our purposes it is sufficient to conflate the regions and use the estimates in $C_{1,\delta}$ and $D_{1,\delta}$ for all points in $I_2$ and $I_4$, respectively. The apparent singularities in the formulae below for $z = \pm 1$ are of course removable (see e.g. (4.10) and (4.12) below). In what follows, notation of the type $O(1/N)$ as $N \to \infty$, means that the estimate holds for $x$ uniformly in the respective region (4.6) under consideration. Also the estimates are uniform for $\delta$ in *compact* subsets of $(0, \delta_0]$: in this connection see the important Remark 4.3 following (4.136) below. Finally, the constants $c_1, c_2, \cdots$ below may depend on $\delta$, but they are independent of $x$ and $N, P$.

We are ready to state [DKMVZ2, Theorem 2.2]. First, we consider the "exponential" regions $I_{1,5}$. Uniformly for all $|x| > 1+\delta$

$$\begin{aligned} \phi_N(c_N x + d_N) =& \frac{1}{\sqrt{4\pi c_N}}\, e^{-(N/2)\int_1^x |y^2-1|^{1/2} h_N(y)dy} \\ &\times \Big( \Big|\frac{x-1}{x+1}\Big|^{1/4} + \Big|\frac{x+1}{x-1}\Big|^{1/4} \Big) \Big(1 + O\Big(\frac{1}{N}\Big)\Big), \qquad x > 1+\delta \end{aligned}$$

and

$$\begin{aligned} \phi_N(c_N x + d_N) =& \frac{(-1)^N}{\sqrt{4\pi c_N}}\, e^{-(N/2)\int_x^{-1} |y^2-1|^{1/2} h_N(y)dy} \\ &\times \Big( \Big|\frac{x-1}{x+1}\Big|^{1/4} + \Big|\frac{x+1}{x-1}\Big|^{1/4} \Big) \Big(1 + O\Big(\frac{1}{N}\Big)\Big), \qquad x < -1-\delta. \end{aligned}$$

In particular, as

$$h_N(x) \ge h_{\min} > 0, \qquad x \in \mathbb{R}, \qquad N \ge N_1(V) \tag{4.7}$$

([DKMVZ2, Prop. 5.3]), we have for $|x| \ge 1+\delta$

$$|\phi_N(c_N x + d_N)| \le \frac{c_1}{\sqrt{4\pi c_N}}\, e^{-Nc_3}\, e^{-(N/4)c_2\,(x^2-(1+\delta)^2)}, \tag{4.8}$$

where we have used the fact that $|y^2-1|^{1/2} \ge c_2|y|$ for $|y| \ge 1+\delta$.

Next, consider the "Airy" region on the right, $I_2$. Uniformly for all $1-\delta < x < 1+\delta$

$$\begin{aligned} \phi_N(c_N x + d_N) = \frac{1}{\sqrt{c_N}} \Big( &\Big|\frac{x+1}{x-1}\Big|^{1/4} |f_N(x)|^{1/4}\, \mathrm{Ai}(f_N(x)) \Big(1 + O\Big(\frac{1}{N}\Big)\Big) \\ &- \Big|\frac{x-1}{x+1}\Big|^{1/4} \frac{1}{|f_N(x)|^{1/4}}\, \mathrm{Ai}'(f_N(x)) \Big(1 + O\Big(\frac{1}{N}\Big)\Big)\Big) \end{aligned} \tag{4.9}$$

where

$$f_N(x) = \alpha_N N^{2/3}\,(x-1)\hat{f}_N(x) \tag{4.10}$$

and in turn (see (the proof of) [DKMVZ2, Proposition 7.3])

(1) $\hat{f}_N(x)$ is real analytic on $(1-\delta, 1+\delta)$, and to any order $q = 0, 1, 2, \cdots$

$$\hat{f}_N(x) = \sum_{j=0}^{q} N^{-j/(2m)}\, \hat{f}_{(j)}(x) + O(N^{-(q+1)/(2m)})$$

uniformly for $x$ in the interval. Moreover, the functions $\hat{f}_{(j)}(x)$ are also real analytic on $1-\delta < x < 1+\delta$

(2) to any order $q = 1, 2, \cdots$

$$\alpha_N \equiv \big(h_N^2(1)/2\big)^{1/3} = 2m^{2/3} + \sum_{j=1}^{q} N^{-j/(2m)}\, \alpha_{(j)} + O(N^{-(q+1)/(2m)})$$

(3) $f_N'(x) = -\alpha_N N^{2/3} U_N(x)$, where $U_N(x) = \hat{f}_N(x) + (x-1)\hat{f}_N'(x)$ also has an expansion uniform in $x$ to any order $q = 0, 1, 2, \cdots$ as above

$$U_N(x) = \sum_{j=0}^{q} N^{-j/(2m)}\, U_{(j)}(x) + O(N^{-(q+1)/(2m)}).$$

The terms $U_{(j)}(x)$ are real analytic on $1-\delta < x < 1+\delta$

(4) $\max_{k=0,1,2} \max_{1-\delta \le x \le 1+\delta} |d^k \hat{f}_N(x)/dx^k| \le M < \infty$ for $N \ge N_2(V)$

(5) $\hat{f}_N(1) = 1 = U_N(1)$ and $\min_{1-\delta \le x \le 1+\delta} \hat{f}_N(x) \ge \frac{1}{2}$ for $N \ge N_2(V)$. Also $\hat{f}_{(0)}(1) = 1 = U_{(0)}(1)$.

Similarly, in the left "Airy" region $I_4$ we have uniformly for all $-1-\delta < x < -1+\delta$

(4.11)

$$\begin{aligned}\phi_N(c_N x + d_N) = \frac{(-1)^N}{\sqrt{c_N}} \bigg( \Big|\frac{x-1}{x+1}\Big|^{1/4} |\tilde{f}_N(x)|^{1/4}\, \mathrm{Ai}(-\tilde{f}_N(x)) \Big(1 + O\Big(\frac{1}{N}\Big)\Big) \\ - \Big|\frac{x+1}{x-1}\Big|^{1/4} \frac{1}{|\tilde{f}_N(x)|^{1/4}}\, \mathrm{Ai}'(-\tilde{f}_N(x)) \Big(1 + O\Big(\frac{1}{N}\Big)\Big)\bigg)\end{aligned}$$

where

$$\tilde{f}_N(x) = \tilde{\alpha}_N N^{2/3}\, (x+1)\hat{\tilde{f}}_N(x) \tag{4.12}$$

and in turn

(1) $\hat{\tilde{f}}_N(x)$ is real analytic on $(-1-\delta, -1+\delta)$, and to any order $q = 0, 1, 2, \cdots$

$$\hat{\tilde{f}}_N(x) = \sum_{j=0}^{q} N^{-j/(2m)}\, \hat{\tilde{f}}_{(j)}(x) + O(N^{-(q+1)/(2m)})$$

uniformly for $x$ in the interval. Moreover, the functions $\hat{\tilde{f}}_{(j)}(x)$ are also real analytic on $-1-\delta < x < -1+\delta$

(2) to any order $q = 1, 2, \cdots$

$$\tilde{\alpha}_N \equiv \big(h_N^2(-1)/2\big)^{1/3} = 2m^{2/3} + \sum_{j=1}^{q} N^{-j/(2m)}\, \tilde{\alpha}_{(j)} + O(N^{-(q+1)/(2m)})$$

(3) $\tilde{f}_N'(x) = -\tilde{\alpha}_N N^{2/3} \tilde{U}_N(x)$, where $\tilde{U}_N(x) = \hat{\tilde{f}}_N(x) + (x+1)\hat{\tilde{f}}_N'(x)$ also has an expansion uniform in $x$ to any order $q = 0, 1, 2, \cdots$ as above

$$\tilde{U}_N(x) = \sum_{j=0}^{q} N^{-j/(2m)}\, \tilde{U}_{(j)}(x) + O(N^{-(q+1)/(2m)}).$$

The terms $\tilde{U}_{(j)}(x)$ are real analytic on $1-\delta < x < 1+\delta$

(4) $\max_{k=0,1,2} \max_{-1-\delta \le x \le -1+\delta} |d^k \hat{\tilde{f}}_N(x)/dx^k| \le M < \infty$ for $N \ge N_2(V)$

(5) $\hat{\tilde{f}}_N(-1) = 1 = \tilde{U}_N(-1)$ and $\min_{-1-\delta \le x \le -1+\delta} \hat{\tilde{f}}_N(x) \ge \frac{1}{2}$ for $N \ge N_2(V)$. Also $\hat{\tilde{f}}_{(0)}(-1) = 1 = \tilde{U}_{(0)}(-1)$.

Finally, in the middle region $I_3$ we have uniformly for all $-1+\delta < x < 1-\delta$

(4.13)

$$\begin{aligned}\phi_N(c_N x + d_N) = {} & \sqrt{\frac{2}{\pi c_N}} \frac{1}{|1-x^2|^{1/4}} \\ & \times \Big( \cos\Big( \frac{N}{2} \int_1^x |1-y^2|^{1/2} h_N(y)dy + \frac{1}{2}\arcsin x \Big) \Big(1 + O\Big(\frac{1}{N}\Big)\Big) \\ & \qquad + \sin\Big( \frac{N}{2} \int_1^x |1-y^2|^{1/2} h_N(y)dy - \frac{1}{2}\arcsin x \Big) O\Big(\frac{1}{N}\Big)\Big).\end{aligned}$$

In the proof of Theorem 2.2 (and Theorem 3.1) we need the following two results that hold as $N \to \infty$,

(4.14)

$$\begin{aligned}\int_{-\infty}^{+\infty} \phi_N(y)\, dy \equiv c_N \int_{-\infty}^{+\infty} \phi_N(c_N y + d_N)\, dy \\ = c_N^{1/2}\, N^{-1/2}\, (2m)^{-1/2}\, (1 + (-1)^N + O(N^{-1/2}) + O(N^{-1/(2m)}))\end{aligned}$$

and uniformly for $x \in \mathbb{R}$

$$\Big| c_N \int_{-\infty}^{x} \phi_N(c_N y + d_N)\, dy \Big| = O\big(c_N^{1/2}\, N^{-1/2}\big) \tag{4.15}$$

or, equivalently, for any interval $K \subset \mathbb{R}$

$$\Big| c_N \int_{K} \phi_N(c_N y + d_N)\, dy \Big| = O\big(c_N^{1/2}\, N^{-1/2}\big).$$

**Notational remark:** In (4.14) above the reader may wonder why we write $O(N^{-1/2}) + O(N^{-1/(2m)})$ rather than just $O(N^{-1/(2m)})$. The reason is that the $O(N^{-1/(2m)})$ term arises simply from the evaluation of the constants $\alpha_N, \tilde{\alpha}_N$ arising in the asymptotics using (4.10)(2), (4.12)(2). The term $O(N^{-1/2})$, however, constitutes the detailed estimate of errors involved in evaluating the integral over all the different asymptotic regions (4.6). In order to separate out these two very different sources of error we have adopted the convention of writing error estimates in the form $O(N^{-1/2}) + O(N^{-1/(2m)})$ as above. We use this convention in all the estimates that follow in this Section. In the proof of Theorem 2.2, error estimates of the type $O(N^{-1/(2m)})$ also arise from the approximation of $h_N(x)$ in (2.7) (see e.g. (4.129) below).

Note that (4.14), (4.15) are direct consequences of the following more detailed statement, which is also used in the proof of Theorem 2.2.

**Proposition 4.3.** *The following holds uniformly for $x$ in the respective regions as $N \to \infty$. In $I_1$:*

$$\begin{aligned}\left|c_N \int_{1+\delta}^{x} \phi_N(c_N y + d_N)\, dy\right| &\le c_N \int_{1+\delta}^{x} \left|\phi_N(c_N y + d_N)\right| dy \\ &= O\big((c_N/N)^{1/2}\, e^{-Nc_3}\big), \qquad x \ge 1+\delta;\end{aligned} \tag{4.16}$$

*in $I_2$:*

(4.17)

$$c_N \int_{1-\delta}^{1+\delta} \phi_N(c_N y + d_N)\, dy = c_N^{1/2} N^{-1/2} (2m)^{-1/2} (1 + O(N^{-1/2}) + O(N^{-1/(2m)}))$$

$$\left|c_N \int_{1-\delta}^{x} \phi_N(c_N y + d_N)\, dy\right| \le O\Big(\frac{c_N^{1/2}}{N^{1/2}}\Big), \qquad 1-\delta \le x \le 1+\delta;$$

*in $I_3$:*

$$\left|c_N \int_{-1+\delta}^{x} \phi_N(c_N y + d_N)\, dy\right| \le O\Big(\frac{c_N^{1/2}}{N}\Big), \qquad -1+\delta \le x \le 1-\delta; \tag{4.18}$$

*in $I_4$:*

(4.19)

$$c_N \int_{-1-\delta}^{-1+\delta} \phi_N(c_N y + d_N)\, dy = c_N^{1/2} N^{-1/2} (2m)^{-1/2} (-1)^N (1 + O(N^{-1/2}) + O(N^{-1/(2m)}))$$

$$\left|c_N \int_{-1-\delta}^{x} \phi_N(c_N y + d_N)\, dy\right| \le O\Big(\frac{c_N^{1/2}}{N^{1/2}}\Big), \qquad -1-\delta \le x \le -1+\delta;$$

*in $I_5$:*

$$\begin{aligned}\left|c_N \int_{-\infty}^{x} \phi_N(c_N y + d_N)\, dy\right| &\le c_N \int_{-\infty}^{x} \left|\phi_N(c_N y + d_N)\right| dy \\ &= O\Big((c_N/N)^{1/2}\, e^{-Nc_3}\Big), \qquad x \le -1-\delta.\end{aligned} \tag{4.20}$$

*Proof.* We refer to (4.8) and note that

$$\int_{1+\delta}^{\infty} e^{-(N/4)c_2\,(y^2-(1+\delta)^2)}\, dy \le O(N^{-1/2})$$

which proves (4.16). The proof of (4.20) is similar.

We now prove (4.18). Uniformly for $-1+\delta \le x \le 1-\delta$, by (4.13),

$$\begin{aligned}
&c_N \int_{-1-\delta}^{x} \phi_N(c_N y + d_N)\, dy \\
&\quad = c_N^{1/2} \sqrt{\frac{2}{\pi}} \int_{-1+\delta}^{x} \Big[\cos\Big(\frac{N}{2}\int_1^y |1-t^2|^{1/2} h_N(t) dt + \frac{1}{2}\arcsin y\Big)\Big(1 + O\Big(\frac{1}{N}\Big)\Big) \\
&\qquad\qquad + \sin\Big(\frac{N}{2}\int_1^y |1-t^2|^{1/2} h_N(t) dt - \frac{1}{2}\arcsin y\Big) O\Big(\frac{1}{N}\Big)\Big] \frac{dy}{|1-y^2|^{1/4}} \\
&\quad = c_N^{1/2} \sqrt{\frac{2}{\pi}} \int_{-1+\delta}^{x} \cos\Big(\frac{N}{2}\int_1^y |1-t^2|^{1/2} h_N(t) dt + \frac{1}{2}\arcsin y\Big) \frac{dy}{|1-y^2|^{1/4}} \\
&\qquad\quad + O\Big(c_N^{1/2} N^{-1}\Big).
\end{aligned}$$

Integrating by parts in the last integral we find

$$\begin{aligned}
&c_N^{1/2}\int_{-1+\delta}^{x}\cos\Big(\frac{N}{2}\int_1^y|1-t^2|^{1/2}h_N(t)dt+\frac{1}{2}\arcsin y\Big)\frac{dy}{|1-y^2|^{1/4}}\\
&\qquad=c_N^{1/2}\frac{\sin\Big(\frac{N}{2}\int_1^y|1-t^2|^{1/2}h_N(t)dt+\frac{1}{2}\arcsin y\Big)}{\frac{N}{2}|1-y^2|^{3/4}h_N(y)+\frac{1}{2}|1-y^2|^{-1/4}}\Bigg|_{y=-1+\delta}^{x}\\
&\qquad\quad-c_N^{1/2}\int_{-1+\delta}^{x}\sin\Big(\frac{N}{2}\int_1^y|1-t^2|^{1/2}h_N(t)dt+\frac{1}{2}\arcsin y\Big)\\
&\qquad\qquad\times\frac{d}{dy}\bigg(\frac{1}{\frac{N}{2}|1-y^2|^{3/4}h_N(y)+\frac{1}{2}|1-y^2|^{-1/4}}\bigg)dy\\
&\qquad=O(c_N^{1/2}N^{-1})
\end{aligned}\tag{4.21}$$

uniformly for $x\in I_3$, which proves (4.18). (We have used (4.7) and also the uniform boundedness of $h_N'(x)$ on $I_3$, as $N\to\infty$.)

Now we prove (4.19). Uniformly for $-1-\delta\le x\le -1+\delta$, by (4.11),

$$\begin{aligned}
&c_N\int_{-1-\delta}^{x}\phi_N(c_Ny+d_N)\,dy\\
&\quad=(-1)^Nc_N^{1/2}\int_{-1-\delta}^{x}\bigg[\Big|\frac{y-1}{y+1}\Big|^{1/4}|\tilde f_N(y)|^{1/4}\operatorname{Ai}(-\tilde f_N(y))\Big(1+O\Big(\frac{1}{N}\Big)\Big)\\
&\qquad\quad-\Big|\frac{y+1}{y-1}\Big|^{1/4}\frac{1}{|\tilde f_N(y)|^{1/4}}\operatorname{Ai}'(-\tilde f_N(y))\Big(1+O\Big(\frac{1}{N}\Big)\Big)\bigg]\,dy.
\end{aligned}$$

Now

$$\begin{aligned}
&\bigg|c_N^{1/2}\int_{-1-\delta}^{x}\Big|\frac{y-1}{y+1}\Big|^{1/4}|\tilde f_N(y)|^{1/4}\operatorname{Ai}(-\tilde f_N(y))\,O\Big(\frac{1}{N}\Big)\,dy\bigg|\\
&\qquad\le C_1c_N^{1/2}O\Big(\frac{1}{N}\Big)\int_{-1-\delta}^{-1+\delta}\Big|\frac{y-1}{y+1}\Big|^{1/4}dy=O(c_N^{1/2}N^{-1})
\end{aligned}$$

where (cf. [AbSt])

$$C_1\equiv\sup_{y\in\mathbb{R}}|y|^{1/4}|\operatorname{Ai}(y)|<\infty.$$

Next, in view of (4.12) and the properties of $\hat{\tilde f}_N$,

$$\begin{aligned}
&\bigg|c_N^{1/2}\int_{-1-\delta}^{x}\Big|\frac{y+1}{y-1}\Big|^{1/4}\frac{\operatorname{Ai}'(-\tilde f_N(y))}{|\tilde f_N(y)|^{1/4}}\,O\Big(\frac{1}{N}\Big)\,dy\bigg|\\
&\qquad\le C_2c_N^{1/2}O\Big(\frac{1}{N}\Big)\int_{-1-\delta}^{-1+\delta}\Big|\frac{y+1}{y-1}\Big|^{1/4}\frac{1+|\tilde f_N(y)|^{1/4}}{|\tilde f_N(y)|^{1/4}}\,dy\\
&\qquad=C_2c_N^{1/2}O\Big(\frac{1}{N}\Big)\int_{-1-\delta}^{-1+\delta}\Big|\frac{y+1}{y-1}\Big|^{1/4}\frac{1+|\tilde\alpha_NN^{2/3}(y+1)\hat{\tilde f}_N(y)|^{1/4}}{|\tilde\alpha_NN^{2/3}(y+1)\hat{\tilde f}_N(y)|^{1/4}}\,dy\\
&\qquad\le\text{const}\cdot c_N^{1/2}N^{-1}\int_{-1-\delta}^{-1+\delta}\Big(N^{-1/6}+|y+1|^{1/4}\Big)\,dy=O(c_N^{1/2}N^{-1}),
\end{aligned}$$

where (cf. [AbSt])

$$C_2\equiv\sup_{y\in\mathbb{R}}(1+|y|^{1/4})^{-1}|\operatorname{Ai}'(y)|<\infty.\tag{4.22}$$

and we have used (see [DKMVZ2, (8.72)] and (4.12)(2))

$$\tilde{\alpha}_N \equiv \left(h_N^2(-1)/2\right)^{1/3} = 2m^{2/3} + O(N^{-1/(2m)}). \tag{4.23}$$

We consider next

$$c_N^{1/2} \int_{-1-\delta}^{x} \Big|\frac{y+1}{y-1}\Big|^{1/4} \frac{\mathrm{Ai}'(-\tilde{f}_N(y))}{|\tilde{f}_N(y)|^{1/4}}\, dy. \tag{4.24}$$

Make the change of variables (cf. (4.12), (4.23))

$$u(y) \equiv -\tilde{f}_N(y) = -\tilde{\alpha}_N N^{2/3}(y+1)\hat{\tilde{f}}_N(y) \tag{4.25}$$

which implies (cf. (4.12)(3))

$$\begin{aligned} \frac{du}{dy} &= -\tilde{\alpha}_N N^{2/3}\tilde{U}_N(y) \\ \tilde{U}_N(y) &\equiv \hat{\tilde{f}}_N(y) + (y+1)\hat{\tilde{f}}'_N(y). \end{aligned} \tag{4.26}$$

Note also that

$$y(u) + 1 = \frac{-u}{\tilde{\alpha}_N N^{2/3}\hat{\tilde{f}}_N(y(u))}. \tag{4.27}$$

Now set

$$B_N(y) \equiv \frac{1}{|y-1|^{1/4}|\hat{\tilde{f}}_N(y)|^{1/4}\,\tilde{U}_N(y)} \tag{4.28}$$

where the denominator is bounded away from 0 by (4.12) for $\delta > 0$ small enough. Then (4.24) becomes

(4.29)

$$\begin{aligned} &-\frac{c_N^{1/2}}{\tilde{\alpha}_N^{5/4}N^{5/6}} \int_{u(-1-\delta)}^{u(x)} \mathrm{Ai}'(u)\, B_N(y(u))\, du \\ &= -\frac{c_N^{1/2}}{\tilde{\alpha}_N^{5/4}N^{5/6}} \left[ \mathrm{Ai}(u)\, B_N(y(u))\Big|_{u(-1-\delta)}^{u(x)} + \frac{1}{\tilde{\alpha}_N N^{2/3}} \int_{u(-1-\delta)}^{u(x)} \mathrm{Ai}(u)\, \frac{B'(y(u))}{\tilde{U}_N(y(u))}\, du \right] \\ &= O(c_N^{1/2}N^{-5/6}) \end{aligned}$$

since the boundary term and the last integrand are uniformly bounded for $x \in I_4$ and the length of the interval of integration is $O(N^{2/3})$.

Finally we consider the integral of Ai. Set

$$L_N(y) \equiv \frac{|y-1|^{1/4}|\hat{\tilde{f}}_N(y)|^{1/4}}{\tilde{U}_N(y)} \tag{4.30}$$

and for future reference note that (see (4.12)(5))

$$L_N(-1) = 2^{1/4}. \tag{4.31}$$

Making the same change of variables (4.25), (4.26), (4.27) we find

$$\begin{aligned} c_N^{1/2}(-1)^N &\int_{-1-\delta}^{x} \Big|\frac{y+1}{y-1}\Big|^{1/4} |\tilde{f}_N(y)|^{1/4}\, \mathrm{Ai}(-\tilde{f}_N(y))\, dy \\ &= \frac{c_N^{1/2}(-1)^N}{\tilde{\alpha}_N^{3/4}N^{1/2}} \int_{u(x)}^{u(-1-\delta)} \mathrm{Ai}(u)\, L_N(y(u))\, du. \end{aligned} \tag{4.32}$$

In what follows we use the standard estimates (see [AbSt])

$$\begin{aligned} &|\operatorname{Ai}(t)| \le Ce^{-(2/3)t^{3/2}}, \qquad |\operatorname{Ai}'(t)| \le C(1+t^{1/4})e^{-(2/3)t^{3/2}}, \qquad t \ge 0,\\ &|\operatorname{Ai}(t)| \le C(1+|t|)^{-1/4}, \qquad |\operatorname{Ai}'(t)| \le C(1+|t|)^{1/4}, \qquad t<0 \end{aligned} \tag{4.33}$$

and

$$\Big|\int_{-\infty}^{t} \operatorname{Ai}(s)\,ds\Big| \le C(1+|t|)^{-3/4}, \qquad t \le 0. \tag{4.34}$$

Note that by (4.25), $u(y) > 0$ for $y < -1$ and $u(y) < 0$ for $y > -1$. Consider first (4.32) for $x < -1$:

$$\begin{aligned} &\Big|\frac{c_N^{1/2}}{\tilde\alpha_N^{3/4}N^{1/2}}(-1)^N \int_{u(x)}^{u(-1-\delta)} \operatorname{Ai}(u)\,L_N(y(u))\,du\Big| \\ &\qquad \le \frac{c_N^{1/2}}{\tilde\alpha_N^{3/4}N^{1/2}}\text{const}\int_0^\infty |\operatorname{Ai}(u)|\,du = O(c_N^{1/2}N^{-1/2}). \end{aligned} \tag{4.35}$$

Next we consider $x = -1$. Note that $u(-1) = 0$ and $\lim_{N\to\infty} u(-1-\delta) = +\infty$. Also for any fixed $u \in [0, u(-1-\delta)]$, $\lim_{N\to\infty} L_N(y(u)) = L_N(-1) = 2^{1/4}$ (see (4.27), (4.30), (4.31), and also (4.23)). We have

$$\begin{aligned} \int_0^{u(-1-\delta)} \operatorname{Ai}(u)\,L_N(y(u))\,du &= L_N(-1)\int_0^{u(-1-\delta)} \operatorname{Ai}(u)\,du \\ &\quad + \int_0^{u(-1-\delta)} \operatorname{Ai}(u)\,\big(L_N(y(u)) - L_N(-1)\big)\,du \equiv Q_{0,1} + Q_{0,2}. \end{aligned}$$

Now by (4.27)

$$\begin{aligned} |Q_{0,2}| &\le \int_0^{u(-1-\delta)} |\operatorname{Ai}(u)|\Big(\max_{-1-\delta\le t\le -1} |L_N'(t)|\Big)\,|y(u)+1|\,du \\ &\le c\int_0^{u(-1-\delta)} |\operatorname{Ai}(u)|\,\frac{u}{|\tilde\alpha_N N^{2/3}\hat{\hat{f}}(y(u))|}\,du \\ &\le cN^{-2/3}\int_0^\infty |u\operatorname{Ai}(u)|\,du = O(N^{-2/3}). \end{aligned}$$

Also $Q_{0,1} = 2^{1/4}\big(\int_0^\infty - \int_{u(-1-\delta)}^\infty\big)\operatorname{Ai}(u)\,du$. Taking (4.25), (4.33), (4.12)(2) into account, we conclude that

$$\begin{aligned} &\frac{c_N^{1/2}(-1)^N}{\tilde\alpha_N^{3/4}N^{1/2}}\int_0^{u(-1-\delta)} \operatorname{Ai}(u)\,L_N(y(u))\,du \\ &\qquad = \frac{c_N^{1/2}}{N^{1/2}}\frac{2^{1/4}}{(2m^{2/3})^{3/4}}(-1)^N\Big(\int_0^\infty \operatorname{Ai}(u)\,du + O(N^{-2/3}) + O(N^{-1/(2m)}) + O(e^{-cN})\Big). \end{aligned} \tag{4.36}$$

Now consider $-1 < x < -1+\delta$. We have $u(x) < 0$. Then (4.32) becomes

$$\frac{c_N^{1/2}(-1)^N}{\tilde\alpha_N^{3/4}N^{1/2}}\Big(\int_0^{u(-1-\delta)} + \int_{u(x)}^0\Big)\operatorname{Ai}(u)\,L_N(y(u))\,du \tag{4.37}$$

where the first integral was evaluated in (4.36), and for the second we write

$$\begin{aligned}\int_{u(x)}^{0} \mathrm{Ai}(u)\, L_N(y(u))\, du =& \Big(\int_{-\infty}^{u} \mathrm{Ai}(t)dt\Big)\, L_N(y(u))\Big|_{u(x)}^{0} \\ &- \int_{u(x)}^{0} \Big(\int_{-\infty}^{u} \mathrm{Ai}(t)dt\Big)\, \frac{L_N'(y(u))}{-\tilde{\alpha}_N N^{2/3} \tilde{U}_N(y(u))}\, du \\ =& 2^{1/4}\Big(\int_{-\infty}^{0} \mathrm{Ai}(t)dt\Big) - \Big(\int_{-\infty}^{u(x)} \mathrm{Ai}(t)dt\Big)\, L_N(y(u(x))) \\ &+ O\Big(\int_{u(x)}^{0} \frac{1}{(1+|u|)^{3/4}}\, \frac{du}{N^{2/3}}\Big)\end{aligned} \tag{4.38}$$

by (4.34) and the uniform boundness of $L_N'$. From this we draw two conclusions. First, uniformly for $-1 \leq x \leq -1+\delta$

$$\int_{u(x)}^{0} \mathrm{Ai}(u)\, L_N(y(u))\, du = O(1), \qquad N \to \infty, \tag{4.39}$$

and, second, again using (4.34), for any fixed $-1 < x \leq -1+\delta$, and in particular for $x = -1+\delta$,

$$\int_{u(x)}^{0} \mathrm{Ai}(u)\, L_N(y(u))\, du = 2^{1/4}\Big(\int_{-\infty}^{0} \mathrm{Ai}(t)dt\Big) + O(N^{-1/2}) \tag{4.40}$$

since

$$u(x) = -\tilde{\alpha}_N N^{2/3}(x+1)\hat{\hat{f}}_N(x) \leq -C(x)\, N^{2/3} \to -\infty. \tag{4.41}$$

Hence we conclude that the uniform estimate in (4.19) holds. Also recalling (4.36), (4.23) we find

$$\begin{aligned}&\frac{c_N^{1/2}(-1)^N}{\tilde{\alpha}_N^{3/4} N^{1/2}} 2^{1/4}\Big(\int_{0}^{+\infty} \mathrm{Ai}(t)\, dt + \int_{-\infty}^{0} \mathrm{Ai}(t)\, dt + O(N^{-1/2})\Big) \\ &\qquad = \frac{c_N^{1/2}}{N^{1/2}}\, (2m)^{-1/2}\, \Big((-1)^N + O(N^{-1/2}) + O(N^{-1/(2m)})\Big), \qquad N \to \infty,\end{aligned}$$

since $\int_{-\infty}^{\infty} \mathrm{Ai}(t)\, dt = 1$ (see [AbSt]), which proves the asymptotic formula in (4.19).

The proof of (4.17) is similar. □

4.2.2. *Proof of Theorem 3.1.* First we prove the second statement. By the definition of $\epsilon$

$$\begin{aligned}\epsilon\phi_N(x) &= \frac{1}{2}\Big[\int_{-\infty}^{x} \phi_N(y)\, dy - \int_{x}^{+\infty} \phi_N(y)\, dy\Big] \\ &= \frac{1}{2}\int_{-\infty}^{+\infty} \phi_N(y)\, dy - \int_{x}^{+\infty} \phi_N(y)\, dy.\end{aligned} \tag{4.42}$$

Making the change of variables $y \to c_N y + d_N$, we see that we have to estimate

$$\frac{1}{2} c_N \int_{-\infty}^{+\infty} \phi_N(c_N y + d_N)\, dy \tag{4.43}$$

and

$$c_N \int_{(x-d_N)/c_N}^{+\infty} \phi_N(c_N y + d_N)\, dy \tag{4.44}$$

The second statement now follows from (4.14), (4.15) above.

Now set $y = c_N x + d_N$, i.e. $x = (y - d_N)/c_N$. Note that if $y \in [-a, a]$ where $a = o(c_N)$, then $x = o(1)$, and hence $\phi_N(y) = \phi_N(c_N x + d_N)$ is given asymptotically by (4.13). Thus $\|\phi_N(y)\|_{L^\infty([-a,a])} = O(c_N^{-1/2})$ which proves the first statement.

Finally to prove the third statement (which is not used in this paper, but is of independent interest) we again introduce for $y$ the rescaled variable $x$, $y = c_N x + d_N$, and note that by (4.8),

$$\sup_{|x| \geq 1+\delta} |\phi_N(c_N x + d_N)| = O(c_N^{-1/2}\, e^{-N c_3}),$$

and by (4.13),

$$\sup_{|x| \leq 1-\delta} |\phi_N(c_N x + d_N)| = O(c_N^{-1/2}).$$

Thus $\|\phi_N\|_{L^\infty(\mathbb{R})}$ is determined by the Airy regions. Let us consider $-1 - \delta < x < -1 + \delta$ (the neighborhood of $x = 1$ is treated in the same way). We refer to (4.11) and note that by (4.25) the Ai term is of order

(4.45)
$$\big|c_N^{-1/2} |x+1|^{-1/4} |\tilde{\alpha}_N N^{2/3} (x+1) \hat{\hat{\tilde{f}}}_N(x)|^{1/4}\ \mathrm{Ai}(u(x))(1 + O(N^{-1})\big| = O(c_N^{-1/2} N^{1/6})$$

since Ai is bounded (and is not of a smaller order since e.g. $\mathrm{Ai}(u(-1)) = \mathrm{Ai}(0) \neq 0$). Also by (4.22) for the Ai$'$ term

$$\begin{aligned} &\left| c_N^{-1/2} |x+1|^{1/4} \frac{\mathrm{Ai}'(-\tilde{f}_N(x))}{|\tilde{f}_N(x)|^{1/4}} \,(1 + O(N^{-1}) \right| \\ &\qquad \leq C_2 c_N^{-1/2} |x+1|^{1/4} \frac{(1 + |\tilde{f}_N(x)|^{1/4})}{|\tilde{f}_N(x)|^{1/4}} \,(1 + O(N^{-1}) = O(c_N^{-1/2}) \end{aligned} \tag{4.46}$$

by (4.25).

*Remark* 4.2. Note that in [Sz] the estimate $\|\phi_N\|_{L^\infty(\mathbb{R})} = O(N^{-1/12})$ is proved for the case of Hermite polynomials, $V(x) = x^2$. Of course in view of (2.3) for $m = 1$ this is consistent with our estimate $O(c_N^{-1/2} N^{1/6})$. Also from our estimate and (2.3) it follows that, for all $m \geq 2$, $\|\phi_N\|_{L^\infty(\mathbb{R})}$ blows up as $N^{1/6 - 1/(4m)}$, as $N \to \infty$.

4.2.3. *Proof of Theorem 2.2.* From (2.10) we anticipate the leading order terms to be of order $\frac{c_N}{N} \sim \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}}$. Using (4.42) we find

$$\begin{aligned} (\phi_N, \epsilon \phi_P) =& \frac{1}{2} \Big( \int_{-\infty}^{+\infty} \phi_N(x)\, dx \Big) \Big( \int_{-\infty}^{+\infty} \phi_P(y)\, dy \Big) \\ &- \int_{-\infty}^{+\infty} \phi_N(x)\, dx \int_{x}^{+\infty} \phi_P(y)\, dy. \end{aligned} \tag{4.47}$$

By (4.14), as $N, P \dot{\to} \infty$, the first term in (4.47) equals

$$\begin{aligned} &\frac{1}{2} \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}} \frac{1}{2m} \,(1 + (-1)^N + O(N^{-1/2}) + O(N^{-1/(2m)})) \\ &\qquad\qquad \times (1 + (-1)^P + O(P^{-1/2}) + O(P^{-1/(2m)})). \end{aligned} \tag{4.48}$$

The estimation of the second term in (4.47) involves an extended region-by-region calculation, and concludes eventually at equation (4.136) below.

The following result will be used repeatedly throughout this Subsection.

**Lemma 4.4.** *For $V(x) = \kappa_{2m}x^{2m} + \cdots$, $\kappa_{2m} > 0$ as before, as $N, P \dot{\to} \infty$*

$$\frac{c_N}{c_P} - 1 = \frac{1}{2m}\frac{N-P}{P} + O(P^{-1-1/(2m)}) \tag{4.49}$$

*and*

$$\frac{d_N - d_P}{c_P} = O(P^{-1-1/m}). \tag{4.50}$$

*Proof.* Note first that for any fixed $\alpha > 0$, as $N, P \dot{\to} \infty$

$$\begin{aligned}\frac{1}{N^\alpha} - \frac{1}{P^\alpha} = \frac{1}{(P + (N-P))^\alpha} - \frac{1}{P^\alpha} &= \frac{1}{P^\alpha}\Big[-\alpha\frac{N-P}{P} + O\Big(\frac{1}{P^2}\Big)\Big] \\ &= O(P^{-1-\alpha}).\end{aligned} \tag{4.51}$$

This together with (2.3) for $q = 2m$ implies

$$\begin{aligned} c_N - c_P &= c_{(-1)}\big[N^{1/(2m)} - P^{(1/2m)}\big] + \sum_{k=1}^{2m}\big[c_{(k)}\cdot(N^{-k/(2m)} - P^{-k/(2m)})\big] + O(P^{-1-1/(2m)}) \\ &= c_{(-1)}\big[N^{1/(2m)} - P^{(1/2m)}\big] + \sum_{k=1}^{2m}\big[c_{(k)}\cdot O(N^{-1-k/(2m)})\big] + O(P^{-1-1/(2m)}). \end{aligned}$$

Therefore by (4.51)

$$\begin{aligned}\frac{c_N - c_P}{c_P} &= \frac{c_{(-1)}\big[N^{1/(2m)} - P^{(1/2m)}\big] + O(P^{-1-1/(2m)})}{c_{(-1)}P^{1/(2m)}(1 + O(P^{-1/(2m)})} \\ &= \big[((N/P)^{1/(2m)} - 1) + O(P^{-1-1/m})\big]\,(1 + O(P^{-1/(2m)})) \\ &= \Big[\frac{1}{2m}\frac{N-P}{P} + O(P^{-2}) + O(P^{-1-1/m})\Big]\,(1 + O(P^{-1/(2m)})) \\ &= \frac{1}{2m}\frac{N-P}{P} + O(P^{-1-1/(2m)}).\end{aligned}$$

The second statement follows similarly from (4.51) and (2.4) with $q = 2m$. □

We begin the estimation of the second term in (4.47) by making the change of variables $x \to c_N x + d_N$, $y \to c_P y + d_N$ in (4.47) to find

$$\begin{aligned}\int_{-\infty}^{+\infty}\phi_N(x)\,dx\int_x^{+\infty}\phi_P(y)\,dy &= c_N c_P\int_{-\infty}^{+\infty}\phi_N(c_N x + d_N)\,dx\int_{X(x)}^{+\infty}\phi_P(c_P y + d_P)\,dy \\ &= c_N c_P\sum_{j=1}^{5}\int_{I_j}\phi_N(c_N x + d_N)\,dx\int_{X(x)}^{+\infty}\phi_P(c_P y + d_P)\,dy\end{aligned} \tag{4.52}$$

where

$$X(x) \equiv x\frac{c_N}{c_P} + \frac{d_N - d_P}{c_P}. \tag{4.53}$$

Note that by Lemma 4.4

$$\frac{c_N}{c_P} = 1 + \frac{N-P}{2m}\frac{1}{P} + O(P^{-1-1/(2m)}) > 1, \qquad \frac{d_N - d_P}{c_P} = O(P^{-1-1/m})$$

and so for any fixed $x_0 > 0$ (e.g. $x_0 = 1/2$) and $N$, $P$ large enough

$$X(x) > x \quad \text{for } x \geq x_0, \qquad X(x) < x \quad \text{for } x \leq -x_0. \tag{4.54}$$

Now (4.16), (4.20), (4.15) imply

$$\begin{aligned} &c_N c_P \Big| \Big( \int_{I_1} + \int_{I_5} \Big) \phi_N(c_N x + d_N)\, dx \int_{X(x)}^{+\infty} \phi_P(c_P y + d_P)\, dy \Big| \\ &\quad \leq c_N c_P \Big( \Big( \int_{I_1} + \int_{I_5} \Big) \big| \phi_N(c_N x + d_N) \big|\, dx \Big) \cdot \sup_{x \in \mathbb{R}} \Big| \int_{x}^{+\infty} \phi_P(c_P y + d_P)\, dy \Big| \\ &\quad = O\big( (c_N c_P)^{1/2} (NP)^{-1/2}\, e^{-N c_3} \big). \end{aligned} \tag{4.55}$$

Hence in (4.52) we are left with three types of integrals

$$\begin{aligned} &c_N c_P \Big( \int_{I_4} + \int_{I_3} + \int_{I_2} \Big) \phi_N(c_N x + d_N)\, dx \int_{X(x)}^{+\infty} \phi_P(c_P y + d_P)\, dy \\ &\qquad \equiv J_4 + J_3 + J_2. \end{aligned} \tag{4.56}$$

By (4.54), uniformly for $x \in I_4$, we have $X(x) < x$ for $N$, $P$ sufficiently large. So

$$\begin{aligned} J_4 &= c_N c_P \int_{I_4} \phi_N(c_N x + d_N)\, dx \Big( \int_{X(x)}^{-1+\delta} + \int_{I_3} + \int_{I_2} + \int_{I_1} \Big) \phi_P(c_P y + d_P)\, dy \\ &\equiv J_{44} + J_{43} + J_{42} + J_{41}. \end{aligned} \tag{4.57}$$

Now $J_{43}$, $J_{42}$, $J_{41}$ are products of (4.19) with (4.18), (4.17), (4.16) respectively, and hence

$$J_{42} = \frac{(-1)^N}{2m} \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}} \left(1 + O(N^{-1/2}) + O(N^{-1/(2m)})\right) \tag{4.58}$$

and

$$J_{43} + J_{41} = \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}}\, O\big(P^{-1/2} + e^{-P c_3}\big) = \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}}\, O\big(P^{-1/2}\big). \tag{4.59}$$

Next define $x_{N,P} \in I_4$ such that

$$X(x_{N,P}) \equiv x_{N,P} \frac{c_N}{c_P} + \frac{d_N - d_P}{c_P} = -1 - \delta. \tag{4.60}$$

By Lemma 4.4, we see for $N, P \dot{\to} \infty$

$$x_{N,P} = -(1+\delta) \Big( 1 - \frac{1}{2m} \frac{N-P}{P} \Big) + O(P^{-1-1/(2m)}) \tag{4.61}$$

and so $x_{N,P}$ indeed lies in $I_4 = (-1-\delta, -1+\delta)$. With this notation

$$\begin{aligned} J_{44} &= \Big( \int_{-1-\delta}^{x_{N,P}} + \int_{x_{N,P}}^{-1+\delta} c_N \phi_N(c_N x + d_N)\, dx \Big( \int_{X(x)}^{-1+\delta} c_P \phi_P(c_P y + d_P)\, dy \Big) dx \\ &\equiv J'_{44} + J''_{44}. \end{aligned} \tag{4.62}$$

Observe that in $J''_{44}$, $-1-\delta < X(x) < -1+\delta$ and so all the points $y$ in the inner integral lie in $I_4$.

We need the bounds

$$\sup_{\min(|x+1|, |x-1|) \geq \delta} \big| \phi_N(c_N x + d_N) \big| \leq C\, c_N^{-1/2} \tag{4.63}$$

and

$$\sup_{|x\pm 1|\le\delta}\big|\phi_N(c_Nx+d_N)\big| \le \frac{C}{c_N^{1/2}}\,\frac{N^{1/6}}{(1+|x\pm 1|N^{2/3})^{1/4}} \tag{4.64}$$

which were almost proved in (4.45), (4.46). Indeed, the two inequalities preceding (4.45) imply (4.63), and as

$$|\operatorname{Ai}(f_N(x))| \le C\cdot(1+|f_N(x)|)^{-1/4}, \qquad |\operatorname{Ai}'(f_N(x))| \le C\cdot(1+|f_N(x)|)^{1/4},$$

we have for $|x-1|\le\delta$, by (4.10),

$$\begin{aligned}\frac{|f_N(x)|^{1/4}|\operatorname{Ai}(f_N(x))|}{|x-1|^{1/4}} &\le \frac{C|f_N(x)|^{1/4}}{|x-1|^{1/4}(1+|f_N(x)|)^{1/4}} \le \frac{C(N^{2/3})^{1/4}}{(1+|x-1|N^{2/3})^{1/4}}\\ \frac{|x-1|^{1/4}|\operatorname{Ai}'(f_N(x))|}{|f_N(x)|^{1/4}} &\le \frac{C(1+|x-1|N^{2/3})^{1/4}}{N^{1/6}} \le \text{const.}\end{aligned} \tag{4.65}$$

Inserting (4.65) and the analogous inequalities for $|x+1|\le\delta$ into (4.9), we obtain (4.64).

Now by (4.15), (4.64)

$$\begin{aligned}\big|J_{44}'\big| &\le c_N\int_{-1-\delta}^{x_{N,P}}\frac{C\,N^{1/6}}{c_N^{1/2}}\frac{C\,c_P^{1/2}}{P^{1/2}}\,dx\\ &= \text{const}\cdot\frac{(c_Nc_P)^{1/2}}{P^{1/2}}\,N^{1/6}\big(x_{N,P}+(1+\delta)\big).\end{aligned}$$

But by (4.61)

$$x_{N,P}+1+\delta = \frac{d_P-d_N}{c_P}+x_{N,P}\,\frac{c_P-c_N}{c_P} = O(P^{-1}).$$

Thus

$$\big|J_{44}'\big| \le \text{const}\cdot\frac{(c_Nc_P)^{1/2}}{P^{1/2}}\,N^{1/6}P^{-1} = \frac{(c_Nc_P)^{1/2}}{(NP)^{1/2}}\,O\big(N^{-1/3}\big). \tag{4.66}$$

Next

$$\begin{aligned}J_{44}'' = (-1)^Nc_N^{1/2}\int_{x_{N,P}}^{-1+\delta}&\bigg(\Big|\frac{x-1}{x+1}\Big|^{1/4}|\tilde f_N(x)|^{1/4}\operatorname{Ai}(-\tilde f_N(x))\Big(1+O\Big(\frac{1}{N}\Big)\Big)\\ &\quad-\Big|\frac{x+1}{x-1}\Big|^{1/4}\frac{1}{|\tilde f_N(x)|^{1/4}}\operatorname{Ai}'(-\tilde f_N(x))\Big(1+O\Big(\frac{1}{N}\Big)\Big)\bigg)\,dx\\ &\times\int_{X(x)}^{-1+\delta}c_P\phi_P(c_py+d_P)\,dy\end{aligned} \tag{4.67}$$

We consider first the two $O(N^{-1})$ terms. As

$$\big||\tilde f_N(x)|^{1/4}\operatorname{Ai}(-\tilde f_N(x))\big| \le C' \tag{4.68}$$

we have

$$\begin{aligned}
&c_N^{1/2}\int_{x_{N,P}}^{-1+\delta}\Big|\frac{x-1}{x+1}\Big|^{1/4}|\tilde f_N(x)|^{1/4}|\operatorname{Ai}(-\tilde f_N(x))|\,O\Big(\frac{1}{N}\Big)\,dx\\
&\qquad\qquad\times\sup_{x\in\mathbb{R}}\Big|\int_{X(x)}^{-1+\delta}c_P\phi_P(c_p y+d_P)\,dy\Big|\\
&\qquad\qquad\le \operatorname{const}C'\frac{(c_Nc_P)^{1/2}}{P^{1/2}}O(N^{-1})\int_{-1-\delta}^{-1+\delta}\Big|\frac{x-1}{x+1}\Big|^{1/4}dx=\frac{(c_Nc_P)^{1/2}}{(NP)^{1/2}}O(N^{-1/2}).
\end{aligned}\tag{4.69}$$

Also by the analog of the second inequality in (4.65) for $|x+1|\le\delta$,

$$\frac{|x+1|^{1/4}}{|\tilde f_N(x)|^{1/4}}\big|\operatorname{Ai}'(-\tilde f_N(x))\big|\le C'\tag{4.70}$$

and hence

$$\begin{aligned}
&c_N^{1/2}\int_{x_{N,P}}^{-1+\delta}\Big|\frac{x+1}{x-1}\Big|^{1/4}\frac{1}{|\tilde f_N(x)|^{1/4}}|\operatorname{Ai}'(-\tilde f_N(x))|\,O\Big(\frac{1}{N}\Big)\,dx\\
&\qquad\qquad\times\sup_{x\in\mathbb{R}}\Big|\int_{X(x)}^{-1+\delta}c_P\phi_P(c_p y+d_P)\,dy\Big|\\
&\qquad\qquad\le \operatorname{const}C'\frac{(c_Nc_P)^{1/2}}{P^{1/2}}O(N^{-1})\int_{-1-\delta}^{-1+\delta}\frac{1}{|x+1|^{1/4}}\,dx=\frac{(c_Nc_P)^{1/2}}{(NP)^{1/2}}O(N^{-1/2}).
\end{aligned}\tag{4.71}$$

Now we substitute (4.11) for $\phi_P(c_Py+d_P)$ in (4.67). Using an obvious schematic notation we note that, in view of (4.69), (4.71), we have shown

$$\begin{aligned}
J''_{44}=&\int_{x_{N,P}}^{-1+\delta}\big(\operatorname{Ai}+\operatorname{Ai}'\big)\,dx\int_{X(x)}^{-1+\delta}\big(\operatorname{Ai}+\operatorname{Ai}'\big)\,dy\\
&+\int_{x_{N,P}}^{-1+\delta}\big(\operatorname{Ai}+\operatorname{Ai}'\big)\,dx\int_{X(x)}^{-1+\delta}\big(\operatorname{Ai}\cdot O(P^{-1})+\operatorname{Ai}'\cdot O(P^{-1})\big)\,dy\\
&+\frac{(c_Nc_P)^{1/2}}{(NP)^{1/2}}O(N^{-1/2}),\qquad N,P\dot{\to}\infty.
\end{aligned}\tag{4.72}$$

To estimate the second integral in (4.72) we interchange the order of integration. Set $y_0\equiv X(-1+\delta)\in(-1-\delta,-1+\delta)$ and note that $X(x)$ is a 1-1 function from $[x_{N,P},-1+\delta]$ onto $[-1-\delta,y_0]$. We conclude that the second integral in (4.72) takes the form

(4.73)
$$
\begin{aligned}
(-1)^{N+P}(c_Nc_P)^{1/2}\int_{-1-\delta}^{y_0}&\Big(\Big|\frac{y-1}{y+1}\Big|^{1/4}|\tilde f_P(y)|^{1/4}\,\mathrm{Ai}(-\tilde f_P(y))\,O\Big(\frac{1}{P}\Big)\\
&\qquad-\Big|\frac{y+1}{y-1}\Big|^{1/4}\frac{1}{|\tilde f_P(y)|^{1/4}}\,\mathrm{Ai}'(-\tilde f_P(y))\,O\Big(\frac{1}{P}\Big)\Big)\,dy\\
\times\int_{x_{N,P}}^{X^{-1}(y)}&\Big(\Big|\frac{x-1}{x+1}\Big|^{1/4}|\tilde f_N(x)|^{1/4}\,\mathrm{Ai}(-\tilde f_N(x))\\
&\qquad-\Big|\frac{x+1}{x-1}\Big|^{1/4}\frac{1}{|\tilde f_N(x)|^{1/4}}\,\mathrm{Ai}'(-\tilde f_N(x))\Big)\,dx\\
+(-1)^{N+P}(c_Nc_P)^{1/2}\int_{y_0}^{-1+\delta}&\Big(\Big|\frac{y-1}{y+1}\Big|^{1/4}|\tilde f_P(y)|^{1/4}\,\mathrm{Ai}(-\tilde f_P(y))\,O\Big(\frac{1}{P}\Big)\\
&\qquad-\Big|\frac{y+1}{y-1}\Big|^{1/4}\frac{1}{|\tilde f_P(y)|^{1/4}}\,\mathrm{Ai}'(-\tilde f_P(y))\,O\Big(\frac{1}{P}\Big)\Big)\,dy\\
\times\int_{x_{N,P}}^{-1+\delta}&\Big(\Big|\frac{x-1}{x+1}\Big|^{1/4}|\tilde f_N(x)|^{1/4}\,\mathrm{Ai}(-\tilde f_N(x))\\
&\qquad-\Big|\frac{x+1}{x-1}\Big|^{1/4}\frac{1}{|\tilde f_N(x)|^{1/4}}\,\mathrm{Ai}'(-\tilde f_N(x))\Big)\,dx.
\end{aligned}
$$

Note that it follows from (4.29), (4.35), (4.36), (4.37), (4.39) that, uniformly for all intervals $K\subset[-1-\delta,-1+\delta]$,

$$
\begin{aligned}
\Big|\int_K\Big|\frac{y+1}{y-1}\Big|^{1/4}\frac{\mathrm{Ai}'(-\tilde f_N(y))}{|\tilde f_N(y)|^{1/4}}\,dy\Big|&\le C\,N^{-5/6}\\
\Big|\int_K\Big|\frac{y-1}{y+1}\Big|^{1/4}|\tilde f_N(y)|^{1/4}\;\mathrm{Ai}(-\tilde f_N(y))\,dy\Big|&\le C\,N^{-1/2}.
\end{aligned}
$$

This together with (4.68), (4.70) implies that (4.73) is of order

$$
(c_Nc_P)^{1/2}P^{-1}N^{-1/2}\int_{-1-\delta}^{-1+\delta}\frac{dx}{|x+1|^{1/4}}=\frac{(c_Nc_P)^{1/2}}{(NP)^{1/2}}\,O(N^{-1/2}). \tag{4.74}
$$

Thus (cf. (4.72)) we have shown (again schematically)

$$
\begin{aligned}
J''_{44}=&\int_{x_{N,P}}^{-1+\delta}\big(\,\mathrm{Ai}+\mathrm{Ai}'\,\big)\,dx\int_{X(x)}^{-1+\delta}\big(\,\mathrm{Ai}+\mathrm{Ai}'\,\big)\,dy\\
&+\frac{(c_Nc_P)^{1/2}}{(NP)^{1/2}}O(N^{-1/2}),\qquad N,P\stackrel{\cdot}{\to}\infty.
\end{aligned}\tag{4.75}
$$

Now we consider the terms of the form $\mathrm{Ai}'\times\mathrm{Ai}'$. After changing variables

$$
v=-\tilde f_P(y)=-\tilde\alpha_P P^{2/3}(y+1)\hat{\tilde f}_P(y),\qquad u=-\tilde f_N(x) \tag{4.76}
$$

and recalling (4.25), (4.26), (4.27) and (4.28), we find that the $\mathrm{Ai}' \times \mathrm{Ai}'$ term equals $(-1)^{N+P}(c_N c_P)^{1/2}$ times

$$\begin{aligned}
&\int_{x_{N,P}}^{-1+\delta} \Big|\frac{x+1}{x-1}\Big|^{1/4} \frac{1}{|\tilde{f}_N(x)|^{1/4}} \,\mathrm{Ai}'(-\tilde{f}_N(x))\,dx \\
&\qquad\qquad \times \int_{X(x)}^{-1+\delta} \Big|\frac{y+1}{y-1}\Big|^{1/4} \frac{1}{|\tilde{f}_P(y)|^{1/4}} \,\mathrm{Ai}'(-\tilde{f}_P(y))\,dy \\
&= \frac{1}{(\tilde{\alpha}_N\tilde{\alpha}_P)^{5/4}} \frac{1}{(NP)^{5/6}} \int_{u(-1+\delta)}^{u(x_{N,P})} B_N(x(u))\; \mathrm{Ai}'(u)\,du \\
&\qquad\qquad \times \int_{v(-1+\delta)}^{v(X(x(u)))} B_P(y(v))\; \mathrm{Ai}'(v)\,dv \\
&= \frac{1}{(\tilde{\alpha}_N\tilde{\alpha}_P)^{5/4}} \frac{1}{(NP)^{5/6}} \int_{u(-1+\delta)}^{u(x_{N,P})} B_N(x(u))\; \mathrm{Ai}'(u)\,du \\
&\qquad\qquad \times \Bigg( B_P(y(v(X(x(u)))))\,\mathrm{Ai}(v(X(x(u)))) \\
&\qquad\qquad\quad - B_P(y(v(-1+\delta)))\,\mathrm{Ai}(v(-1+\delta)) \\
&\qquad\qquad\quad + \frac{1}{\tilde{\alpha}_P P^{2/3}} \int_{v(-1+\delta)}^{v(X(x(u)))} \frac{B_P'(y(v))\; \mathrm{Ai}(v)}{\tilde{U}_P(y(v))}\,dv \Bigg) \\
&\equiv Q_{1,1} + Q_{1,2} + Q_{1,3}.
\end{aligned} \tag{4.77}$$

Note from (4.61) that $x_{N,P}+1 = -\delta + o(1)$ as $N, P \dot{\to} \infty$, and hence by (4.25)

$$u(x_{N,P}) \to +\infty, \qquad u(x_{N,P}) = O(N^{2/3}). \tag{4.78}$$

Now

$$\begin{aligned}
Q_{1,2} &= -\frac{1}{(\tilde{\alpha}_N\tilde{\alpha}_P)^{5/4}} \frac{1}{(NP)^{5/6}} B_P(y(v(-1+\delta)))\,\mathrm{Ai}(v(-1+\delta)) \\
&\qquad\qquad \times \Bigg( B_N(x(u))\,\mathrm{Ai}(u)\Big|_{u(-1+\delta)}^{u(x_{N,P})} \\
&\qquad\qquad\quad + \frac{1}{\tilde{\alpha}_N N^{2/3}} \int_{u(-1+\delta)}^{u(x_{N,P})} \frac{B_N'(x(u))\; \mathrm{Ai}(u)}{\tilde{U}_N(x(u))}\,du \Bigg) \\
&\le \frac{\mathrm{const}}{(NP)^{5/6}}
\end{aligned}$$

since the boundary term and the integrand are uniformly bounded and also the length of the interval of integration is $O(N^{2/3})$. Hence

$$|Q_{1,2}| \le \frac{\mathrm{const}}{(NP)^{1/2}}\, O(N^{-2/3}). \tag{4.79}$$

In $Q_{1,3}$ we integrate $\mathrm{Ai}'(u)$ and differentiate the other factors to obtain

$$\begin{aligned}\frac{1}{(\tilde{\alpha}_N\tilde{\alpha}_P)^{5/4}}\frac{1}{(NP)^{5/6}}\frac{1}{\tilde{\alpha}_P P^{2/3}}\Bigg[&\left(B_N(x(u))\ \mathrm{Ai}(u)\int_{v(-1+\delta)}^{v(X(x(u)))}\frac{B_P'(y(v))\ \mathrm{Ai}(v)}{\tilde{U}_P(y(v))}\,dv\right)\Bigg|_{u(-1+\delta)}^{u(x_{N,P})}\\ &-\int_{u(-1+\delta)}^{u(x_{N,P})}\mathrm{Ai}(u)\,\frac{B_N'(x(u))}{-\tilde{\alpha}_N N^{2/3}\tilde{U}_N(x(u))}\,du\int_{v(-1+\delta)}^{v(X(x(u)))}\frac{B_P'(y(v))\ \mathrm{Ai}(v)}{\tilde{U}_P(y(v))}\,dv\\ &-\int_{u(-1+\delta)}^{u(x_{N,P})}\mathrm{Ai}(u)\,B_N(x(u))\,\frac{B_P'(X(x(u)))\,\mathrm{Ai}(v(X(x(u))))}{\tilde{U}_P(X(x(u)))}\\ &\qquad\times\frac{d}{du}\big[v(X(x(u)))\big]\,du\Bigg]\equiv Q_{1,3}'+Q_{1,3}''+Q_{1,3}'''\end{aligned}\tag{4.80}$$

where we have used

$$y\circ v=\mathrm{id}.\tag{4.81}$$

Set

$$\beta(u)\equiv v(X(x(u))).\tag{4.82}$$

**Lemma 4.5.** *Uniformly for $u(-1+\delta)<u<u(-1-\delta)$, we have as $N,P\overset{\cdot}{\to}\infty$*

$$\beta(u)=O(P^{-1/3})+(1+O(P^{-1}))\,u\tag{4.83}$$

*and*

$$\frac{d}{du}\beta(u)=1+O\Big(\frac{1}{P}\Big).\tag{4.84}$$

*Proof.* We have in view of (4.26), (4.53)

$$\begin{aligned}\frac{d}{du}\beta(u)&=v'(X(x(u)))\,X'(x(u))\,x'(u)\\ &=\frac{P^{2/3}}{N^{2/3}}\frac{\tilde{\alpha}_P}{\tilde{\alpha}_N}\frac{c_N}{c_P}\frac{\tilde{U}_P(X(x(u)))}{\tilde{U}_N(x(u))}\\ &=\frac{P^{2/3}}{N^{2/3}}\frac{\tilde{\alpha}_P}{\tilde{\alpha}_N}\frac{c_N}{c_P}\left(\frac{\tilde{U}_P(x(u))}{\tilde{U}_N(x(u))}+\frac{\tilde{U}_P(X(x(u)))-\tilde{U}_P(x(u))}{\tilde{U}_N(x(u))}\right).\end{aligned}$$

By Lemma 4.4, $\frac{c_N}{c_P}-1=O(P^{-1})$, and also $X(x(u))-x(u)=(\frac{c_N}{c_P}-1)x(u)+(d_N-d_P)/c_P=O(P^{-1})$ since $x(u)\in[-1-\delta,-1+\delta]$ is uniformly bounded. Also by (4.12) $\max_{u\in[u(-1+\delta),u(-1-\delta)]}\tilde{U}_P'(u)\le\text{const}$. In addition, $\tilde{\alpha}_N$ also has a complete expansion in powers of $N^{-1/(2m)}$ (see (4.12)), and therefore by the same argument as in the proof of Lemma 4.4, $\frac{\tilde{\alpha}_P}{\tilde{\alpha}_N}=1+O(P^{-1})$, as $N,P\overset{\cdot}{\to}\infty$. Finally as $\tilde{U}_N(x)$ has a complete expansion (see (4.12)) in powers of $N^{-1/(2m)}$, uniformly for $-1-\delta\le x\le-1+\delta$, we again find $\frac{\tilde{U}_P(x(u))}{\tilde{U}_N(x(u))}=1+O(P^{-1})$. Thus

$$\begin{aligned}\frac{d}{du}\beta(u)&=\big(1+O(P^{-1})\big)\,\big(1+O(P^{-1})\big)\,\big(1+O(P^{-1})\big)\,\big(1+O(P^{-1})\big)\\ &=1+O(P^{-1})\end{aligned}$$

uniformly for $u(-1+\delta)\le u\le u(-1-\delta)$ which proves (4.84).

Finally by (4.84)

$$\beta(u)=\beta(0)+\int_0^u\beta'(t)\,dt=\beta(0)+u\cdot\big(1+O(P^{-1})\big)$$

which together with

$$\begin{aligned}\beta(0) &= v(X(-1)) = v\big(-c_N/c_P + (d_N - d_P)/c_P\big) = v\big(-1 + O(P^{-1})\big)\\ &= -\tilde{\alpha}_P P^{2/3}\big(1 - 1 + O(P^{-1})\big)\,\hat{\tilde{f}}_P\big(-1 + O(P^{-1})\big) = O(P^{-1/3})\end{aligned}$$

proves (4.83). □

Now we use the uniform boundedness of the functions Ai, $B_N$, $B'_N$, $\frac{1}{\tilde{U}_N}$, and also (4.84), to conclude the following. Note that the second estimate in (4.33) holds on the whole of $\mathbb{R}$ and that the lengths of the intervals of integration in (4.80) are of order $O(N^{2/3})$. Hence the integral in $Q'_{1,3}$ (and also the whole boundary term) is bounded by $O((N^{2/3})^{3/4}) = O(N^{1/2})$. The term $Q''_{1,3}$ is estimated in a similar way. Finally, in $Q'''_{1,3}$ we just estimate the integrand by a constant (note (4.84)). We then obtain

$$\begin{aligned}|Q'_{1,3}| &\le \frac{\text{const}}{P^{7/3}}\,N^{1/2} = \frac{1}{(NP)^{1/2}}\,O(P^{-5/6})\\ |Q''_{1,3}| &\le \frac{\text{const}}{P^{7/3}}\,\frac{1}{N^{2/3}}\,(N^{2/3})^{3/4}\,(N^{2/3})^{3/4} = \frac{1}{(NP)^{1/2}}\,O(P^{-1})\\ |Q'''_{1,3}| &\le \frac{\text{const}}{P^{7/3}}\,N^{2/3} = \frac{1}{(NP)^{1/2}}\,O(P^{-2/3}).\end{aligned}\tag{4.85}$$

Finally we consider the term $Q_{1,1}$ in (4.77). Using (4.81), (4.82) we rewrite $Q_{1,1}$ as

$$\begin{aligned}&\frac{1}{(\tilde{\alpha}_N\tilde{\alpha}_P)^{5/4}}\,\frac{1}{(NP)^{5/6}}\left(\int_{u(-1+\delta)}^{-1} + \int_{-1}^{u(x_{N,P})}\right) B_N(x(u))\ \mathrm{Ai}'(u)\\ &\qquad\qquad\qquad\qquad \times B_P(X(x(u)))\,\mathrm{Ai}(\beta(u))\,du\\ &\quad \equiv Q'_{1,1} + Q''_{1,1}.\end{aligned}$$

Now in $Q''_{1,1}$, $u(x_{N,P}) \to +\infty$, and we use the estimate for Ai$'$ in (4.33) (which clearly also holds on $[-1, +\infty)$) together with the boundedness of the other factors to find

$$|Q''_{1,1}| \le \frac{\text{const}}{(NP)^{5/6}} = \frac{1}{(NP)^{1/2}}O(N^{-2/3}).\tag{4.86}$$

Note that by (4.83), for $P$ and $N$ large enough,

$$C^{-1} \le \frac{u}{\beta(u)} \le C \qquad \text{and} \qquad \beta(u) \le -1/2\tag{4.87}$$

uniformly for $u(-1+\delta) < u < -1$. Hence using (4.33), and recalling that $B_N$, $B'_N$ are uniformly bounded, and also using the properties of $\hat{\tilde{f}}_N$ after (4.12) (cf. (4.41)), we find

$$|Q'_{1,1}| \le \frac{\text{const}}{(NP)^{5/6}}\int_{-\delta N^{2/3}\hat{\tilde{f}}_N(-1+\delta)}^{-1}\left|\frac{u}{\beta(u)}\right|^{1/4} du \le \text{const}\frac{\delta}{(NP)^{1/2}}.\tag{4.88}$$

We will see below that it is important that we can choose $\delta$ to be arbitrarily small. Collecting the estimates (4.86), (4.88), (4.79), (4.85) in (4.77) and recalling (4.75) we conclude (schematically)

$$
\begin{aligned}
J''_{44} = &\int_{x_{N,P}}^{-1+\delta} \mathrm{Ai}\ dx \int_{X(x)}^{-1+\delta} \mathrm{Ai}\ dy \\
&+ \int_{x_{N,P}}^{-1+\delta} \mathrm{Ai}\ dx \int_{X(x)}^{-1+\delta} \mathrm{Ai}'\ dy + \int_{x_{N,P}}^{-1+\delta} \mathrm{Ai}'\ dx \int_{X(x)}^{-1+\delta} \mathrm{Ai}\ dy \\
&+ \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}} O(\delta + N^{-1/2}), \qquad N, P \dot{\to} \infty.
\end{aligned} \tag{4.89}
$$

Now we consider the term of the form $\mathrm{Ai} \times \mathrm{Ai}'$ in (4.89). After changing variables as in (4.76) and using (4.81), we find that the $\mathrm{Ai} \times \mathrm{Ai}'$ term equals $(-1)^{N+P}(c_N c_P)^{1/2}$ times

$$
\begin{aligned}
&\int_{x_{N,P}}^{-1+\delta} \Big|\frac{x-1}{x+1}\Big|^{1/4} |\tilde{f}_N(x)|^{1/4}\ \mathrm{Ai}(-\tilde{f}_N(x))\, dx \\
&\qquad \times \int_{X(x)}^{-1+\delta} \Big|\frac{y+1}{y-1}\Big|^{1/4} \frac{1}{|\tilde{f}_P(y)|^{1/4}}\ \mathrm{Ai}'(-\tilde{f}_P(y))\, dy \\
&= \frac{1}{\tilde{\alpha}_N^{3/4} \tilde{\alpha}_P^{5/4}} \frac{1}{N^{1/2} P^{5/6}} \int_{u(-1+\delta)}^{u(x_{N,P})} L_N(x(u))\ \mathrm{Ai}(u)\, du \\
&\qquad \times \int_{v(-1+\delta)}^{v(X(x(u)))} B_P(y(v))\ \mathrm{Ai}'(v)\, dv \\
&= \frac{1}{\tilde{\alpha}_N^{3/4} \tilde{\alpha}_P^{5/4}} \frac{1}{N^{1/2} P^{5/6}} \int_{u(-1+\delta)}^{u(x_{N,P})} L_N(x(u))\ \mathrm{Ai}(u)\, du \\
&\qquad \times \Big( B_P(X(x(u)))\, \mathrm{Ai}(v(X(x(u)))) \\
&\qquad\quad - B_P(-1+\delta)\, \mathrm{Ai}(v(-1+\delta)) \\
&\qquad\quad + \frac{1}{\tilde{\alpha}_P P^{2/3}} \int_{v(-1+\delta)}^{v(X(x(u)))} \frac{B'_P(y(v))\ \mathrm{Ai}(v)}{\tilde{U}_P(y(v))}\, dv \Big) \\
&\equiv Q_{2,1} + Q_{2,2} + Q_{2,3}.
\end{aligned} \tag{4.90}
$$

In $Q_{2,3}$ we split both integrals into the parts with positive and negative arguments. Using (4.33) and (4.12) we find

$$
\begin{aligned}
|Q_{2,3}| \le &\frac{\text{const}}{N^{1/2} P^{5/6}} \frac{1}{P^{2/3}} \Big( \text{const} + \int_{u(-1+\delta)}^{0} \frac{du}{(1+|u|)^{1/4}} \Big) \\
&\qquad \times \Big( \text{const} + \int_{v(-1+\delta)}^{0} \frac{dv}{(1+|v|)^{1/4}} \Big) \\
&\le \frac{\text{const}}{N^{1/2} P^{3/2}} \big( C + (\delta N^{2/3})^{3/4} \big) \big( C + (\delta P^{2/3})^{3/4} \big) \\
&\le \frac{1}{(NP)^{1/2}}\, O\big(\delta^{3/2} + P^{-1}\big).
\end{aligned} \tag{4.91}
$$

Next by (4.81)

$$
\begin{aligned}
|Q_{2,2}| \le & \frac{C}{N^{1/2}P^{5/6}} \Big( \int_{u(-1+\delta)}^{u(x_{N,P})} L_N(x(u) \ \mathrm{Ai}(u)\, du \Big) B_P(-1+\delta)\, \mathrm{Ai}(v(-1+\delta)) \\
\le & \frac{C}{(NP)^{2/3}} \Big( \mathrm{const} + \int_{u(-1+\delta)}^{-1} L_N(x(u) \ \mathrm{Ai}(u)\, du \Big) O(1) \\
= & \frac{1}{(NP)^{2/3}}\, O(1) \Big[ \mathrm{const} + \Big( \int_{-\infty}^{u} \mathrm{Ai}(t) dt \Big) L_N(x(u)) \Big|_{u(-1+\delta)}^{-1} \\
& \qquad + \frac{1}{\tilde{\alpha}_N N^{2/3}} \int_{u(-1+\delta)}^{-1} \Big( \int_{-\infty}^{u} \mathrm{Ai}(t) dt \Big) \frac{L_N'(x(u))}{\tilde{U}_N(x(u))}\, du \Big] \\
\le & \frac{1}{(NP)^{2/3}}\, O(1)\, \mathrm{const} = \frac{1}{(NP)^{1/2}}\, O(P^{-1/3})
\end{aligned} \tag{4.92}
$$

since the length of the interval of integration is $O(N^{2/3})$ (again by (4.12)) and since $\int_{-\infty}^{u} \mathrm{Ai}(t) dt$ and all other functions are uniformly bounded.

Finally splitting the interval as in (4.91), and using (4.12), (4.87) and (4.33), we find

$$
\begin{aligned}
|Q_{2,1}| \le & \frac{\mathrm{const}}{N^{1/2}P^{5/6}} \Big( \mathrm{const} + \int_{-\delta N^{2/3} \hat{\tilde{f}}_N(-1+\delta)}^{-1} |\, \mathrm{Ai}(u)|\, |\, \mathrm{Ai}(\beta(u))|\, du \Big) \\
\le & \frac{\mathrm{const}}{(NP)^{1/2}} \frac{1}{P^{1/3}} \Big( \mathrm{const} + \int_{1}^{\delta N^{2/3} \hat{\tilde{f}}_N(-1+\delta)} \frac{du}{u^{1/2}} \Big) \\
\le & \frac{\mathrm{const}}{(NP)^{1/2}} \frac{1}{P^{1/3}} \big( \mathrm{const} + (\delta N^{2/3})^{1/2} \big) \\
\le & \frac{C}{(NP)^{1/2}}\, O\big(\delta^{1/2} + N^{-1/3}\big).
\end{aligned} \tag{4.93}
$$

Note that the integral of the form $\mathrm{Ai}' \times \mathrm{Ai}$ in (4.89) can be estimated similarly after changing the order of integration as in (4.72) above: the estimate is then the same as for the integral of the form $\mathrm{Ai} \times \mathrm{Ai}'$. Thus collecting the estimates (4.93), (4.92), (4.91) in (4.90), and recalling (4.89), we conclude (schematically)

$$
\begin{aligned}
J_{44}'' = & \int_{x_{N,P}}^{-1+\delta} \mathrm{Ai}\ dx \int_{X(x)}^{-1+\delta} \mathrm{Ai}\ dy \\
& + \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}} O(\delta^{1/2} + \delta + \delta^{3/2} + P^{-1/3}), \qquad N, P \overset{\cdot}{\to} \infty.
\end{aligned} \tag{4.94}
$$

Note for future reference that we will eventually take a limit $\delta \to 0$, and hence we leave only the term $O(\delta^{1/2})$ in the above estimate (and in similar ones below).

It remains to analyze the $\mathrm{Ai} \times \mathrm{Ai}$ integral in (4.94). As we will see, this is the only term that contributes to leading order in $J_{44}$. Making the same change of variables that led to (4.90), and recalling (4.30), (4.82), we write the $\mathrm{Ai} \times \mathrm{Ai}$ integral as

$(-1)^{N+P}\sqrt{c_N c_P}$ times

$$\begin{aligned}
&\frac{1}{(\tilde{\alpha}_N\tilde{\alpha}_P)^{3/4}}\frac{1}{(NP)^{1/2}}\Big(\int_{u(-1+\delta)}^{0}+\int_{0}^{u(x_{N,P})}\Big)L_N(x(u))\ \mathrm{Ai}(u)\,du\\
&\qquad\qquad\times\int_{v(-1+\delta)}^{\beta(u)}L_P(y(v))\ \mathrm{Ai}(v)\,dv\\
&\qquad\equiv\frac{1}{(\tilde{\alpha}_N\tilde{\alpha}_P)^{3/4}}\frac{1}{(NP)^{1/2}}\Big(Q_{3,1}+Q_{3,2}\Big).
\end{aligned}\tag{4.95}$$

Note that in $Q_{3,2}$, $u\geq 0$ (recall $\lim_{N,P\dot{\to}\infty}u(x_{N,P})=+\infty$). It follows from (4.83) that there is a number $u_0>0$, $u_0=O(P^{-1/3})$ such that $\beta(u)\geq 0$ for $u\geq u_0$. We write

$$\begin{aligned}
Q_{3,2}=&\int_0^{u_0}L_N(x(u))\ \mathrm{Ai}(u)\,du\int_{v(-1+\delta)}^{\beta(u)}L_P(y(v))\ \mathrm{Ai}(v)\,dv\\
&+\int_{u_0}^{u(x_{N,P})}L_N(x(u))\ \mathrm{Ai}(u)\,du\\
&\qquad\times\Big(\int_{v(-1+\delta)}^{0}+\int_0^{\beta(u)}\Big)L_P(y(v))\ \mathrm{Ai}(v)\,dv\\
&\equiv Q_{4,1}+Q_{4,2}+Q_{4,3}.
\end{aligned}\tag{4.96}$$

Now

$$\begin{aligned}
|Q_{4,1}|=&\Bigg|\int_0^{u_0}L_N(x(u))\,\mathrm{Ai}(u)\,du\Bigg[L_P(y(v))\Big(\int_{-\infty}^{v}\mathrm{Ai}(t)dt\Big)\Bigg|_{v(-1+\delta)}^{\beta(u)}\\
&\qquad+\frac{1}{\tilde{\alpha}_P P^{2/3}}\int_{v(-1+\delta)}^{\beta(u)}\frac{L_P'(y(v))}{\tilde{U}_P(y(v))}\Big(\int_{-\infty}^{v}\mathrm{Ai}(t)dt\Big)\,dv\Bigg]\Bigg|\\
&\leq\frac{\mathrm{const}}{P^{1/3}}
\end{aligned}\tag{4.97}$$

since $u_0=O(P^{-1/3})$, all the integrands in the $du$ and $dv$ integrals are uniformly bounded, and the length of the inner interval of integration is $O(P^{2/3})$ (recall (4.83)).

To estimate $Q_{4,3}$ recall first from (4.27) that $y(v)-(-1)=-\frac{v}{\tilde{\alpha}_P P^{2/3}\tilde{U}_P(y(v))}$. Also by (4.12), $|L_P'|$ is uniformly bounded. Hence

$$\big|L_P(y(v))-L_P(-1)\big|\leq\mathrm{const}\cdot|y(v)-(-1)|\leq\mathrm{const}\cdot|v|\cdot P^{-2/3}.\tag{4.98}$$

Thus, since $\beta(u)\geq 0$, using (4.26), (4.33) and the fact that $\hat{\hat{f}}_N(-1)=1$, we obtain

$$\begin{aligned}
\int_0^{\beta(u)}L_P(y(v))\ \mathrm{Ai}(v)\,dv=&L_P(-1)\int_0^{\beta(u)}\mathrm{Ai}(v)\,dv+O\Big(\int_0^{\beta(u)}\frac{v\,e^{-(2/3)v^{3/2}}}{P^{2/3}}\,dv\Big)\\
=&2^{1/4}\int_0^{\beta(u)}\mathrm{Ai}(v)\,dv+O(P^{-2/3}),
\end{aligned}$$

and so

$$
\begin{aligned}
Q_{4,3} &= \int_{u_0}^{u(x_{N,P})} L_N(x(u))\ \mathrm{Ai}(u)\Big(2^{1/4}\int_0^{\beta(u)} \mathrm{Ai}(v)\,dv + O(P^{-2/3})\Big)\,du \\
&= 2^{1/4}\int_{u_0}^{u(x_{N,P})} L_N(x(u))\ \mathrm{Ai}(u)\ \Big(\int_0^{\beta(u)} \mathrm{Ai}(v)\,dv\Big)\,du + O(P^{-2/3})
\end{aligned}
\tag{4.99}
$$

again by (4.33). Next

$$
\begin{aligned}
&\int_{u_0}^{u(x_{N,P})} L_N(x(u))\ \mathrm{Ai}(u)\,du \int_0^{\beta(u)} \mathrm{Ai}(v)\,dv \\
&\qquad = L_N(-1)\int_{u_0}^{u(x_{N,P})} \mathrm{Ai}(u)\,du \int_0^{\beta(u)} \mathrm{Ai}(v)\,dv \\
&\qquad + \int_{u_0}^{u(x_{N,P})} \big(L_N(x(u)) - L_N(-1)\big)\ \mathrm{Ai}(u)\,du \int_0^{\beta(u)} \mathrm{Ai}(v)\,dv \equiv Q_{5,1} + Q_{5,2}.
\end{aligned}
\tag{4.100}
$$

Now by the uniform boundedness of $L_N'$ and (4.27), and since $\int_0^\infty |v^j \,\mathrm{Ai}(v)| dv < \infty$, $j = 0, 1$,

$$
|Q_{5,2}| \le c\int_{u_0}^{u(x_{N,P})} |x(u){+}1|\,|\,\mathrm{Ai}(u)|\,du \le c_1 N^{-2/3}\int_{u_0}^{u(x_{N,P})} |u\,\mathrm{Ai}(u)|\,du = O(N^{-2/3}).
$$

Next

$$
Q_{5,1} = 2^{1/4}\int_{u_0}^{u(x_{N,P})} \mathrm{Ai}(u)\,du\Big(\int_0^u + \int_u^{\beta(u)}\Big)\,\mathrm{Ai}(v)\,dv \equiv Q_{5,1}' + Q_{5,1}''.
$$

We have by (4.83)

$$
\Big|\int_u^{\beta(u)} \mathrm{Ai}(v)\,dv\Big| \le c|\beta(u) - u| = O(P^{-1/3} + P^{-1}|u|)
$$

and hence

$$
|Q_{5,1}''| \le 2^{1/4}\int_0^\infty |\,\mathrm{Ai}(u)|\,O(P^{-1/3} + P^{-1}|u|)\,du = O(P^{-1/3}).
$$

Also

$$
\begin{aligned}
Q_{5,1}' &= 2^{1/4}\Big(\int_0^\infty - \int_0^{u_0} - \int_{u(x_{N,P})}^\infty\Big)\,\mathrm{Ai}(u)\,du\int_0^u \mathrm{Ai}(v)\,dv \\
&= 2^{1/4}\int_0^\infty \mathrm{Ai}(u)\,du\int_0^u \mathrm{Ai}(v)\,dv + O(P^{-1/3}) + O(e^{-cN})
\end{aligned}
$$

where we have used the fact that $u_0 = O(P^{-1/3})$, the uniform boundedness of $\int_0^u \mathrm{Ai}(v)dv$ for $u \ge 0$, and the (super)exponential decay of $\mathrm{Ai}(v)$ for $v \ge 0$. Thus

$$
Q_{4,3} = 2^{1/2}\int_0^{+\infty} \mathrm{Ai}(u)\,du\int_0^u \mathrm{Ai}(v)\,dv + O(P^{-1/3}),
\tag{4.101}
$$

$N, P \dot{\to} \infty$.

To estimate $Q_{4,2}$ we first write

$$\begin{aligned}
&\int_{v(-1+\delta)}^{0} L_P(y(v))\ \mathrm{Ai}(v)\,dv \\
&\qquad = L_P(-1)\int_{-\infty}^{0}\mathrm{Ai}(t)\,dt - L_P(y(v(-1+\delta)))\int_{-\infty}^{v(-1+\delta)}\mathrm{Ai}(t)\,dt \\
&\qquad\quad + \frac{1}{\tilde{\alpha}_P P^{2/3}}\int_{v(-1+\delta)}^{0}\Big(\int_{-\infty}^{v}\mathrm{Ai}(t)\,dt\Big)\,\frac{L_P'(y(v))}{\tilde{U}_P(y(v))}\,dv.
\end{aligned}\tag{4.102}$$

But by (4.34), as $v(-1+\delta) = -\tilde{\alpha}_P P^{2/3}\delta\hat{\hat{\tilde{f}}}_P(-1+\delta) \to -\infty$,

$$\begin{aligned}
\Big| L_P(y(v(-1+\delta)))\int_{-\infty}^{v(-1+\delta)}\mathrm{Ai}(t)\,dt\Big| &\le \mathrm{const}\,|\tilde{\alpha}_P P^{2/3}\delta\hat{\hat{\tilde{f}}}_P(-1+\delta)|^{-3/4} \\
&\le \mathrm{const}\,\frac{1}{\delta^{3/4}P^{1/2}}
\end{aligned}$$

and also

$$\begin{aligned}
&\Big|\frac{1}{\tilde{\alpha}_P P^{2/3}}\int_{v(-1+\delta)}^{0}\Big(\int_{-\infty}^{v}\mathrm{Ai}(t)\,dt\Big)\,\frac{L_P'(y(v))}{\tilde{U}_P(y(v))}\,dv\Big| \\
&\qquad\qquad \le \frac{\mathrm{const}}{P^{2/3}}\int_{-\tilde{\alpha}_P P^{2/3}\delta\hat{\hat{\tilde{f}}}_P(-1+\delta)}^{0}\frac{dv}{(1+|v|)^{3/4}} \\
&\qquad\qquad \le \mathrm{const}\,\frac{\delta^{1/4}}{P^{1/2}}.
\end{aligned}\tag{4.103}$$

Thus

$$\begin{aligned}
Q_{4,2} &= \int_{u_0}^{u(x_{N,P})} L_N(x(u))\ \mathrm{Ai}(u)\,du \\
&\qquad\times\Big[L_P(-1)\int_{-\infty}^{0}\mathrm{Ai}(v)dv + O\Big(\frac{\delta^{1/4}}{P^{1/2}} + \frac{1}{\delta^{3/4}P^{1/2}}\Big)\Big] \\
&= 2^{1/4}\int_{u_0}^{u(x_{N,P})} L_N(x(u))\ \mathrm{Ai}(u)\,du\int_{-\infty}^{0}\mathrm{Ai}(v)dv + O\Big(\frac{1}{\delta^{3/4}P^{1/2}}\Big)
\end{aligned}$$

where we have again used the estimate in (4.33) as in (4.99). Now as in the analysis of (4.100) we find

$$\begin{aligned}
&\int_{u_0}^{u(x_{N,P})} L_N(x(u))\ \mathrm{Ai}(u)\,du \\
&\qquad = L_N(-1)\int_{u_0}^{u(x_{N,P})}\mathrm{Ai}(u)\,du + \int_{u_0}^{u(x_{N,P})}\big(L_N(x(u)) - L_N(-1)\big)\ \mathrm{Ai}(u)\,du \\
&\qquad = 2^{1/4}\int_{0}^{\infty}\mathrm{Ai}(u)\,du + O(P^{-1/3} + e^{-cN}) + O(N^{-2/3}).
\end{aligned}$$

We conclude that

$$Q_{4,2} = 2^{1/2}\int_{0}^{+\infty}\mathrm{Ai}(u)\,du\int_{-\infty}^{0}\mathrm{Ai}(v)dv + O(P^{-1/3}) + O\Big(\frac{1}{\delta^{3/4}P^{1/2}}\Big).\tag{4.104}$$

Thus as $N, P \dot{\to} \infty$, the second integral in (4.95) behaves as

$$Q_{3,2} = \int_0^{+\infty} \mathrm{Ai}(u)\,du \int_{-\infty}^{u} \mathrm{Ai}(v)dv + O(P^{-1/3}) + O(\delta^{-3/4}P^{-1/2}). \tag{4.105}$$

The first integral in (4.95) is given by (recall (4.81))

$$\begin{aligned}
Q_{3,1} =& \int_{u(-1+\delta)}^{0} L_N(x(u))\,\mathrm{Ai}(u)\,du \int_{v(-1+\delta)}^{\beta(u)} L_P(y(v))\ \mathrm{Ai}(v)\,dv \\
=& L_N(x(u))\Big(\int_{-\infty}^{u} \mathrm{Ai}(t)dt\Big)\Big(\int_{v(-1+\delta)}^{\beta(u)} L_P(y(v))\,\mathrm{Ai}(v)\,dv\Big)\Big|_{u(-1+\delta)}^{0} \\
&+ \frac{1}{\tilde{\alpha}_N N^{2/3}} \int_{u(-1+\delta)}^{0} \Big(\int_{-\infty}^{u} \mathrm{Ai}(t)dt\Big)\frac{L_N'(x(u))}{\tilde{U}_N(x(u))}\,du \\
&\qquad\qquad \times \Big(\int_{v(-1+\delta)}^{\beta(u)} L_P(y(v))\,\mathrm{Ai}(v)\,dv\Big) \\
&- \int_{u(-1+\delta)}^{0} \Big(\int_{-\infty}^{u} \mathrm{Ai}(t)dt\Big)\, L_N(x(u))\, L_P(X(x(u)))\ \mathrm{Ai}(\beta(u))\,\beta'(u)\,du \\
\equiv& Q_{6,1} + Q_{6,2} + Q_{6,3}.
\end{aligned} \tag{4.106}$$

Observe that $u(-1+\delta) \le u \le 0$, and so $\beta(u) \le \text{const} < \infty$ by (4.83),

$$\begin{aligned}
\Big|\int_{v(-1+\delta)}^{\beta(u)} L_P(y(v))\ \mathrm{Ai}(v)\,dv\Big| \le& \Big| L_P(y(v))\Big(\int_{-\infty}^{v} \mathrm{Ai}(t)dt\Big)\Big|_{v(-1+\delta)}^{\beta(u)}\Big| \\
&+ \frac{1}{\tilde{\alpha}_P P^{2/3}}\Big|\int_{v(-1+\delta)}^{\beta(u)} \Big(\int_{-\infty}^{v} \mathrm{Ai}(t)dt\Big)\frac{L_P'(y(v))}{\tilde{U}_P(y(v))}\,dv\Big| \\
\le& \text{const} + O\big((N^{2/3})^{1/4}P^{-2/3}\big) \le \text{const}
\end{aligned} \tag{4.107}$$

where we have used the bound on the integrand (4.34). Thus again using (4.34),

$$|Q_{6,2}| \le \frac{\text{const}}{N^{2/3}} \int_{v(-1+\delta)}^{0} \frac{C}{(1+|u|)^{3/4}}\,du = O(N^{-1/2}). \tag{4.108}$$

Note next that in

$$\begin{aligned}
Q_{6,1} =& L_N(x(0))\Big(\int_{-\infty}^{0} \mathrm{Ai}(t)dt\Big)\cdot \int_{v(-1+\delta)}^{\beta(0)} L_P(y(v))\ \mathrm{Ai}(v)\,dv \\
&- L_N(x(u(-1+\delta)))\Big(\int_{-\infty}^{u(-1+\delta)} \mathrm{Ai}(t)dt\Big)\cdot \int_{v(-1+\delta)}^{\beta(u(-1+\delta))} L_P(y(v))\ \mathrm{Ai}(v)\,dv
\end{aligned}$$

the second term is $O(P^{-1/2})$ by (4.34), (4.41), (4.107). As in (4.102), the first term can be written as

$$\begin{aligned}
L_N(x(0))\Big(\int_{-\infty}^{0} \mathrm{Ai}(t)dt\Big)\cdot L_P(y(\beta(0)))\Big[\Big(\int_{-\infty}^{\beta(0)} \mathrm{Ai}(v)\,dv\Big) + O(\delta^{-3/4}P^{-1/2})\Big] \\
= 2^{1/2}\Big(\int_{-\infty}^{0} \mathrm{Ai}(t)dt\Big)\Big(\int_{-\infty}^{0} \mathrm{Ai}(v)\,dv\Big) + O(\delta^{-3/4}P^{-1/2} + P^{-1/3})
\end{aligned}$$

in view of (4.27) and (4.83). Thus

$$Q_{6,1} = 2^{1/2}\Big(\int_{-\infty}^{0} \mathrm{Ai}(t)dt\Big)^2 + O(\delta^{-3/4}P^{-1/2} + P^{-1/3}). \tag{4.109}$$

Finally,

$$\begin{aligned} Q_{6,3} &= -\int_{u(-1+\delta)}^{0} \Big(\int_{-\infty}^{u} \mathrm{Ai}(t)dt\Big)\, L_N(x(u))\, L_P(X(x(u)))\ \mathrm{Ai}(\beta(u))\,\beta'(u)\,du \\ &= -\int_{u(-1+\delta)}^{0} \Big(\int_{-\infty}^{u} \mathrm{Ai}(t)dt\Big)\, L_N(x(u))\, L_P(X(x(u)))\ \mathrm{Ai}(\beta(u))\,du + O(P^{-1/3}) \end{aligned} \tag{4.110}$$

by (4.84), the uniform boundedness of the remaining factors in the integrand, and since the length of the interval of integration is $O(N^{2/3})$. Denote

$$A(u) \equiv \int_{u(-1+\delta)}^{u} \mathrm{Ai}(\beta(s))\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\,ds.$$

Since

$$\frac{d}{du}A(u) = \Big(\int_{-\infty}^{u} \mathrm{Ai}(t)dt\Big)\ \mathrm{Ai}(\beta(u))$$

the remaining integral in (4.110) after integration by parts becomes

$$\begin{aligned} &-A(u)\, L_N(x(u))\, L_P(X(x(u)))\Big|_{u(-1+\delta)}^{0} \\ &\qquad -\frac{1}{\tilde{\alpha}_N N^{2/3}}\int_{u(-1+\delta)}^{0} A(u)\,\frac{L_N'(x(u))}{\tilde{U}_N(x(u))}\, L_P(X(x(u)))\,du \\ &\qquad -\frac{1}{\tilde{\alpha}_N N^{2/3}}\int_{u(-1+\delta)}^{0} A(u)\, L_N(x(u))\frac{L_P'(X(x(u)))\,X'(x(u))}{\tilde{U}_P(x(u))}\,du. \end{aligned} \tag{4.111}$$

We need the following result.

**Lemma 4.6.** *As* $N,P\dot{\to}\infty$*, we have uniformly for* $u(-1+\delta)\le u\le 0$

$$A(u) = \int_{-\infty}^{u} \mathrm{Ai}(s)\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\,ds + O(\delta^{1/2}) + O(\delta^{-3/2}N^{-1}). \tag{4.112}$$

*Proof.* For $u(-1+\delta)\le u\le -1$ we recall (4.76)

$$u(-1+\delta) = -\tilde{\alpha}_N\delta\hat{\tilde{f}}_N(-1+\delta) = -c\delta N^{2/3}(1+o(1)), \qquad c>0. \tag{4.113}$$

For $u\le 0$

$$\begin{aligned} &\Big|\int_{u(-1+\delta)}^{u} \mathrm{Ai}(\beta(s))\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\,ds - \int_{-\infty}^{u} \mathrm{Ai}(s)\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\,ds\Big| \\ &\qquad \le \Big|\int_{-\infty}^{u(-1+\delta)} \mathrm{Ai}(s)\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\,ds\Big| \\ &\qquad\quad + \Big|\int_{u(-1+\delta)}^{u} \big[\,\mathrm{Ai}(\beta(s)) - \mathrm{Ai}(s)\big]\,\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\,ds\Big|. \end{aligned} \tag{4.114}$$

Note that

$$\int_{-\infty}^{u} \mathrm{Ai}(s)\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\,ds = \frac{1}{2}\Big(\int_{-\infty}^{u} \mathrm{Ai}(s)\,ds\Big)^2 \tag{4.115}$$

and so, by (4.34), (4.113), the first term in the RHS in (4.114) equals

$$\frac{1}{2}\Big(\int_{-\infty}^{u(-1+\delta)} \mathrm{Ai}(s)\,ds\Big)^2 = O((\delta N^{2/3})^{-3/2}) = O(\delta^{-3/2}N^{-1}).$$

In view of (4.22), (4.83), we bound the second term in (4.114) by

$$\begin{aligned}\int_{u(-1+\delta)}^{u} &\big|\,\mathrm{Ai}(\beta(s)) - \mathrm{Ai}(s)\big| \cdot \Big|\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big|\, ds \\ &\leq \int_{u(-1+\delta)}^{0} \Big(\sup_{t\in[\beta(s),s]} |\,\mathrm{Ai}'(t)|\Big) \cdot |\beta(s)-(s)| \cdot \Big|\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big|\, ds \\ &\leq \mathrm{const}\int_{u(-1+\delta)}^{1} (1+|s|)^{1/4}\Big[O(P^{-1/3}) + |s|O(P^{-1})\Big]\frac{ds}{1+|s|^{3/4}} \\ &\leq O(P^{-1/3}(\delta N^{2/3})^{1/2}) + O(P^{-1}(\delta N^{2/3})^{3/2}) = O(\delta^{1/2})\end{aligned}$$

uniformly for $u(-1+\delta) \leq u \leq 0$. □

From (4.115), Lemma 4.6, (4.111), and the fact that $A(u(-1+\delta)) = 0$, we conclude that

$$Q_{6,3} = -\frac{2^{1/2}}{2}\Big(\int_{-\infty}^{0} \mathrm{Ai}(s)\,ds\Big)^2 + O(\delta^{1/2}) + O(\delta^{-3/2}N^{-1}) \tag{4.116}$$

again by the uniform boundedness of the terms in the integrands, the integrability of $\big(\int_{-\infty}^{u} \mathrm{Ai}(s)\,ds\big)^2 \sim |u|^{-3/2}$, $u \to -\infty$, and the fact that $u(-1+\delta) = O(N^{2/3})$. By (4.116), (4.108), (4.109), (4.115) we see that (4.106) is given by

$$\begin{aligned}Q_{3,1} = &\int_{-\infty}^{0} \mathrm{Ai}(s)\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\, ds \\ &+ O(\delta^{1/2}) + O(\delta^{-3/4}N^{-1/2}) + O(\delta^{-3/2}N^{-1}) + O(P^{-1/3}).\end{aligned} \tag{4.117}$$

Thus by (4.95), (4.117), (4.105), the Ai × Ai integral in (4.94) equals $(-1)^{N+P}\frac{(c_N c_P)^{1/2}}{(\tilde{\alpha}_N\tilde{\alpha}_P)^{3/4}(NP)^{1/2}}$ times

$$\begin{aligned}&\Big(\int_{-\infty}^{0} + \int_{0}^{+\infty}\Big)\mathrm{Ai}(s)\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\, ds + O(P^{-1/3}) + O\bigg(\delta^{1/2} + \frac{1}{\delta^{3/4}P^{1/2}} + \frac{1}{\delta^{3/2}P}\bigg) \\ &= \int_{-\infty}^{+\infty} \mathrm{Ai}(s)\Big(\int_{-\infty}^{s} \mathrm{Ai}(t)dt\Big)\, ds + O(P^{-1/3}) + O\bigg(\delta^{1/2} + \frac{1}{\delta^{3/4}P^{1/2}} + \frac{1}{\delta^{3/2}P}\bigg) \\ &= \frac{1}{2}\Big(\int_{-\infty}^{+\infty} \mathrm{Ai}(s)\,dx\Big)^2 + O(P^{-1/3}) + O\bigg(\delta^{1/2} + \frac{1}{\delta^{3/4}P^{1/2}} + \frac{1}{\delta^{3/2}P}\bigg) \\ &= \frac{1}{2}\bigg[1 + O(P^{-1/3}) + O\bigg(\delta^{1/2} + \frac{1}{\delta^{3/4}P^{1/2}} + \frac{1}{\delta^{3/2}P}\bigg)\bigg]\end{aligned} \tag{4.118}$$

by (4.115), and since $\int_{-\infty}^{+\infty}\mathrm{Ai}(t)dt = 1$ (see [AbSt]). Thus from (4.12)(2), (4.62)

$$\begin{aligned} J_{44} = & J'_{44} + J''_{44} = (-1)^{N+P}\frac{(c_N c_P)^{1/2}}{(NP)^{1/2}} \\ & \times \Bigg[\frac{1}{4m} + O(P^{-1/3}) + O(P^{-1/(2m)}) \\ & \qquad\qquad + O\Big(\delta^{1/2} + \delta^{-3/4}P^{-1/2} + \delta^{-3/2}P^{-1}\Big) \qquad \text{by (4.118)} \\ & \quad + O(N^{-1/3}) \qquad \text{by (4.66)} \\ & \quad + O(\delta^{1/2}) + O(P^{-1/3}) \qquad \text{by (4.94)}\Bigg] \\ = & \frac{(-1)^{N+P}}{4m}\frac{(c_N c_P)^{1/2}}{(NP)^{1/2}}\Bigg[1 + O(P^{-1/3}) + O(P^{-1/(2m)}) \\ & \qquad\qquad + O\Big(\delta^{1/2} + \delta^{-3/4}P^{-1/2} + \delta^{-3/2}P^{-1}\Big)\Bigg]. \end{aligned}$$

By this estimate and (4.57), (4.58), (4.59), we find finally that

$$\begin{aligned} J_4 = \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}}\Bigg[&\frac{(-1)^{N+P}}{4m} + \frac{(-1)^N}{2m} + O(P^{-1/3}) + O(P^{-1/(2m)}) \\ & + O\Big(\delta^{1/2} + P^{-1/2}\delta^{-3/4} + P^{-1}\delta^{-3/2}\Big)\Bigg]. \end{aligned} \tag{4.119}$$

In a similar way we obtain

$$\begin{aligned} J_2 \equiv & \int_{1-\delta}^{1+\delta} c_N\phi_N(c_N x + d_N)\Big(\int_{X(x)}^{1+\delta} c_P\phi_P(c_P x + d_P)\Big)\,dx \\ = & \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}}\Bigg[\frac{1}{4m} + O(P^{-1/3}) + O(P^{-1/(2m)}) \\ & + O\Big(\delta^{1/2} + P^{-1/2}\delta^{-3/4} + P^{-1}\delta^{-3/2}\Big)\Bigg]. \end{aligned} \tag{4.120}$$

(Note that the integral leading to the term $\frac{(-1)^N}{2m}$ in (4.119), is not present for $J_2$.)

It remains to consider $J_3$. Again assume that $N, P \to \infty$. By (4.56)

$$\begin{aligned} J_3 = & c_N c_P \int_{I_3}\phi_N(c_N x + d_N)\,dx \int_{X(x)}^{+\infty}\phi_P(c_P y + d_P)\,dy \\ = & c_N c_P \int_{I_3}\phi_N(c_N x + d_N)\,dx\Big(\int_{X(x)}^{1-\delta} + \int_{I_2} + \int_{I_1}\Big)\phi_P(c_P y + d_P)\,dy \\ \equiv & J_{33} + J_{32} + J_{31}. \end{aligned}$$

By (4.18), (4.17), (4.16)

$$|J_{32}| + |J_{31}| = c_N^{1/2}N^{-1}\,O\big((c_P/P)^{1/2} + (c_P/P)^{1/2}\,e^{-Pc_3}\big) = \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}}\,O\big(N^{-1/2}\big).$$

To analyze $J_{33}$ we first introduce $x^{\pm}_{N,P} \in I_3$ such that

$$X(x^{\pm}_{N,P}) \equiv x^{\pm}_{N,P}c_N/c_P + (d_N - d_P)/c_P = \pm(1-\delta)$$

which in view of (4.54), implies that for $N, P \dot{\to} \infty$,

$$X(x) \in I_3, \qquad x \in I_3' \equiv [x_{N,P}^-, x_{N,P}^+] \subset I_3. \tag{4.121}$$

Set $I_3'' \equiv I_3 \setminus I_3'$ and note that by Lemma 4.4

$$x_{N,P}^{\pm} = \pm(1-\delta) + O(P^{-1}). \tag{4.122}$$

Then

$$\begin{aligned} J_{33} =& c_N c_P \Big( \int_{I_3'} + \int_{I_3''} \Big) \phi_N(c_N x + d_N)\, dx \int_{X(x)}^{1-\delta} \phi_P(c_P y + d_P)\, dy \\ &\equiv J_{33}' + J_{33}''. \end{aligned}$$

Now

$$|J_{33}''| \le C \int_{I_3''} \frac{c_N}{c_N^{1/2}} \frac{c_N^{1/2}}{N^{1/2}}\, dx = C c_N\, N^{-1/2}\, |I_3''| = \frac{(c_N c_P)^{1/2}}{(NP)^{1/2}}\, O\big(P^{-1/2}\big) \tag{4.123}$$

by (4.63), (4.15), (4.122).

Now taking into account (4.121) and using (4.13)

$$\begin{aligned} J_{33}' =& \frac{2}{\pi} \sqrt{c_N c_P} \int_{I_3'} \frac{dx}{|1-x^2|^{1/4}} \Big[ \Big( \cos f_N^+(x) \Big) \Big( 1 + O(N^{-1} \Big) + \Big( \sin f_N^-(x) \Big) O(N^{-1}) \Big] \\ & \times \int_{X(x)}^{1-\delta} \frac{dy}{|1-y^2|^{1/4}} \Big[ \Big( \cos f_P^+(y) \Big) \Big( 1 + O(P^{-1} \Big) + \Big( \sin f_P^-(y) \Big) O(P^{-1}) \Big] \end{aligned}$$

where

$$f_N^{\pm}(x) \equiv \frac{N}{2} \int_1^x |1-y^2|^{1/2} h_N(y) dy \pm \frac{1}{2} \arcsin x \tag{4.124}$$

and $h_N$ satisfies (4.7) and (2.7).

Clearly a product of two terms, one of which has a factor $O(N^{-1})$ and the other $O(P^{-1})$, gives rise to a smaller order contribution $\frac{\sqrt{c_N c_P}}{\sqrt{NP}} O(N^{-1})$. Consider next the integral

$$\begin{aligned} & \Big| \int_{I_3'} \frac{dx}{|1-x^2|^{1/4}} \Big( \sin f_N^-(x) \Big) O(N^{-1}) \int_{X(x)}^{1-\delta} \frac{dy}{|1-y^2|^{1/4}} \Big( \cos f_P^+(y) \Big) \Big| \\ & \quad \le O(N^{-1}) \int_{I_3'} \frac{dx}{|1-x^2|^{1/4}} \Big| \int_{X(x)}^{1-\delta} \frac{dy}{|1-y^2|^{1/4}} \Big( \cos f_P^+(y) \Big) \Big| \\ & \quad \le O(N^{-1}) \Big( \int_{I_3'} \frac{dx}{|1-x^2|^{1/4}} \Big) \max_{-1+\delta \le x_1 \le 1-\delta} \Big| \int_{x_1}^{1-\delta} \frac{dy}{|1-y^2|^{1/4}} \Big( \cos f_P^+(y) \Big) \Big| \\ & \quad = O((NP)^{-1}), \end{aligned}$$

where we have used (4.21) in the last step. Thus

$$\begin{aligned} & \frac{2}{\pi} \sqrt{c_N c_P} \int_{I_3'} \frac{dx}{|1-x^2|^{1/4}} \Big( \sin f_N^-(x) \Big) O(N^{-1}) \int_{X(x)}^{1-\delta} \frac{dy}{|1-y^2|^{1/4}} \Big( \cos f_P^+(y) \Big) \\ & \qquad = \frac{\sqrt{c_N c_P}}{\sqrt{NP}}\, O(N^{-1}). \end{aligned}$$

Similarly, interchanging the order of integration as in (4.73), we find

$$\begin{aligned}
&\frac{2}{\pi}\sqrt{c_Nc_P}\int_{I_3'}\cos f_N^+(x)\,\frac{dx}{|1-x^2|^{1/4}}\int_{X(x)}^{1-\delta}\Big(\sin f_P^-(y)\Big)\,O(P^{-1})\,\frac{dy}{|1-y^2|^{1/4}}\\
&\qquad=\frac{2}{\pi}\sqrt{c_Nc_P}\int_{-1+\delta}^{1-\delta}\frac{\big(\sin f_P^-(y)\big)\,O(P^{-1})}{|1-y^2|^{1/4}}\,dy\int_{-1+\delta}^{X^{-1}(y)}\frac{\cos f_N^+(x)}{|1-x^2|^{1/4}}\,dx\\
&\qquad=\frac{\sqrt{c_Nc_P}}{\sqrt{NP}}\,O(N^{-1}).
\end{aligned}$$

Thus

$$J_3=\frac{2}{\pi}\sqrt{c_Nc_P}\int_{I_3'}\frac{\cos f_N^+(x)}{|1-x^2|^{1/4}}\,dx\int_{X(x)}^{1-\delta}\frac{\cos f_P^+(y)}{|1-y^2|^{1/4}}\,dy+\frac{\sqrt{c_Nc_P}}{\sqrt{NP}}\,O(N^{-1/2}). \tag{4.125}$$

Note that, in view of (4.7), for a fixed $\delta>0$, there exists $\eta_\delta>0$ such that as $N\to\infty$

$$(f_N^\pm(x))'=\frac{N}{2}|1-x^2|^{1/2}h_N(x)\pm\frac{1}{2|1-x^2|^{1/2}}\geq N\eta_\delta \tag{4.126}$$

uniformly for $x\in I_3'$. The integral in (4.125) is given by $\frac{2}{\pi}\sqrt{c_Nc_P}$ times

$$\begin{aligned}
&\int_{I_3'}\frac{\cos f_N^+(x)}{|1-x^2|^{1/4}}\,dx\bigg[\frac{\sin f_P^+(y)}{(f_P^+(y))'|1-y^2|^{1/4}}\bigg|_{X(x)}^{1-\delta}\\
&\qquad\qquad-\int_{X(x)}^{1-\delta}(\sin f_P^+(y))\Big(\frac{1}{|1-y^2|^{1/4}(f_P^+(y))'}\Big)'\,dy\bigg]\\
&\quad=-\int_{I_3'}\frac{\cos f_N^+(x)}{|1-x^2|^{1/4}}\,\frac{\sin f_P^+(X(x))}{f_P^{+\prime}(X(x))|1-(X(x))^2|^{1/4}}\,dx+O((NP)^{-1})
\end{aligned}\tag{4.127}$$

in view of (4.126), since we can integrate by parts in $x$ once more for the boundary term at $y=1-\delta$, and also integrate by parts again in the integral term $\int_{X(x)}^{1-\delta}\cdots$. Note next that by Lemma 4.4 we have again uniformly for $x\in I_3'$

$$|X(x)-x|=O(P^{-1}).$$

Also, uniformly for $x\in I_3$, $|f_P^{+\prime\prime}(x)|\leq\text{const}\cdot P$ and $|1-x^2|^{-1}\leq c$. Thus, as $X(I_3')\subset I_3$, we conclude that

$$\left|\frac{1}{f_P^{+\prime}(X(x))|1-(X(x))^2|^{1/4}}-\frac{1}{f_P^{+\prime}(x)|1-x^2|^{1/4}}\right|\leq\frac{\text{const}}{P^2}$$

and hence using (4.127) we find

$$J_3=-\frac{2}{\pi}\sqrt{c_Nc_P}\int_{I_3'}\frac{\cos f_N^+(x)\,\sin f_P^+(X(x))}{f_P^{+\prime}(x)|1-x^2|^{1/2}}\,dx+\frac{\sqrt{c_Nc_P}}{\sqrt{NP}}\,O(N^{-1/2}). \tag{4.128}$$

Recalling (4.126), (4.7) and (2.7), we derive from (4.128)

$$J_3=-\frac{4}{\pi}\frac{\sqrt{c_Nc_P}}{\sqrt{NP}}\int_{I_3'}\frac{\cos f_N^+(x)\,\sin f_P^+(X(x))}{(1-x^2)\,h(x)}\,dx+\frac{\sqrt{c_Nc_P}}{\sqrt{NP}}\,O(N^{-1/(2m)}). \tag{4.129}$$

Now the integral above is given by

(4.130)

$$\begin{aligned}\int_{I_3'} \frac{\cos f_N^+(x)\,\sin f_P^+(X(x))}{(1-x^2)\,h(x)}\,dx =&\frac{1}{2}\int_{I_3'} \frac{\sin\left[f_P^+(X(x))+f_N^+(x)\right]}{(1-x^2)\,h(x)}\,dx\\ &+\frac{1}{2}\int_{I_3'} \frac{\sin\left[f_P^+(X(x))-f_N^+(x)\right]}{(1-x^2)\,h(x)}\,dx \equiv Q_{7,1}+Q_{7,2}.\end{aligned}$$

Integrating by parts in $Q_{7,1}$ in the same way as in (4.127) and noting that for sufficiently large $N$, by (4.126),

$$f_P^{+\prime}(X(x))\,X'(x)+f_N^{+\prime}(x)\geq N\tilde{\eta}_\delta$$

uniformly for $x\in I_3'$, we conclude that

$$|Q_{7,1}|\leq O(N^{-1}).$$

Finally consider $Q_{7,2}$. Introduce the convenient notation

$$\theta(x)\equiv\frac{1}{2}\int_0^x |1-t^2|^{1/2}\,h(t)\,dt \tag{4.131}$$

and note that $\theta(1)=\pi/2$ by (2.5), (2.7), the fact that $h(t)$ is even (see (2.8)) and the normalization condition $\int_{-1}^1 \psi_N^{(\mathrm{eq})}(x)\,dx=1$.

**Lemma 4.7.** *Uniformly for $x\in I_3'$ as $N,P\dot{\to}\infty$,*

$$f_P^+(X(x))-f_N^+(x)=(N-P)\left[\frac{1}{2m}\,x\theta'(x)-\theta(x)+\frac{\pi}{2}\right]+O(N^{-1/(2m)}). \tag{4.132}$$

*Proof.* We have

$$f_P^+(X(x))-f_N^+(x)=[f_P^+(X(x))-f_P^+(x)]+[f_P^+(x)-f_N^+(x)]\equiv F_1+F_2.$$

By (4.124)

$$F_2=\frac{1}{2}\int_1^x |1-y^2|^{1/2}[Ph_P(x)-Nh_N(y)]dy.$$

Note that by (2.7) with $q=2m$ and (4.51)

$$\begin{aligned}Ph_P(x)-Nh_N(x)&=P(h_P(x)-h_N(x))+(P-N)h_N(x)\\ &=P\sum_{k=1}^{2m}\Big[h_{(k)}(x)\cdot(P^{-k/(2m)}-N^{-k/(2m)})\Big]+P\,O(P^{-1-1/(2m)})\\ &\quad+(P-N)[h(x)+O(N^{-1/(2m)})]\\ &=(P-N)h(x)+O(N^{-1/(2m)})\end{aligned}$$

uniformly for $-1\leq x\leq 1$. Hence, uniformly for $x\in I_3'$,

$$\begin{aligned}F_2=&(P-N)\frac{1}{2}\int_1^x |1-y^2|^{1/2}h(y)dy+O(N^{-1/(2m)})\\ =&-(N-P)\Big[\theta(x)-\frac{1}{2}\int_0^1 |1-y^2|^{1/2}h(y)dy\Big]+O(N^{-1/(2m)})\\ =&-(N-P)\Big[\theta(x)-\frac{\pi}{2}\Big]+O(N^{-1/(2m)}),\end{aligned}$$

since $\theta(1)=\pi/2$. Next

$$F_1=f_P^{+\prime}(x)\,(X(x)-x)+f_P^{+\prime\prime}(\xi(x))\,(X(x)-x)^2/2$$

for some $\xi(x)$ between $x$ and $X(x)$. Note that by (4.53) and Lemma 4.4

$$X(x) - x = (c_N/c_P - 1)x + (d_N - d_P)/c_P = \frac{1}{2m}\frac{N-P}{P}\,x + O(P^{-1-1/(2m)}).$$

Also by (4.126) and (2.7), $|f_P^{+\prime\prime}(x)| \le \mathrm{const} \cdot P$ uniformly for $x \in I_3$. But

$$f_P^{+\prime}(x) = \frac{P}{2}|1-x^2|^{1/2}h(x)\Big[1 + O(P^{-1/(2m)}) + O(P^{-1})\Big]$$

and hence by (4.131)

$$F_1 = \frac{1}{2}\frac{N-P}{2m}\,x|1-x^2|^{1/2}h(x) + O(P^{-1/(2m)}) = \frac{N-P}{2m}\,x\theta'(x) + O(P^{-1/(2m)})$$

which completes the proof of the lemma. □

We need the following fortunate and remarkable fact.

**Lemma 4.8.** *For any $V(x) = \kappa_{2m}x^{2m} + \cdots$ the function $\theta$ defined by* (4.131) *solves the linear first order ODE*

$$\theta(x) - \frac{1}{2m}x\theta'(x) = \arcsin x, \qquad -1 \le x \le 1.$$

*Proof.* See Remark 6.1 in Subsection 6.3. □

Thus from (4.130), (4.132)

$$J_3 = -\frac{2}{\pi}\frac{\sqrt{c_N c_P}}{\sqrt{NP}}\int_{I_3'} \frac{\sin\big[(P-N)\arcsin x + (N-P)\frac{\pi}{2})\big]}{(1-x^2)\,h(x)}\,dx + \frac{\sqrt{c_N c_P}}{\sqrt{NP}}\,O(N^{-1/(2m)}).$$

Note that by (4.122) we can now replace $I_3'$ with $I_3 \equiv [-1+\delta, 1-\delta]$ introducing an error $\frac{\sqrt{c_N c_P}}{\sqrt{NP}}\,O(N^{-1})$. Now using the formula for the sine of a sum and noting that $\frac{\sin[(P-N)\arcsin x]}{(1-x^2)\,h(x)}$ is an odd function we find

$$\begin{aligned} J_3 = &-\frac{2}{\pi}\frac{\sqrt{c_N c_P}}{\sqrt{NP}}\,\sin\Big(\frac{(N-P)\pi}{2}\Big)\int_{-1+\delta}^{1-\delta}\frac{\cos\big[(P-N)\arcsin x\big]}{(1-x^2)\,h(x)}\,dx \\ &+ \frac{\sqrt{c_N c_P}}{\sqrt{NP}}\,O(N^{-1/(2m)}). \end{aligned}$$

Assume $N-P$ is even. Then $J_3 = \frac{\sqrt{c_N c_P}}{\sqrt{NP}}\,O(N^{-1/(2m)})$. Assume $N-P$ is odd. Then

$$\frac{\cos\big[(P-N)\arcsin x\big]}{(1-x^2)\,h(x)} = O\Big(\frac{1}{|1-x^2|^{1/2}}\Big), \qquad x \to \pm(1-0)$$

and hence for $N-P$ odd

$$\begin{aligned} J_3 = &-\frac{2}{\pi}\frac{\sqrt{c_N c_P}}{\sqrt{NP}}\,\sin\Big(\frac{(N-P)\pi}{2}\Big)\int_{-1}^{1}\frac{\cos\big[(N-P)\arcsin x\big]}{(1-x^2)\,h(x)}\,dx \\ &+ \frac{\sqrt{c_N c_P}}{\sqrt{NP}}\Big(O(N^{-1/(2m)}) + O(\delta^{1/2})\Big). \end{aligned} \tag{4.133}$$

By (4.47), (4.52), (4.56), as $N, P \dot{\to} \infty$,
(4.134)

$$\begin{aligned}
(\phi_N, \epsilon\phi_P) =& \frac{1}{2}\Big(\int_{-\infty}^{+\infty} \phi_N(x)\,dx\Big)\Big(\int_{-\infty}^{+\infty} \phi_P(y)\,dy\Big) \\
&- \Big[c_N c_P \Big(\int_{I_1} + \int_{I_5}\Big)\phi_N(c_N x + d_N)\,dx \int_{X(x)}^{+\infty} \phi_P(c_P y + d_P)\,dy \\
&\qquad + \Big(J_4 + J_2\Big) + J_3\Big] \\
=& \frac{\sqrt{c_N c_P}}{\sqrt{NP}} \Big\{ \frac{1}{2}\,\frac{1}{2m}\,(1 + (-1)^N + O(N^{-1/2}) + O(N^{-1/(2m)})) \\
&\qquad \times (1 + (-1)^P + O(P^{-1/2}) + O(P^{-1/(2m)})) \qquad \text{by (4.48)} \\
&\quad - \Big[ O(e^{-c_3 N}) \qquad \text{by (4.55)} \\
&\qquad + \frac{(-1)^{N+P}}{4m} + \frac{(-1)^N}{2m} + \frac{1}{4m} \\
&\qquad\quad + O(P^{-1/3}) + O(P^{-1/(2m)}) \\
&\qquad\quad + O\Big(\delta^{1/2} + P^{-1/2}\delta^{-3/4} + P^{-1}\delta^{-3/2}\Big) \quad \text{by (4.119), (4.120)} \\
&\qquad - \sin\Big(\frac{(N-P)\pi}{2}\Big)\,\frac{2}{\pi}\int_{-1}^{1} \frac{\cos\big[(N-P)\arcsin x\big]}{(1-x^2)\,h(x)}\,dx \\
&\qquad\quad + O(N^{-1/(2m)}) + O(\delta^{1/2})\Big]\Big\} \quad \text{by (4.133)} \\
=& \frac{\sqrt{c_N c_P}}{\sqrt{NP}} \Big[ \frac{(-1)^P - (-1)^N}{4m} + I(N-P) \\
&\qquad + O(P^{-1/3}) + O(P^{-1/(2m)}) + O\Big(\delta^{1/2} + P^{-1/2}\delta^{-3/4} + P^{-1}\delta^{-3/2}\Big)\Big]
\end{aligned}$$

where

$$I(q) \equiv \begin{cases} \sin\Big(\frac{q\pi}{2}\Big)\,\frac{2}{\pi}\int_{-1}^{1} \frac{\cos(q\arcsin x)}{(1-x^2)\,h(x)}\,dx, & q \text{ odd} \\ 0, & q \text{ even.} \end{cases}$$

As $(\phi_N, \epsilon\phi_P) = -(\epsilon\phi_N, \phi_P) = -(\phi_P, \epsilon\phi_N)$ we see that (4.134) is true also if $N < P$. Hence as $N, P \dot{\to} \infty$

$$(\epsilon\phi_N, \phi_P) = \frac{\sqrt{c_N c_P}}{\sqrt{NP}} \begin{cases} \frac{(-1)^N}{2m} - I(N-P) + O(\delta^{1/2}) + o_\delta(1), & N-P \text{ odd} \\ O(\delta^{1/2}) + o_\delta(1), & N-P \text{ even,} \end{cases}$$

where $o_\delta(1) \equiv O(P^{-1/3}) + O(P^{-1/(2m)}) + O(P^{-1/2}\delta^{-3/4} + P^{-1}\delta^{-3/2})$. Note that some of the smaller order terms in $o_\delta(1)$ are proportional to $\frac{1}{\delta^\alpha}$ for some $\alpha > 0$. Nevertheless we see that e.g. for $N - P$ odd

$$\begin{aligned}
-O(\delta^{1/2}) \le& \liminf_{N,P\dot{\to}\infty} \Big[\frac{\sqrt{NP}}{\sqrt{c_N c_P}}(\epsilon\phi_N, \phi_P) - \Big(\frac{(-1)^N}{2m} - I(N-P)\Big)\Big] \\
&\le \limsup_{N,P\dot{\to}\infty} \Big[\frac{\sqrt{NP}}{\sqrt{c_N c_P}}(\epsilon\phi_N, \phi_P) - \Big(\frac{(-1)^N}{2m} - I(N-P)\Big)\Big] \le O(\delta^{1/2}).
\end{aligned}$$

Letting $\delta \to 0$ we conclude that $\liminf = \limsup$ and hence

$$(\epsilon\phi_N, \phi_P) = \frac{\sqrt{c_N c_P}}{\sqrt{NP}} \left[ \frac{(-1)^N}{2m} - I(N-P) + o(1) \right] \tag{4.135}$$

as $N, P \dot{\to} \infty$, $N - P$ odd. Similarly

$$(\epsilon\phi_N, \phi_P) = \frac{\sqrt{c_N c_P}}{\sqrt{NP}}\, o(1) \tag{4.136}$$

as $N, P \dot{\to} \infty$, $N - P$ even. This completes the proof of Theorem 2.2.

*Remark* 4.3. As noted above, the proof of Theorem 2.2 involves a $\liminf/\limsup$ argument as $N, P \dot{\to} \infty$, followed by a limit as $\delta \to 0$. However, as noted at the beginning of Subsection 4.2.1, the error estimates from [DKMVZ2] in (4.6) et seq., are uniform only for $\delta$ in compact subsets of $(0, \delta_0]$, and the reader may be concerned that the $O(\delta^{1/2})$ term in the $\liminf/\limsup$ argument above, in fact depends on constants, say, that blow up as $\delta \to 0$. But the reader may easily check that all the error terms that arise in the proof of Theorem 2.2, and that could blow up as $\delta \to 0$, are always multiplied by negative power of $N$ or $P$, and hence vanish in the $\liminf/\limsup$ argument: all that remains are $O(\delta^{1/2})$ terms that arise from the evaluation of integrals with explicit $\delta$-independent integrands (see e.g. the proof of (4.133) above), and do not blow up as $\delta \to 0$.

*Remark* 4.4. In the case $m = 1$ the OP's are just the Hermite polynomials. They have the exceptional property that if $N$ is odd then $\epsilon\phi_N$ is a polynomial of degree $N-1$ times $e^{-x^2/2}$, and hence $(\epsilon\phi_N, \phi_P)$ is identically zero for $N$ odd and $N < P$. However, we have

$$\int_{-1}^{1} \frac{\cos\big(q \arcsin x\big)}{1-x^2}\, dx = \begin{cases} \pi, & q = 1, 5, 9, \cdots \\ -\pi, & q = 3, 7, \cdots, \end{cases}$$

and for $m = 1$, $h(x) \equiv 4$ by (2.8). A simple calculation now shows that the leading coefficient in (4.135) is zero for $N$ odd, $N < P$, $P$ even, so that the calculations match.

*Remark* 4.5. The following (nonrigorous) argument is consistent with our asymptotic formulae. We know by Theorem 2.1 that for any polynomial $V$ the matrix $D_\infty$ is banded and looks asymptotically like the product of a Toeplitz matrix, say $\tilde{D}_\infty$, whose diagonals are given by certain binomial coefficients times the diagonal matrix $\tilde{T}_\infty \equiv \mathrm{diag}((m\kappa_{2m} b_j^{2m-1})_{j\geq 0})$, $b_0 \equiv 1$ (see (2.2)). Take a large enough even $N$ and let $\tilde{D}_N$ be an $N \times N$ section of $\tilde{D}_\infty$, i.e. $(\tilde{D}_N)_{i,j} = (\tilde{D}_\infty)_{i,j}$, $0 \leq i, j \leq N-1$. Also set

$$\tilde{T}_N \equiv \mathrm{diag}((m\kappa_{2m} b_j^{2m-1})_{j=0}^{N-1}, \qquad b_0 \equiv 1.$$

Consider a submatrix in the middle of the matrix $(\tilde{D}_N \tilde{T}_N)^{-1}$ near its diagonal which is small compared to $N$. For large $N$, in view of Theorem 2.5(i), we would expect this small submatrix to look like the corresponding submatrix of the matrix $\epsilon_\infty$.

Let us take, e.g., $m = 2$ and $V(x) = x^4$. Then $n = 2m - 1 = 3$. In this case the Toeplitz matrix $\tilde{D}_N$ has diagonals (cf. Theorem 2.1)

$$(\cdots \quad 0 \quad 1 \quad 0 \quad 3 \quad 0 \quad -3 \quad 0 \quad -1 \quad 0 \quad \cdots).$$

In view of the preceding discussion, we would expect that for large enough even $N$ the submatrix of $\tilde{T}_N^{-1}\tilde{D}_N^{-1}$ formed by the rows $M-n, \cdots, M-1$ and the columns $M, \cdots, M+n-1$, $M$ even and $M \sim N/2$, to look like the submatrix of $\epsilon_\infty$ formed by the same rows and columns. In other words, we would expect that this small submatrix of $\tilde{D}_N^{-1}$ would look like the small submatrix of $\tilde{T}_N\epsilon_N$ located at the same position. (Recall that the rows and columns in $\tilde{D}_N^{-1}$ and $\tilde{T}_N\epsilon_N$ are enumerated from 0 to $N-1$ so that the submatrix chosen above corresponds to $B_{12}$ defined in (1.34) about the center $(M, M)$.)

The overall coefficient multiplying the elements in the middle of the matrix $T_N\epsilon_N$ is the same as (2.14), that is $m\kappa_{2m}b_M^{2m-1}\,c_M M^{-1} = \frac{2(m!)^2}{(2m)!}(1+o(1))$, for $M$ large enough. Using Maple software we have computed the matrix $\tilde{D}_N^{-1}$ for $N = 20$. It turns out that the submatrix of $\tilde{D}_N^{-1}$ with rows $7, 8, 9$ and columns $10, 11, 12$ (here $M = N/2 = 10$, $n = 3$) and the corresponding submatrix of $\tilde{T}_N\epsilon_N$ computed using Theorem 2.2 (here $h(x) = \frac{8}{3}(1+2x^2)$ by (2.8)) are given by

$$\begin{pmatrix} -0.01630 & 0 & 0.00435 \\ 0 & 0.15078 & 0 \\ 0.06113 & 0 & -0.01630 \end{pmatrix}, \qquad \begin{pmatrix} -0.01635 & 0 & 0.00438 \\ 0 & 0.15032 & 0 \\ 0.06100 & 0 & -0.01635 \end{pmatrix},$$

respectively. We see that the corresponding nonzero elements differ by at most 0.8%. For $N = 40$, $M = N/2$, the maximal difference is already less than 0.002%. We have also done similar computations for $m = 3, 4$, $n = 2m-1$ again for $N = 40$, $M = N/2$. It turns out that the nonzero elements in the small blocks computed using the matrix inversion and Theorem 2.2 as described above differ by at most 0.005%, 0.02% for $m = 3, 4$, respectively.

4.3. **Convergence of derivatives and integrals of the Christoffel–Darboux kernel for weights $e^{-V(x)}$, $V(x) = \kappa_{2m}x^{2m} + \cdots$, $\kappa_{2m} > 0$.** We start with the convergence of the derivatives in the 12 entries of the kernels $K_{N,\beta}$, $\beta = 1, 4$. Our main results in this direction are (4.150), (4.151) and their Corollaries (4.157), (4.158) below. After that we prove the convergence of the integrals in the 21 entries of the kernels $K_{N,\beta}$, $\beta = 1, 4$. The main result for the integrals can be found in (4.171) below.

4.3.1. *Derivatives.* Fix $L_0 > 0$. For $r$, $\xi$, $\eta$ in the compact set $|\xi|$, $|\eta|$, $|r| \le 2L_0$, define

$$D_{r,N}(\xi,\eta) \equiv \frac{1}{q_N}K_N\left(r + \frac{\xi}{q_N}, r + \frac{\eta}{q_N}\right). \tag{4.137}$$

In what follows, $q_N$ is *any* sequence of numbers with the property

$$q_N = \frac{Nh_N(0)}{2\pi c_N}\left(1 + O(N^{-\alpha})\right) \tag{4.138}$$

for some $0 < \alpha \le 1$, as $N \to \infty$.

*Remark* 4.6. In the calculations that follow we use formulae for $D_{r,N}(\xi,\eta)$ based on (1.7) that holds for $\xi \ne \eta$. But $D_{r,N}(\xi,\eta)$ is continuous in $\xi, \eta$ (see e.g. (1.5)), and it will be clear from the calculations that we can obtain analogous formulae for $D_{r,N}$ on the diagonal by taking the limit $\eta \to \xi$ in the formulae below.

Eventually in (4.137), (4.138) we will take $q_N = R_{N,1,1}(r), R_{N/2,1,4}(r)$ for $\beta = 1, 4$ respectively, but at this stage asymptotics for $R_{N,1,1}(r), R_{N/2,1,4}(r)$ of type

(4.138) must still be proved, see (4.152), (4.154), (4.155) below (cf. [D, p. 240 et seq.] where the analog of (4.138) is proved for $R_{N,1,2}(0) = K_N(0,0)$ in the case $V(x) = x^{2m}$ without lower order terms). By (2.3), $q_N \sim N^{1-1/(2m)}$ to leading order.

We use the notation in [DKMVZ2]. In particular set

$$g(z) \equiv g_N(z) = \int_{-1}^{1} \log(z-x)\, d\mu^{(\mathrm{eq})}(x) = \int_{-1}^{1} \log(z-x)\, \frac{1}{2\pi}|1-x^2|^{1/2}\, h_N(x)\, dx,$$

$z \in \mathbb{C} \setminus (-\infty,-1]$, and for $z \in (-1,1)$ let

$$\begin{aligned} S_+(z) &= \begin{pmatrix} c_N^{-N} & 0 \\ 0 & c_N^{N} \end{pmatrix} e^{-\frac{Nl}{2}\sigma_3}\, Y_+(c_N z + d_N) \\ &\quad \times e^{-N(g_+(z)-\frac{l}{2})\sigma_3} \begin{pmatrix} 1 & 0 \\ -e^{-N(g_+(z)-g_-(z))} & 1 \end{pmatrix}, \end{aligned} \tag{4.139}$$

here ± refer to the boundary values from above/below the real axis, respectively (cf. [DKMVZ2, (4.22)]). Here $Y$ solves the Fokas–Its–Kitaev Riemann–Hilbert problem fot the polynomials orthogonal with respect to the weight $e^{-V(x)}dx$ (see [DKMVZ2, Thm. 3.1], and the constant $l \equiv l_N$ is given by (5.35), loc. cit.). Finally, for $z \in (-1,1)$ let

$$\xi_N(z) \equiv g_+(z) - g_-(z) = i\int_{z}^{1} |1-x^2|^{1/2} h_N(x)\, dx \tag{4.140}$$

and set

$$r_N \equiv (r-d_N)/c_N.$$

Note that $\xi_N(z) \in i\mathbb{R}_+$ and $r_N = O(N^{-1/(2m)})$ by (2.3), (2.4). We need the following properties of $S_+$ which are proved in [DKMVZ2, Sec. 7]:

(i) For $-1 < z < 1$,

$$\det S_+(z) = 1. \tag{4.141}$$

(ii) Let $a_+(z) = \left(\frac{z-1}{z+1}\right)_+^{1/4}$ and set

$$S_+^{(\infty)}(z) \equiv \frac{1}{2}\begin{pmatrix} a_+(z)+a_+(z)^{-1} & i(a_+(z)^{-1}-a_+(z)) \\ i(a_+(z)-a_+(z)^{-1}) & a_+(z)+a_+(z)^{-1} \end{pmatrix}, \qquad -1<z<1.$$

Then as $N \to \infty$, $S_+(z)$ converges with all its derivatives to $S_+^{(\infty)}(z)$ for all $z$ in compact subsets of $(-1,1)$. In particular, for any $0 < \delta < 1$,

$$\sup_N \max_{-1+\delta \le z \le 1-\delta} \left|\frac{d^k}{dz^k} S_+(z)\right| \le c_k < \infty \tag{4.142}$$

for $k = 0, 1, 2, \cdots$.

Now a simple calculation (cf. [D, p. 24]) shows that

$$
\begin{aligned}
&D_{r,N}(\xi,\eta) \qquad (4.143)\\
&= \Bigg[\bigg(S_{+11}\Big(r_N+\frac{\eta}{q_Nc_N}\Big)\,e^{\frac{N}{2}\xi_N\big(r_N+\eta/(q_Nc_N)\big)}+S_{+12}\Big(r_N+\frac{\eta}{q_Nc_N}\Big)\,e^{-\frac{N}{2}\xi_N\big(r_N+\eta/(q_Nc_N)\big)}\bigg)\\
&\quad\times\bigg(S_{+21}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)\,e^{\frac{N}{2}\xi_N\big(r_N+\xi/(q_Nc_N)\big)}+S_{+22}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)\,e^{-\frac{N}{2}\xi_N\big(r_N+\xi/(q_Nc_N)\big)}\bigg)\\
&\qquad-(\xi\leftrightarrow\eta)\Bigg]\Big/\big(2\pi i(\xi-\eta)\big)
\end{aligned}
$$

where $(\xi\leftrightarrow\eta)$ indicates the same terms with $\xi$ and $\eta$ interchanged. Consider the terms

$$
\begin{aligned}
&(4.144)\\
&\bigg[S_{+11}\Big(r_N+\frac{\eta}{q_Nc_N}\Big)\,S_{+22}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)\,e^{\frac{N}{2}\big(\xi_N(r_N+\eta/(q_Nc_N))-\xi_N(r_N+\xi/(q_Nc_N))\big)}\\
&\quad-S_{+11}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)\,S_{+22}\Big(r_N+\frac{\eta}{q_Nc_N}\Big)\,e^{\frac{N}{2}\big(\xi_N(r_N+\xi/(q_Nc_N))-\xi_N(r_N+\eta/(q_Nc_N))\big)}\bigg]\\
&\quad\big/\big(2\pi i(\xi-\eta)\big)\equiv Q_{8,1}+Q_{8,2}+Q_{8,3}
\end{aligned}
$$

where

$$
\begin{aligned}
Q_{8,1}\equiv&\frac{S_{+11}\Big(r_N+\frac{\eta}{q_Nc_N}\Big)-S_{+11}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)}{2\pi i(\xi-\eta)}\\
&\times S_{+22}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)\,e^{\frac{N}{2}\big(\xi_N(r_N+\eta/(q_Nc_N))-\xi_N(r_N+\xi/(q_Nc_N))\big)},
\end{aligned}
$$

and

$$
\begin{aligned}
Q_{8,2}\equiv&S_{+11}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)\,e^{\frac{N}{2}\big(\xi_N(r_N+\eta/(q_Nc_N))-\xi_N(r_N+\xi/(q_Nc_N))\big)}\\
&\times\frac{S_{+22}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)-S_{+22}\Big(r_N+\frac{\eta}{q_Nc_N}\Big)}{2\pi i(\xi-\eta)},
\end{aligned}
$$

and also

$$
\begin{aligned}
Q_{8,3}\equiv&S_{+11}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)\,S_{+22}\Big(r_N+\frac{\eta}{q_Nc_N}\Big)\\
&\times\frac{e^{\frac{N}{2}\big(\xi_N(r_N+\eta/(q_Nc_N))-\xi_N(r_N+\xi/(q_Nc_N))\big)}-e^{\frac{N}{2}\big(\xi_N(r_N+\xi/(q_Nc_N))-\xi_N(r_N+\eta/(q_Nc_N))\big)}}{2\pi i(\xi-\eta)}.
\end{aligned}
$$

Note that

$$
\begin{aligned}
&(4.145)\\
&\frac{S_{+11}\Big(r_N+\frac{\eta}{q_Nc_N}\Big)-S_{+11}\Big(r_N+\frac{\xi}{q_Nc_N}\Big)}{2\pi i(\xi-\eta)}=-\frac{1}{2\pi i}\int_0^1 S'_{+11}\Big(r_N+\frac{\eta+\tau(\xi-\eta)}{q_Nc_N}\Big)\,\frac{d\tau}{q_Nc_N}
\end{aligned}
$$

and

$$
\begin{aligned}
\frac{N}{2}\bigg(\xi_N\big(r_N+\frac{\eta}{q_Nc_N}\big)-\xi_N\big(r_N+\frac{\xi}{q_Nc_N}\big)\bigg)&=i\pi(\xi-\eta)\,G_N(\xi,\eta)\\
&=i\pi(\xi-\eta)+O(N^{-\alpha})
\end{aligned}
\tag{4.146}
$$

as $N \to \infty$, where we have denoted

$$G_N(\xi,\eta) \equiv \frac{N}{2\pi q_N c_N} \int_0^1 \sqrt{1-\Big(\frac{\eta+t(\xi-\eta)}{q_N c_N}\Big)^2}\, h_N\Big(\frac{\eta+t(\xi-\eta)}{q_N c_N}\Big)\, dt. \tag{4.147}$$

and observed that by (4.138),

$$G_N(\xi,\eta) = \frac{N h_N(0)}{2\pi q_N c_N}\,(1+O(N^{-1})) = 1+O(N^{-\alpha}), \qquad N\to\infty. \tag{4.148}$$

From the formulae above and the relation $\frac{iN(\xi-\eta)}{2q_N c_N} = \frac{i\pi(\xi-\eta)}{h_N(0)+o(1)}$, we see that $Q_{8,1}$, and similarly $Q_{8,2}$, and all their $\xi,\eta$ derivatives converge to zero uniformly for $|\xi|,|\eta|,|r| \le 2L_0$, with errors of order $O(N^{-1})$.

We also have the formula

$$\begin{aligned} Q_{8,3} = S_{+11}\Big(r_N + \frac{\xi}{q_N c_N}\Big)\, S_{+22}\Big(r_N+\frac{\eta}{q_N c_N}\Big)& \\ \times G_N(\xi,\eta) \int_0^1 \cos\big(\tau\pi(\xi-\eta)\, G_N(\xi,\eta)\big)\, d\tau.& \end{aligned} \tag{4.149}$$

Together with similar formulae and calculations for the other terms in (4.143) we see that $D_{r,N}(\xi,\eta) = \frac{\sin\left((\xi-\eta)\frac{N h_N(0)}{2\pi q_N c_N}\right)}{\pi(\xi-\eta)} + O(N^{-1}) = K_\infty(\xi,\eta) + O(N^{-\alpha}) + O(N^{-1})$ where $K_\infty(\xi-\eta)$ is the sine kernel $\frac{\sin\pi(\xi-\eta)}{\pi(\xi-\eta)}$. The same is true for all the $\xi,\eta$ derivatives. More precisely we have for any $j,k\ge 0$,

$$\frac{\partial^j}{\partial\xi^j}\,\frac{\partial^k}{\partial\eta^k}\,\frac{1}{q_N} K_N\left(r+\frac{\xi}{q_N}, r+\frac{\eta}{q_N}\right) \to \frac{\partial^j}{\partial\xi^j}\,\frac{\partial^k}{\partial\eta^k}\, K_\infty(\xi-\eta) \tag{4.150}$$

uniformly for $|\xi|,|\eta|,|r| \le 2L_0$ as $N\to\infty$. Note that the terms of the form $S_{+11}(\cdots)S_{+21}(\cdots)$ and $S_{+12}(\cdots)S_{+22}(\cdots)$ do not contribute to leading order, and also $S_{+11}(r_N)S_{+22}(r_N) - S_{+12}(r_N)S_{+21}(r_N) = 1$ by (4.141). Moreover, keeping track of the estimates, we find that

(4.151) the error term in (4.150) is $O(N^{-\alpha})$ uniformly for $|\xi|,|\eta|,|r| \le 2L_0$.

We see from the above that the largest error in (4.152) arises purely from the asymptotic evaluation of $\frac{N h_N(0)}{2\pi q_N c_N}$ using (4.138) (cf. (4.148)): apart from this term the error in (4.152) is $O(N^{-1})$, rather than $O(N^{-\alpha})$.

4.3.2. We can now verify asymptotics for $K_N(r,r)$ of type (4.138). Indeed from the uniform convergence above, taking $q_N \equiv \frac{N h_N(0)}{2\pi c_N}$, for which we may, in turn, take $\alpha = 1$ in (4.138), we have for $|r|\le 2L_0$, as $N\to\infty$,

$$\begin{aligned} K_N(r,r) &= q_N\, D_{r,N}(0,0) = q_N \lim_{\xi\to 0} D_{r,N}(\xi,0) = q_N\left(\lim_{\xi\to 0}\frac{\sin\pi(\xi-0)}{\pi(\xi-0)} + O(N^{-1})\right) \\ &= q_N\big(1+O(N^{-1})\big) = \frac{N h_N(0)}{2\pi c_N}\big(1+O(N^{-1})\big). \end{aligned} \tag{4.152}$$

This means in particular that (4.150), (4.151) are true for $q_N = K_N(r,r)$ with $\alpha = 1$ for any $|r| \le 2L_0$. Hence

$$K_N(r,r) \sim N^{1-1/(2m)}, \qquad \text{for } |r|\le 2L_0. \tag{4.153}$$

Estimating the correction terms for $S_{N,1}(r,r)$ as in (3.5), we find for any $|r| \le 2L_0$

$$\frac{S_{N,1}(r,r)}{K_N(r,r)} = 1 + O(N^{-1/2}). \tag{4.154}$$

Similarly for any $|r| \le 2L_0$

$$\frac{S_{N/2,4}(r,r)}{K_N(r,r)} = 1 + O(N^{-1/2}). \tag{4.155}$$

Note that (4.154), (4.155) together with (4.153) prove (1.23), and hence (1.24).

4.3.3. *Derivatives, continued.* We now consider the term

$$X_N(\xi,\eta) \equiv \frac{1}{q_N^2}(K_N D)\left(r + \frac{\xi}{q_N}, r + \frac{\eta}{q_N}\right), \qquad q_N \text{ as in (4.138)}. \tag{4.156}$$

Terms of this form arise in the 12 entry of $\frac{1}{q_N} K_{N,1}^{(\lambda_N,1)}\big(r + \frac{\xi}{q_N}, r + \frac{\eta}{q_N}\big)$. Clearly $X_N(\xi,\eta) = -\frac{\partial}{\partial\eta} D_{r,N}(\xi,\eta)$ (cf. (4.137)) and so by (4.150) we see that (4.156) satisfies

$$X_N(\xi,\eta) \to -\frac{\partial}{\partial\eta} K_\infty(\xi - \eta) = \frac{\partial}{\partial\xi} K_\infty(\xi-\eta) \tag{4.157}$$

together with all its $\xi, \eta$ derivatives, uniformly for $|\xi|, |\eta|, |r| \le 2L_0$. Moreover as before

(4.158) the error term in (4.157) is $O(N^{-\alpha})$ uniformly for $|\xi|, |\eta|, |r| \le 2L_0$.

4.3.4. *Integrals.* Finally, consider the term

$$Y_N(\xi,\eta) \equiv \frac{1}{2}\int_{-\infty}^{\infty} \operatorname{sgn}\left(r + \frac{\xi}{q_N} - s\right) K_N\left(s,\, r + \frac{\eta}{q_N}\right) ds, \qquad q_N \text{ as in (4.138)}. \tag{4.159}$$

Terms of this form arise in the 21 entry of $\frac{1}{q_N} K_{N,1}^{(\lambda_N,1)}\big(r + \frac{\xi}{q_N}, r + \frac{\eta}{q_N}\big)$. We want to show that $Y_N(\xi,\eta) \to \int_0^{\xi-\eta} K_\infty(t)\,dt$ as $N \to \infty$. Changing variables $s = r + (\eta + t)/q_N$ we write $Y_N$ as

$$Y_N(\xi,\eta) = Q_{9,1} + Q_{9,2} + Q_{9,3} \tag{4.160}$$

where (cf. (4.137))

$$\begin{aligned} Q_{9,1} &\equiv \int_0^{\xi-\eta} D_{r,N}(t+\eta,\eta)\,dt \\ Q_{9,2} &\equiv \frac{1}{2}\int_{-\infty}^{0} D_{r,N}(t+\eta,\eta)\,dt \\ Q_{9,3} &\equiv -\frac{1}{2}\int_{0}^{\infty} D_{r,N}(t+\eta,\eta)\,dt. \end{aligned}$$

Again we consider $\xi, \eta, r$ in the compact set $[-2L_0, 2L_0]$. By (4.150), $Q_{9,1}$ converges, with all its derivatives, to $\int_0^{\xi-\eta} K_\infty(t+\eta-\eta)\,dt = \int_0^{\xi-\eta} K_\infty(t)\,dt$ as $N \to \infty$, and so we must show that $Q_{9,2} + Q_{9,3}$ converges to 0.

To analyze $Q_{9,2}, Q_{9,3}$ it is convenient to consider the cases $|t| \le 1$ and $|t| > 1$ separately. By the above calculations, we have as $N \to \infty$

$$\begin{aligned} &\frac{1}{2}\Big(\int_{-1}^{0} - \int_{0}^{1}\Big) D_{r,N}(t+\eta,\eta)\,dt \\ &\qquad = \frac{1}{2}\Big(\int_{-1}^{0} - \int_{0}^{1}\Big) K_\infty(t+\eta-\eta)\,dt + O(N^{-\alpha}) = O(N^{-\alpha}), \end{aligned} \tag{4.161}$$

as $K_\infty(t)$ is even. As before the term $O(N^{-\alpha})$ is uniform for $|r|, |\eta| \le 2L_0$. Now we must show $Q_{10,1} - Q_{10,2} \to 0$ as $N \to \infty$, where

$$Q_{10,1} \equiv \int_{-\infty}^{-1} D_{r,N}(t+\eta,\eta)\,dt \qquad \text{and} \qquad Q_{10,2} \equiv \int_{1}^{\infty} D_{r,N}(t+\eta,\eta)\,dt.$$

Consider $Q_{10,2}$. We utilize the change of variables

$$r + (t+\eta)/q_N = c_N z + d_N \tag{4.162}$$

i.e.

$$\begin{aligned} t &= -\eta + (d_N - r)q_N + q_N c_N z \\ z &= (r - d_N)/c_N + (t+\eta)/(q_N c_N). \end{aligned} \tag{4.163}$$

Fix $\delta > 0$ small as in Subsection 4.2, set $t_\pm \equiv -\eta + (d_N - r)q_N + q_N c_N(1 \pm \delta)$, and write $Q_{10,2} \equiv Q_{11,1} + Q_{11,2} + Q_{11,3}$ where

$$Q_{11,1} \equiv \int_{1}^{t_-}, \qquad Q_{11,2} \equiv \int_{t_-}^{t_+}, \qquad Q_{11,3} \equiv \int_{t_+}^{\infty}.$$

Note that $t_\pm \sim N$ as $N \to \infty$.

We have by (4.137), (1.7)

$$Q_{11,3} = \int_{t_+}^{\infty} b_{N-1}\Big[\phi_N\Big(r+\frac{t+\eta}{q_N}\Big)\phi_{N-1}\Big(r+\frac{\eta}{q_N}\Big) - \phi_{N-1}\Big(r+\frac{t+\eta}{q_N}\Big)\phi_N\Big(r+\frac{\eta}{q_N}\Big)\Big]\frac{dt}{t}.$$

Changing variables $t \to z$ as in (4.163), the first term takes the form

$$\int_{1+\delta}^{\infty} b_{N-1}\phi_N(c_N z + d_N)\,\phi_{N-1}(c_N z_0 + d_N)\,\frac{q_N c_N\,dz}{q_N c_N z + q_N(d_N - r) - \eta}$$

where $z_0 = z(t=0) = (r-d_N)/c_N + \eta/(q_N c_N) = O(N^{-1/(2m)})$. As $b_{N-1} \sim c_N/2$, using (3.1), (4.16) we see that the above term is bounded by

$$\text{const} \cdot ((c_N/N)^{1/2} e^{-cN}) \cdot (1/c_N^{1/2}) \le \text{const} \cdot e^{-cN}$$

for some $c = c(\delta) > 0$. There is a similar estimate for the second term in $Q_{11,3}$, and we have

$$|Q_{11,3}| = O(e^{-cN}), \qquad c = c(\delta) > 0, \tag{4.164}$$

uniformly for $|r|, |\eta| \le 2L_0$.

Changing variables $t \to z$ as in (4.163) in the first term of $Q_{11,2}$, and then integrating by parts, we obtain

$$\begin{aligned} b_{N-1}\phi_{N-1}(c_N z_0 + d_N)\Big[&\Big(\int_{1-\delta}^{1+\delta}\phi_N(c_N w + d_N)\,dw\Big)\frac{1}{1+\delta+\frac{d_N-r}{c_N}-\frac{\eta}{q_N c_N}} \\ &+ \int_{1-\delta}^{1+\delta}\Big(\int_{1-\delta}^{z}\phi_N(c_N w + d_N)\,dw\Big)\frac{dz}{(z+\frac{d_N-r}{c_N}-\frac{\eta}{q_N c_N})^2}\Big] \end{aligned}$$

which is bounded by

$$\text{const} \cdot c_N \cdot c_N^{-1/2} \cdot (c_N^{-1/2} N^{-1/2}) = O(N^{-1/2})$$

in view of (3.1), (4.17) (of course $z \sim 1$ in the above integrals, so there is no singularity in the integrand). Again there is a similar estimate for the second term in $Q_{11,2}$, and so

$$|Q_{11,2}| = O(N^{-1/2}) \tag{4.165}$$

uniformly for $|r|, |\eta| \leq 2L_0$.

Let $0 < \alpha' < \min\left(\frac{1}{2}, \alpha\right)$ and set $t_N \equiv N^{\alpha'}$. Thus $1 < t_N < t_-$ as $N \to \infty$. Let $Q_{11,1} = Q'_{11,1} + Q''_{11,1}$ where

$$Q'_{11,1} \equiv \int_1^{t_N} D_{r,N}(t+\eta,\eta)\,dt \qquad \text{and} \qquad Q''_{11,1} \equiv \int_{t_N}^{t_-} D_{r,N}(t+\eta,\eta)\,dt.$$

First we consider $Q'_{11,1}$. From (4.143) we again see that $D_{r,N}(t+\eta,\eta)$ is a sum of terms of the form $S_{+11}(\cdots)S_{+22}(\cdots)$, $S_{+12}(\cdots)S_{+21}(\cdots)$, $\cdots$. Express the term $S_{+11}S_{+22}$ as a sum

$$Q_{8,1}(t+\eta,\eta) + Q_{8,2}(t+\eta,\eta) + Q_{8,3}(t+\eta,\eta)$$

as in (4.144). As in the proof of (4.150), we have $|Q_{8,1}(t+\eta,\eta)|, |Q_{8,2}(t+\eta,\eta)| = O(N^{-1})$ as $N \to \infty$ uniformly for $1 < t < t_N$ and $|r|, |\eta| \leq 2L_0$. Thus the contribution of $Q_{8,1}(t+\eta,\eta), Q_{8,2}(t+\eta,\eta)$ to $Q'_{11,1}$ is $O(t_N N^{-1}) = O(N^{-1+\alpha'})$ uniformly for $|r|, |\eta| \leq 2L_0$. From (4.149)

$$Q_{8,3}(t+\eta,\eta) = S_{+11}\Big(r_N + \frac{t+\eta}{q_N c_N}\Big)\, S_{+22}\Big(r_N + \frac{\eta}{q_N c_N}\Big)\, \frac{\sin(\pi t\, G_N(t+\eta,\eta))}{\pi t}$$

where $G_N$ is as in (4.147). By (4.142)

$$\begin{aligned} Q_{8,3}(t+\eta,\eta) &= S_{+11}(r_N)\, S_{+22}(r_N)\, \frac{\sin(\pi t\, G_N(t+\eta,\eta))}{\pi t} + O\bigg(\frac{1}{N} \max_{t\geq 1} \Big|\frac{t+|\eta|}{t}\Big|\bigg) \\ &= S_{+11}(r_N)\, S_{+22}(r_N)\, \frac{\sin \pi t}{\pi t}\, \cos\big[\pi t(G_N(t+\eta,\eta)-1)\big] \\ &\quad + S_{+11}(r_N)\, S_{+22}(r_N)\, \frac{\cos \pi t}{\pi t}\, \sin\big[\pi t(G_N(t+\eta,\eta)-1)\big] + O(N^{-1}). \end{aligned}$$

But by (4.138), (4.152), we find from (4.147)

$$G_N(t+\eta,\eta) = \frac{N h_N(0)}{2\pi q_N c_N}\,(1 + O(t/N)) = 1 + O(t/N) + O(N^{-\alpha})$$

uniformly for $1 \leq t \leq t_N$, $|\eta| \leq 2L_0$. Thus

$$\left|\frac{\sin\big[\pi t(G_N(t+\eta,\eta)-1)\big]}{\pi t}\right| \leq |G_N(t+\eta,\eta) - 1| = O(t/N) + O(N^{-\alpha})$$

and similarly

$$\big|\cos\big[\pi t(G_N(t+\eta,\eta)-1)\big] - 1\big| = O(t^4/N^2) + O(t^2/N^{2\alpha}).$$

Thus as $\alpha \leq 1$,

$$Q_{8,3}(t+\eta,\eta) = S_{+11}(r_N)\, S_{+22}(r_N)\, \frac{\sin \pi t}{\pi t} + O\bigg(\frac{t^3}{N^2} + \frac{t}{N} + \frac{t}{N^{2\alpha}} + \frac{1}{N^{\alpha}}\bigg)$$

and so

$$\begin{aligned}\int_1^{t_N} Q_{8,3}(t+\eta,\eta)\,dt =& S_{+11}(r_N)\,S_{+22}(r_N)\int_1^{t_N}\frac{\sin\pi t}{\pi t}\,dt\\ &+O\left(\frac{N^{4\alpha'}}{N^2}+\frac{N^{2\alpha'}}{N}+\frac{N^{2\alpha'}}{N^{2\alpha}}+\frac{N^{\alpha'}}{N^{\alpha}}\right).\end{aligned}$$

Assembling the above estimates we find for $0<\alpha'<\min\left(\frac{1}{2},\alpha\right)$ as $N\to\infty$, the contribution of the $S_{+11}S_{+22}$ term to $Q'_{11,1}$ is given by

$$S_{+11}(r_N)\,S_{+22}(r_N)\int_1^{t_N}\frac{\sin\pi t}{\pi t}\,dt+O(N^{-1+2\alpha'})+O(N^{-\alpha+\alpha'})$$

uniformly for $|r|,|\eta|\le 2L_0$. Analyzing the other contributions $S_{+12}S_{+21},\ \cdots$ we obtain as $N\to\infty$

$$Q'_{11,1}=\int_1^{t_N}D_{r,N}(t+\eta,\eta)\,dt=\int_1^{t_N}\frac{\sin\pi t}{\pi t}\,dt+O(N^{-1+2\alpha'})+O(N^{-\alpha+\alpha'}) \tag{4.166}$$

uniformly for $|r|,|\eta|\le 2L_0$. Again we have used (4.141).

Finally we consider

$$Q''_{11,1}=\int_{t_N}^{t_-}b_{N-1}\left[\phi_N\left(r+\frac{t+\eta}{q_N}\right)\phi_{N-1}\left(r+\frac{\eta}{q_N}\right)-\phi_{N-1}\left(r+\frac{t+\eta}{q_N}\right)\phi_N\left(r+\frac{\eta}{q_N}\right)\right]\frac{dt}{t}.$$

Changing variables $t\to z$ as in (4.163) and then integrating by parts, we obtain for the first term in $Q''_{11,1}$

$$\begin{aligned}b_{N-1}\phi_{N-1}(r+\eta/q_N)\Bigg[&\left(\int_{z(t_N)}^{1-\delta}\phi_N(c_Nw+d_N)\,dw\right)\frac{q_Nc_N}{q_Nc_N(1-\delta)+(d_N-r)q_N-\eta}\\ &+\int_{z(t_N)}^{1-\delta}\left(\int_{z(t_N)}^{z}\phi_N(c_Nw+d_N)\,dw\right)\frac{(q_Nc_N)^2\,dz}{(q_Nc_Nz+(d_N-r)q_N-\eta)^2}\Bigg].\end{aligned}$$

Note that $z(t_N)\sim N^{-1+\alpha}$ and recall (3.1), (4.18) to conclude that the first term in $Q''_{11,1}$ is bounded by

$$\text{const}\cdot b_{N-1}\cdot c_N^{-1/2}\cdot(c_N^{-1/2}N^{-1})\cdot\left(\text{const}+q_Nc_N\int_{t_N}^{t_-}\frac{dt}{t^2}\right)=O(N^{-\alpha'}).$$

The same is true for the second term in $Q''_{11,1}$, and so together with (4.166) we find

$$Q_{11,1}=\int_1^{t_-}D_{r,N}(t+\eta,\eta)\,dt=\int_1^{t_N}\frac{\sin\pi t}{\pi t}\,dt+O(N^{-1+2\alpha'}+N^{-\alpha'}+N^{-\alpha+\alpha'}) \tag{4.167}$$

and hence, as $\alpha'<1/2$

$$\begin{aligned}Q_{10,2}&=\int_1^{\infty}D_{r,N}(t+\eta,\eta)\,dt\\ &=\int_1^{t_N}\frac{\sin\pi t}{\pi t}\,dt+O(N^{-1+2\alpha'}+N^{-\alpha'}+N^{-1/2}+N^{-\alpha+\alpha'}+e^{-cN})\\ &=\int_1^{t_N}\frac{\sin\pi t}{\pi t}\,dt+O(N^{-1+2\alpha'}+N^{-\alpha'}+N^{-\alpha+\alpha'}).\end{aligned}$$

The best error estimate is clearly obtained for $\alpha' = \alpha^* \equiv \min\left(\frac{\alpha}{2}, \frac{1}{3}\right) < \alpha$. Thus

$$\int_1^\infty D_{r,N}(t+\eta,\eta)\,dt = \int_1^{t_N} \frac{\sin \pi t}{\pi t}\,dt + O(N^{-\alpha^*}) \tag{4.168}$$

and similarly

$$\int_{-\infty}^{-1} D_{r,N}(t+\eta,\eta)\,dt = \int_{-t_N}^{-1} \frac{\sin \pi t}{\pi t}\,dt + O(N^{-\alpha^*}). \tag{4.169}$$

But $\frac{\sin \pi t}{\pi t}$ is even, and together with (4.161) we see that

$$Q_{9,2} + Q_{9,3} = \frac{1}{2}\left(\int_{-\infty}^{0} D_{r,N}(t+\eta,\eta)\,dt - \int_0^\infty D_{r,N}(t+\eta,\eta)\,dt\right) = O(N^{-\alpha^*}) \tag{4.170}$$

as $N \to \infty$ uniformly for $|r|, |\eta| \le 2L_0$. We have proved the following (cf. (4.159)):

$$\begin{aligned} &Y_N(\xi,\eta) = A_N(\xi,\eta) + B_N(\eta), \qquad \text{where as } N \to \infty \\ &\text{(a)}\quad A_N(\xi,\eta) \text{ converges with all its } \xi,\eta \text{ derivatives to } \int_0^{\xi-\eta} K_\infty(t)\,dt \\ &\qquad\qquad \text{with error } O(N^{-\alpha}) \text{ uniformly for } \xi,\eta,r \in [-2L_0, 2L_0]; \\ &\text{(b)}\quad B_N(\eta) = O(N^{-\alpha^*}) \text{ uniformly for } \eta, r \in [-2L_0, 2L_0]. \end{aligned} \tag{4.171}$$

Here $\alpha, \alpha^*$ are as in (4.138), (4.168), respectively.

## 5. Proofs of Theorems 2.5, 2.3 and 2.4

5.1. **Proof of Theorem 2.5.** We know by Lemma 4.1 that $D\phi_k$ is a (finite) linear combination of $\phi_{\max(0,k-n)}, \cdots, \phi_{k+n}$, $n = \deg V'$. Hence

$$D\phi_k = \sum_{j=\max(0,k-n)}^{k+n} (D\phi_k, \phi_j)\,\phi_j. \tag{5.1}$$

Note next that since $\phi_k$ and its derivative are rapidly decaying,

$$\epsilon D\phi_k = \phi_k, \qquad k \in \mathbb{Z}_{\ge 0}. \tag{5.2}$$

Apply $\epsilon$ to (5.1) and take into account (5.2) to find

$$\phi_k = \epsilon D\phi_k = \sum_{j=\max(0,k-n)}^{k+n} (D\phi_k, \phi_j)\,\epsilon\phi_j. \tag{5.3}$$

Finally, take the inner product of (5.3) with any $\phi_l$ to find

$$\delta_{k,l} = \sum_{j=\max(0,k-n)}^{k+n} (D\phi_k, \phi_j)\,(\epsilon\phi_j, \phi_l) \tag{5.4}$$

which shows that $I_\infty = D_\infty \epsilon_\infty$ (recall that $(D\phi_k, \phi_j) = 0$ for $|j-k| > n$). Taking the transposes and using the fact that $D_\infty$ and $\epsilon_\infty$ are both skew symmetric gives the second statement in Theorem 2.5(i).

Now for the proof of Theorem 2.5(ii). Denote the $N \times (N+n)$ section of $\epsilon_\infty$ by $\epsilon_{N,N+n}$ and denote the last $n$ columns of $\epsilon_{N,N+n}$ by

$$\begin{pmatrix} \epsilon^*_{N-n,n} \\ B_{12} \end{pmatrix} \tag{5.5}$$

($\epsilon^*_{N-n,n}$ is the same matrix as in (2.18)). Denote the $(N+n)\times N$ section of $D_\infty$ by $D_{N+n,N}$. Note that the last $n$ rows of this matrix are $\begin{pmatrix}0_{n,N-n} & D_{21}\end{pmatrix}$. By Theorem 2.5(i)

$$\begin{pmatrix} I_{N-n} & 0 \\ \\ 0 & I_n \end{pmatrix} = \epsilon_{N,N+n} D_{N+n,N} = \epsilon_N D_N + \left( 0_{N,N-n} \middle| \begin{matrix} \epsilon^*_{N-n,n} D_{21} \\ B_{12} D_{21} \end{matrix} \right)$$

which proves Theorem 2.5(ii).

5.2. **Proof of Theorem 2.3.** Recall the definitions of the block matrices $A$, $B$, and $C$. Note first that

$$BA = \begin{pmatrix} B_{12}A_{21} & B_{11}A_{12} \\ \\ B_{22}A_{21} & B_{21}A_{12} \end{pmatrix} = \begin{pmatrix} -B_{12}D_{21} & B_{11}D_{12} \\ \\ -B_{22}D_{21} & B_{21}D_{12} \end{pmatrix}.$$

Recall from Section 2 that $R$ denotes the $n \times n$ matrix with all zeros and ones on the anti-diagonal. Let $G^T$ denote the transpose of $G$ and $G^\perp$ be the transpose with respect to the anti-diagonal. An application of $R$ to a matrix from both sides interchanges the order of all rows and columns in the matrix. We have for any $G$

$$RGR = (G^\perp)^T = (G^T)^\perp \quad \text{and also} \quad RR = I_n. \tag{5.6}$$

It is convenient to denote for $j,k = 1,2$

$$\tilde{B}_{jk} \equiv N^{1-1/(2m)}\, B_{jk}, \qquad \tilde{D}_{jk} \equiv N^{-1+1/(2m)}\, D_{jk}, \qquad \tilde{A}_{jk} \equiv N^{-1+1/(2m)}\, A_{jk}.$$

Then by Theorem 2.1 and 2.2 the matrices with the tilde tend to certain explicit limiting matrices, as $N \to \infty$, see e.g. (2.13), (2.3). Note that (2.13) implies that as $N \to \infty$, $N$ even,

$$\tilde{B}_{12}^\perp = \tilde{B}_{12} + o(1), \qquad \tilde{D}_{21}^\perp = \tilde{D}_{21} + o(1) \tag{5.7}$$

and also from the skew symmetry of $\epsilon_\infty$ and $D_\infty$

$$B_{12}^T = -B_{21}, \qquad D_{21}^T = -D_{12}.$$

We prove now the first statement of Theorem 2.3:

$$\begin{aligned} -R(BA)_{11}R &= RB_{12}D_{21}R = RB_{12}RRD_{21}R = (B_{12}^\perp)^T (D_{21}^\perp)^T = (\tilde{B}_{12}^\perp)^T (\tilde{D}_{21}^\perp)^T \\ &= (\tilde{B}_{12}^T + o(1))\,(\tilde{D}_{21}^T + o(1)) = (\tilde{B}_{21} + o(1))\,(\tilde{D}_{12} + o(1)) \\ &= \tilde{B}_{21}\tilde{D}_{12} + o(1) = B_{21}D_{12} + o(1) = (BA)_{22} + o(1). \end{aligned} \tag{5.8}$$

Next we prove the second statement of Theorem 2.3. First note that

$$BAC = \begin{pmatrix} (BAC)_{11} & (BAC)_{12} \\ \\ (BAC)_{21} & (BAC)_{22} \end{pmatrix}$$

where

$$\begin{aligned} (BAC)_{11} &= B_{12}A_{21} + B_{12}A_{21}B_{12}A_{21} + B_{11}A_{12}B_{22}A_{21} \\ (BAC)_{12} &= B_{12}A_{21}B_{11}A_{12} + B_{11}A_{12}B_{21}A_{12} \\ (BAC)_{21} &= B_{22}A_{21} + B_{22}A_{21}B_{12}A_{21} + B_{21}A_{12}B_{22}A_{21} \\ (BAC)_{22} &= B_{22}A_{21}B_{11}A_{12} + B_{21}A_{12}B_{21}A_{12}. \end{aligned} \tag{5.9}$$

Let us prove first that $(BAC)_{12} = 0$. Since $\det A_{12} \neq 0$ (by Theorem 2.1 and since $\det \tilde{A}_{12} \neq 0$) this is the same as proving

$$B_{12}A_{21}B_{11} + B_{11}A_{12}B_{21} = 0. \tag{5.10}$$

We claim that to prove (5.10) is in turn the same as to prove that $B_{12}A_{21}B_{11}$ is skew symmetric. This is because

$$B_{11}A_{12}B_{21} = B_{11}^T A_{21}^T B_{12}^T = (B_{12}A_{21}B_{11})^T$$

(recall $A_{12} = A_{21}^T$). Now since $B_{11}$ is skew, we note that $B_{12}A_{21}B_{11}$ is skew if and only if

$$(I_n + B_{12}A_{21})B_{11} = (I_n - B_{12}D_{21})B_{11}$$

is skew. But the last matrix is skew because it is the lower right $n \times n$ corner of the skew matrix $\epsilon_N D_N \epsilon_N$. (Here we used (2.18) and the fact that $B_{11}$ is the lower right $n \times n$ corner of $\epsilon_N$.)

Next we show that $(BAC)_{11} = 0$. Since $\det A_{21} \neq 0$ this is the same as proving that

$$(I_n + B_{12}A_{21})B_{12} + B_{11}A_{12}B_{22} = 0. \tag{5.11}$$

Recall that $A_{12} = D_{12}$, $A_{21} = -D_{21}$ and note that we have to prove

$$(I_n - B_{12}D_{21})B_{12} = -B_{11}D_{12}B_{22} \tag{5.12}$$

The matrix $(I - B_{12}D_{21})$ as we know is the lower right $n \times n$ corner of $\epsilon_N D_N$. In view of (2.18) and (5.5), $(I - B_{12}D_{21})B_{12}$ equals the product of the last $n$ rows of $(\epsilon_N D_N)$ with $\begin{pmatrix} \epsilon^*_{N-n,n} \\ B_{12} \end{pmatrix}$. But this is the same as the product of the last $n$ rows of $\epsilon_N$ with $D_N \begin{pmatrix} \epsilon^*_{N-n,n} \\ B_{12} \end{pmatrix}$. We now derive a different expression for the latter product. To this end we consider the product of the rows $0, \cdots, N-1$ in $D_\infty$ with the columns $N, \cdots, N+n-1$ in $\epsilon_\infty$. Since $D_\infty \epsilon_\infty = I_\infty$ this product equals $0_{N,n}$. But using the fact that $D_\infty$ is banded, and the expression (5.5) for $\epsilon_{N,N+n}$, this relation becomes

$$D_N \begin{pmatrix} \epsilon^*_{N-n,n} \\ B_{12} \end{pmatrix} = \begin{pmatrix} 0_{N-n,n} \\ -D_{12}B_{22} \end{pmatrix}.$$

Multiplying the latter expression from the left by the last $n$ rows of $\epsilon_N$ we see that the product is $-B_{11}D_{12}B_{22}$, and so (5.12) is established.

Our next goal is to show $(BAC)_{22} = (BA)_{22} + o(1)$. In view of (5.8) we must prove that $(BAC)_{22}$ equals

$$\begin{aligned} -R(BA)_{11}R + o(1) &= -RB_{12}A_{21}R + o(1) = -(RB_{12}R)(RA_{21}R) + o(1) \\ &= -(RB_{12}R)(A_{21}^{\perp})^T + o(1) = -(RB_{12}R)A_{21}^T + o(1) \\ &= -(RB_{12}R)A_{12} + o(1). \end{aligned}$$

Now using (5.9) and inserting tildes, we see that (recall $\det \tilde{A}_{12} \neq 0$) we must prove

$$\tilde{B}_{22}\tilde{A}_{21}\tilde{B}_{11} + \tilde{B}_{21}\tilde{A}_{12}\tilde{B}_{21} + R\tilde{B}_{12}R = o(1).$$

After noting that $R\tilde{B}_{12}R = -\tilde{B}_{21} + o(1)$, and then taking transposes, we see that we must prove

$$\tilde{B}_{12} + \tilde{B}_{11}\tilde{A}_{12}\tilde{B}_{22} + \tilde{B}_{12}\tilde{A}_{21}\tilde{B}_{12} = o(1).$$

But this follows from (5.11) (which holds exactly) and which we have already proved in the $(BAC)_{11}$ part.

Finally we show $(BAC)_{21} = (BA)_{21} + o(1)$ (which is not really needed in this paper; our proof of the fact that $\lim_{N\to\infty} \det(I_{2n} + BAC) \neq 0$ in Section 2 only uses $(BAC)_{22} = (BA)_{22} + o(1)$). By (5.9), since $(BA)_{21} = B_{22}A_{21}$ and again $\det \tilde{A}_{21} \neq 0$, we must show

$$\tilde{B}_{22}\tilde{A}_{21}\tilde{B}_{12} + \tilde{B}_{21}\tilde{A}_{12}\tilde{B}_{22} = o(1).$$

Since $\tilde{B}_{22} = \tilde{B}_{11}^{\perp} + o(1)$, this reduces to showing

$$\tilde{B}_{11}^{\perp}\tilde{A}_{21}\tilde{B}_{12} + \tilde{B}_{21}\tilde{A}_{12}\tilde{B}_{11}^{\perp} = o(1).$$

Now we apply the transposition across the anti-diagonal and use the property $(GH)^{\perp} = H^{\perp}G^{\perp}$ together with (5.7) to find

$$\tilde{B}_{11}\tilde{A}_{12}\tilde{B}_{21} + \tilde{B}_{12}\tilde{A}_{21}\tilde{B}_{11} = o(1).$$

But this follows from (5.10) (the latter holds exactly) which we proved in the $(BAC)_{12}$ part. The proof of Theorem 2.3 is now complete.

5.3. **Proof of Theorem 2.4.** Assume $N$ is even. Let us consider first the case when $V(x)$ is even. Then all the even-numbered OP's and also the functions $\phi_j$ are even, and all the odd-numbered ones are odd functions on $\mathbb{R}$. As the maps $f \to \epsilon f$, $f \to f'$ reverse parity, it follows that the $(i, j)$ entries of $\epsilon_N$ and $D_N$ are zero if $i$ and $j$ have the same parity. In turn we see that the $(i, j)$ entries of $W_N = \epsilon_N D_N$ are zero if $i$ and $j$ have the opposite parity. Let $W_N^{(1)}$, $W_N^{(2)}$ be the $N/2 \times N/2$ matrices constructed from the rows and columns $1, 3, \cdots, N-1$ (respectively, $0, 2, \cdots, N-2$) of $W_N$. Clearly $\det W_N = \det W_N^{(1)} \det W_N^{(2)}$. But more is true: by (2.18) and the asymptotics of $T_m'$, $T_{m-1}$ in (2.16), (2.17) we must have that $\det W_N^{(1)} = \det T_m' + o(1)$ and $\det W_N^{(2)} = \det T_{m-1} + o(1)$. Now $W_N^{(1)} = \epsilon_N^{(1)} D_N^{(1)}$ and $W_N^{(2)} = \epsilon_N^{(2)} D_N^{(2)}$ where $\epsilon_N^{(i)}$, $D_N^{(i)}$, $i = 1, 2$, are the following $N/2 \times N/2$ matrices:

- $\epsilon_N^{(1)}$, $\epsilon_N^{(2)}$ are formed from rows $1, 3, \cdots, N-1$ (respectively, $0, 2, \cdots, N-2$) and columns $0, 2, \cdots, N-2$ (respectively, $1, 3, \cdots, N-1$) of $\epsilon_N$
- $D_N^{(1)}$, $D_N^{(2)}$ are formed from rows $0, 2, \cdots, N-2$ (respectively, $1, 3, \cdots, N-1$) and columns $1, 3, \cdots, N-1$ (respectively, $0, 2, \cdots, N-2$) of $D_N$.

Thus $\det T_m' = \det \epsilon_N^{(1)} \det D_N^{(1)} + o(1)$ and $\det T_{m-1} = \det \epsilon_N^{(2)} \det D_N^{(2)} + o(1)$. But as $\epsilon_N$, $D_N$ are skew symmetric, a simple calculation shows that

$$\epsilon_N^{(1)} = -\big(\epsilon_N^{(2)}\big)^T, \qquad D_N^{(1)} = -\big(D_N^{(2)}\big)^T.$$

Letting $N \to \infty$, it follows that $\det T_m' = \det T_{m-1}$.

Now for the general case when $V(x)$ is not assumed to be even. Note that the $N$-independent matrices $T_m'$ and $T_{m-1}$ depend only on the degree $2m$ of $V(x)$ (and not on $\kappa_{2m} > 0$). But this means that $\det T_m'$ and $\det T_{m-1}$ are equal to the corresponding determinants for the *even* weight $\kappa_{2m}x^{2m}$. By the above argument, $\det T_m' = \det T_{m-1}$. This completes the proof of Theorem 2.4.

## 6. DIFFERENTIAL EQUATIONS FOR $h(x)$, $y_m(\theta)$, AND PROOF OF THEOREM 2.6

6.1. **Plan of the proof.** The goal in this Section is to prove that for all $m \geq 2$, $\det T_{m-1} \neq 0$, where $T_{m-1}$ is defined in (2.17). The key object to understand is $I(q)$ in (2.11) for $q = 3, 5, \cdots, 2n-3 = 4m-5$. After introducing some convenient notation we proceed to the proof. As noted in Remark 2.3, we use different arguments for $m$ "small" and "large," and for a given $m$, we must estimate $I(q)$ for different ranges of $q$. More precisely, in view of Proposition 6.1 below, what we really need is a bound on $|1+\tilde{I}(q)|$, where $\tilde{I}(q) \equiv mI(q) - \frac{1}{2}$ (see (2.20) and (6.3) below). For some values of $q$ and $m$, we estimate $|\tilde{I}(q)|$, and hence $|1+\tilde{I}(q)| \leq 1+|\tilde{I}(q)|$, but in other ranges we must consider the absolute value of the combination $1+\tilde{I}(q)$. Of course if we could show that

$$|\tilde{I}(q)| < 1, \tag{6.1}$$

for any $m \geq 2$ and all $q = 3, 5, \cdots, 4m-5$, then Theorem 2.6 would follow immediately from Proposition 6.1: unfortunately we are unable to prove (6.1) for all $q, m$ as above, and in fact it may not even be true for all $q, m$ in the range.

In Subsection 6.3, we prove a crude a priori bound on the absolute value of $\tilde{I}(q)$ (see (6.15) below) that is valid for all $m \geq 2$ and all $q = 3, 5, \cdots, 4m-5$. The proof of this bound follows from the hypergeometric ODE for $h(x)$ mentioned above (cf. (6.8) below).

In Subsection 6.4, we use a more refined argument to show that the absolute value of $\tilde{I}(q)$ is bounded by 1 for all $m \geq 2$ and for all $q$ in a region $O(\sqrt{m}) \leq q \leq O(m)$, see (6.21) below. To this end we use the properties of $y_m(\theta)$ (see (2.21)) that follow from the fact that it satisfies a certain Riccati equation (see Proposition 6.4): as noted in Remark 2.3 this Riccati equation in turn is a consequence of the hypergeometric ODE for $h(x)$ mentioned above.

Finally, in Subsection 6.5 we use a certain integral representation for $h(x)$ to find, for large enough $m$, an accurate approximation for $\tilde{I}(q)$ which yields a bound (see (6.57) below) on $|1+\tilde{I}(q)|$ for $q$ in the region $3 \leq q \leq O(\sqrt{m})$.

The above estimates are sufficient to prove $\det T_{m-1} \neq 0$ for all $m \geq 2$. The crude bound in Subsection 6.3 is sufficient to give a simple proof of the result for $m \leq 51$ (see Subsection 6.3). The bound in Subsection 6.4 for $O(\sqrt{m}) \leq q \leq O(m)$, together with the crude bound in Subsection 6.3 applied to $q$ in the range $3 \leq q \leq O(\sqrt{m})$, gives a proof of the result for $m \leq 99$ (see Subsection 6.4). Finally the bound in Subsection 6.5 for large $m$ and $3 \leq q \leq O(\sqrt{m})$, together with the bound in Subsection 6.4 for $O(\sqrt{m}) \leq q \leq O(m)$, proves $\det T_{m-1} \neq 0$ for $m \geq 38$ (see Subsection 6.5). In this way the estimates in Subsections 6.3, 6.4 and 6.5 cover the entire range $m \geq 2$, with overlap and some redundancy.

### 6.2. **Notation.**

Recall the definition of $h(x)$ in (2.8). We will repeatedly use the change of variable $x = \sin\theta$ in (2.11). We find

$$I(q) = \frac{1}{2m} + \frac{2}{m\pi}\sin\frac{q\pi}{2}\int_{-\pi/2}^{\pi/2}\cos(q\theta)\, y_m(\theta)\, d\theta \tag{6.2}$$

where $y_m$ is defined in (2.21), and as indicated above we introduce the notation

$$\tilde{I}(q) \equiv mI(q) - \frac{1}{2} = \frac{2}{\pi}\sin\frac{q\pi}{2}\int_{-\pi/2}^{\pi/2}\cos(q\theta)\, y_m(\theta)\, d\theta, \qquad q = 3, 5, \cdots 4m-5. \tag{6.3}$$

In (6.2) we have used the following, simple identities for $q$ in the above range,

$$\int_{-\pi/2}^{\pi/2}\cos(q\theta)\cos\theta\, d\theta = 0, \qquad \frac{2}{\pi}\sin\frac{q\pi}{2}\int_{-\pi/2}^{\pi/2}\frac{\cos(q\theta)}{\cos\theta}\, d\theta = 2. \tag{6.4}$$

Set

$$\gamma_m \equiv \frac{2(m!)^2}{m(2m)!}$$

and let $Y_{m-1} \equiv (Y_{jk})_{j,k=1}^{m-1}$ denote the very last matrix (of binomial coefficients) in (2.17). Set

$$X_{m-1} \equiv (X_{jk})_{j,k=1}^{m-1} \equiv \begin{pmatrix} 1 & 1 & \cdots & 1 \\ 1 & 1 & \cdots & 1 \\ \cdots & & & \\ 1 & 1 & \cdots & 1 \end{pmatrix} + \begin{pmatrix} \tilde{I}(n) & \tilde{I}(n+2) & \cdots & \tilde{I}(2n-3) \\ \tilde{I}(n-2) & \tilde{I}(n) & \cdots & \tilde{I}(2n-5) \\ \cdots & & & \\ \tilde{I}(3) & \tilde{I}(5) & \cdots & \tilde{I}(n) \end{pmatrix}.$$

Then (2.17) becomes

$$T_{m-1} = I_{m-1} - \gamma_m X_{m-1} Y_{m-1}. \tag{6.5}$$

We treat the second term as a perturbation and it turns out to be important to choose the norm $\|\cdot\|$ on $\mathbb{R}^{m-1}$ appropriately. We take $\|\cdot\|$ to be the maximum norm on $\mathbb{R}^{m-1}$, $\|x = (x_1, \cdots, x_{m-1})\| = \max_{1\le j\le m-1} |x_j|$. We show eventually that acting as an operator on $(\mathbb{R}^{m-1}, \|\cdot\|)$, the second term in (6.5) has norm $< 1$ for all $m$. This of course implies that $\det T_{m-1} \neq 0$ for all $m \geq 2$.

**Proposition 6.1.** *Assume that for some* $C > 0$, $\max_{j,k=1,\cdots,m-1} |X_{j,k}| \leq C$. *Then*

$$\big\|\gamma_m X_{m-1} Y_{m-1}\big\| \leq C\Big(\frac{1}{2} - \frac{(m!)^2}{m(2m)!} \cdot 2^{2m-2}\Big). \tag{6.6}$$

*Proof.* By the definition of the norm $\|\cdot\|$ on $\mathbb{R}^{m-1}$

$$\begin{aligned} \|X_{m-1}Y_{m-1}\| &\leq \max_{i=1,\cdots,m-1} \sum_{j=1}^{m-1} |(X_{m-1}Y_{m-1})_{ij}| \\ &\leq \max_{i=1,\cdots,m-1} \sum_{j,k=1}^{m-1} |X_{ik}|\,|Y_{kj}| \leq C \sum_{j,k=1}^{m-1} |Y_{kj}| \\ &= C \sum_{i=0}^{m-2} \sum_{l=0}^{m-2-i} \binom{2m-1}{l} = C \sum_{l=0}^{m-1} \binom{2m-1}{l} (m-l-1). \end{aligned}$$

Substituting $\sum_{l=0}^{m-1} \binom{2m-1}{l} = 2^{2m-2}$ and

$$\sum_{l=0}^{m-1} \binom{2m-1}{l} (m-l) = \frac{m}{2} \binom{2m-1}{m-1} + 2^{2m-3}$$

which follows from the identity

$$2\sum_{l=0}^{q} \binom{p}{l} l = p \sum_{l=0}^{q-1} \binom{p}{l} + q \binom{p}{q}, \qquad q \leq p, \tag{6.7}$$

with $q = m-1$, $p = 2m-1$, we arrive at (6.6). It remains to prove (6.7). Note that

$$\begin{aligned}\sum_{l=0}^{q}\binom{p}{l}l &= \sum_{l=1}^{q}\frac{p(p-1)\cdots(p-(l-2))}{(l-1)!}\,(p-(l-1))\\ &= \sum_{l=1}^{q}\binom{p}{l-1}(p-(l-1)) = p\sum_{l=0}^{q-1}\binom{p}{l} - \sum_{l=0}^{q-1}\binom{p}{l}l\end{aligned}$$

from which (6.7) follows. □

6.3. **ODE for** $h(x)$**.** The *final* estimate (6.16) in this Subsection is completely superceded by the *final* estimate (6.22) in the next Subsection, which gives the desired result for $m$ up till 99. However both (6.16) and (6.22) utilize the estimate (6.15) that is proved below in this Subsection.

**Proposition 6.2.** *For any* $m \in \mathbb{N}$ *the (even) function* $h(x)$ *satisfies the ODE*

$$x(x^2-1)h' + ((2m-1) - 2(m-1)x^2)h = 4m, \qquad 0 \le x \le 1 \tag{6.8}$$

*Also*

$$h(0) = \frac{4m}{2m-1}, \qquad h(1) = 4m \tag{6.9}$$

*which together with* (6.8) *implies*

$$h(x) = \frac{4m}{x\sqrt{1-x^2}}\int_x^1 \left(\frac{x}{t}\right)^{2m}\frac{dt}{\sqrt{1-t^2}}, \qquad 0 < x < 1. \tag{6.10}$$

*Finally, the function* $h(x)$ *is strictly increasing on* $[0,1]$, *and hence it is* $\ge \frac{4m}{2m-1} > 0$ *on* $[-1,1]$.

*Proof.* Assuming (6.8) and either of the conditions (6.9) the integral representation (6.10) can be found using the integrating factor method. The explicit formula (2.8) readily implies (6.9) after elementary algebraic computations, and also that $h(x)$ is strictly increasing on $[0,1]$.

To prove (6.8), note that from (2.8)

$$\beta_k = \frac{4m}{2m-1}\frac{(-m+1)_k}{(-m+3/2)_k}$$

where $(a)_k \equiv a(a+1)\cdots(a+k-1)$ denotes the Pochhammer symbol. As $(1)_k = k!$,

$$\begin{aligned}h(x) &= \frac{4m}{2m-1}\sum_{k=0}^{m-1}\frac{(1)_k(-m+1)_k}{(-m+3/2)_k}\frac{(x^2)^k}{k!}\\ &= \frac{4m}{2m-1}\,{}_2F_1(1,-m+1,-m+3/2;x^2)\end{aligned} \tag{6.11}$$

(the series in ${}_2F_1$ terminates at $k = m-1$). Thus $w(z) \equiv h(\sqrt{z})$ satisfies the hypergeometric equation

$$z(1-z)w'' + (c-(a+b+1)z)w' - abw = 0$$

with $a = 1$, $b = -m+1$, $c = -m+3/2$ which becomes

$$z(1-z)w'' + ((-m+3/2) + (m-3)z)w' + (m-1)w = 0. \tag{6.12}$$

In turn this equation can be rewritten as

$$\frac{d}{dz}\Big(z(z-1)w' + ((m-1/2)-(m-1)z)w\Big) = 0 \tag{6.13}$$

and so

$$z(z-1)w' + ((m-1/2)-(m-1)z)w = \text{const.} \tag{6.14}$$

Letting $z \to 1$ and using (6.9) we conclude that const $= 2m$. But (6.14) is just (6.8) for $w(z) = h(\sqrt{z})$. Of course (6.8) can also be proved by substituting (2.8) directly into (6.8). □

*Remark* 6.1. Differentiating (4.131) and using (6.8), we immediately obtain the proof of Lemma 4.8.

Now from (6.8) it follows that

$$\frac{1}{1-x^2}\Big(\frac{4m}{h(x)}-1\Big) = 2(m-1) - \frac{xh'(x)}{h(x)}$$

which together with (6.3) and (2.21) gives for $q = 3, 5, \cdots$

$$\begin{aligned}\tilde{I}(q) &= \frac{m}{4m}\frac{2}{\pi}\sin\frac{q\pi}{2}\int_{-1}^{1}\frac{\cos(q\arcsin x)}{1-x^2}\Big(\frac{4m}{h(x)}-1\Big)\,dx\\ &= \frac{1}{2\pi}\sin\frac{q\pi}{2}\int_{-1}^{1}\cos(q\arcsin x)\,\Big(2(m-1)-\frac{xh'(x)}{h(x)}\Big)\,dx.\end{aligned}$$

The first integral vanishes for odd $q \geq 3$ by (6.4), and estimating the second by the absolute value of the integrand (using $h' \geq 0$ on $[0,1]$ and the fact that $h$ is even) we obtain the a priori estimate

$$|\tilde{I}(q)| \leq \frac{1}{\pi}\int_0^1 \frac{xh'(x)}{h(x)}\,dx = \frac{1}{\pi}\int_0^1 x\Big(\log\frac{h(x)}{h(1)}\Big)'\,dx = \frac{1}{\pi}\int_0^1 \log\frac{4m}{h(x)}\,dx$$

by (6.9). Now we use the estimate

$$\frac{4m}{h(x)} \leq (2m-1) - 2(m-1)x^2, \qquad 0 \leq x \leq 1,$$

which follows from (6.8) since $h$, $h' \geq 0$, to find

$$|\tilde{I}(q)| \leq L(m) \tag{6.15}$$

where

$$L(m) \equiv \frac{\log(2m-1) - 2 + (t_m+1)\log(1+t_m^{-1}) - (t_m-1)\log(1-t_m^{-1})}{\pi}$$

and $t_m \equiv \sqrt{\frac{2m-1}{2m-2}}$. Using Maple, for example, it is straightforward to check that for $m = 2, 3, \cdots, 51$

$$\big(1+L(m)\big)\Big(\frac{1}{2} - \frac{(m!)^2}{m(2m)!}\cdot 2^{2m-2}\Big) < 1 \tag{6.16}$$

and hence by Proposition 6.1, $T_{m-1}$ is invertible and therefore $\det T_{m-1} \neq 0$ for $m \leq 51$.

*Remark* 6.2. By Stirling's formula, $\frac{(m!)^2}{m(2m)!}\cdot 2^{2m-2} \sim \frac{\sqrt{\pi}}{4\sqrt{m}}$, and so the LHS of (6.16) grows logarithmically as $m \to \infty$.

6.4. **Riccati equation for $y_m(\theta)$.** A consequence of the calculations in this Subsection is that

$$\|\gamma_m X_{m-1}Y_{m-1}\|_{(\mathbb{R}^{m-1},\|\cdot\|)\to(\mathbb{R}^{m-1},\|\cdot\|)} < 1$$

for $m$ up to (the larger value) 99. In place of Proposition 6.1 we will use the following result. The proof is immediate.

**Proposition 6.3.** *Assume that for some matrix $\tilde{X}$ with nonnegative elements we have for all $j,k = 1,\cdots,m-1$ that $|X_{j,k}| \le \tilde{X}_{j,k}$. Then*

$$\|\gamma_m X_{m-1}Y_{m-1}\| \le \gamma_m \max_{i=1,\cdots,m-1} \sum_{j,k=1}^{m-1} \tilde{X}_{ik}Y_{kj}. \tag{6.17}$$

The main goal in this Subsection is to prove the estimate (6.21) below on $\tilde{I}(q)$. This estimate is useful for $q$ in the range $O(\sqrt{m}) \le q \le O(m)$. The matrix $\tilde{X}$ in Proposition 6.3 is then constructed by "interpolating" between this estimate and the estimate (6.15) above, as indicated in (6.22) below. Estimate (6.21) will be derived from the Riccati equation (2.22) for $y_m(\theta) = \frac{m}{\cos\theta}\big(\frac{1}{h(\sin\theta)} - \frac{1}{4m} - \frac{\cos^2\theta}{2}\big)$ given in Proposition 6.4 below. Note from (2.8) that $y_m(\theta)$ is even.

*Remark* 6.3. The specific form of $y_m(\theta)$ has the following consequences for $\tilde{I}(q) = \frac{2}{\pi}\sin\frac{q\pi}{2}\int_{-\pi/2}^{\pi/2}\cos(q\theta)\,y_m(\theta)\,d\theta$. One can show that as $m\to\infty$, $\frac{1}{h(x)} \to \frac{1-x^2}{2} = \frac{\cos^2\theta}{2}$ (formally at least, divide (6.8) by $m$ and let $m\to\infty$). Hence for a fixed $q$, as $m\to\infty$, we expect $\tilde{I}(q) \to -\frac{1}{2}$ (recall (6.4)). On the other hand for odd $q = 2l+1$,

$$\cos(q\theta)\,y_m(\theta) = \cos(2l\theta)\,(\cos\theta\,y_m(\theta)) - \sin(2l\theta)\,(\sin\theta\,y_m(\theta))$$

and so $\tilde{I}(q)$ is a sum of Fourier coefficients of the real analytic $\pi$-periodic functions $(\cos\theta)\,y_m(\theta)$, $(\sin\theta)\,y_m(\theta)$, and so for a fixed $m$, $\tilde{I}(q)$ decays exponentially as $q\to\infty$ by the Paley–Wiener theorem. As discussed in Remark 2.3 above, this indicates that there should be a transition in the $(q,m)$-plane between these two kinds of behavior. And indeed, as we will see below, the transition region is in the range $q \sim \sqrt{m}$.

**Proposition 6.4.** *For any $m \ge 1$ the function $y_m(\theta)$ defined in* (2.21) *satisfies a Riccati differential equation* (2.22)

$$y_m' = \frac{4}{\sin\theta}\left(y_m + \frac{2m+1}{4}\cos\theta\right)\left(y_m + \frac{1}{2\cos\theta}\right).$$

*Also $y_m(0) = -1/2$ and $y_m(\pi/2) = 0$, see Fig. 1.*

*Proof.* This is a straightforward computation using the ODE (6.8) for $h(x)$. The interesting fact here is that the RHS in the Riccati equation has real roots which, moreover, have simple expressions. The proof that $y_m(0) = -1/2$ follows directly from (2.9) and the fact that $y_m(\pi/2) = 0$ follows from (2.9) together with L'Hôpital's rule. □

We will show next that $y_m$ is a unimodal nonpositive function on $[0,\pi/2]$, and that its minimum satisfies a certain bound (see Fig. 2).

**Proposition 6.5.** *For any $m \ge 2$ the function $y_m(\theta)$ on the interval $[0,\pi/2]$ strictly decreases from $y_m(0) = -1/2$ to the value $y_{m,\min}$ which is achieved at a unique point $\theta_{\min} \in (0,\pi/2)$, and then strictly increases to $0$ at the point $\theta = \pi/2$. The value*



FIGURE 1. Graphs of $y_m(\theta)$, $-\frac{2m+1}{4}\cos\theta$ for $m = 10$ and $-1/(2\cos\theta)$ (the latter for $0 \leq \theta \leq \frac{\pi}{2} - 0.1$)



FIGURE 2. Graphs of $y_m(\theta)$ for $m = 10$, 100, 1000 and $-\frac{1}{2\cos\theta}$ (the latter for $0 \leq \theta \leq \frac{\pi}{2} - 0.04$)

$y_{m,\min}$ *lies above the $y$-coordinate of the point of intersection of the curves* $-\frac{1}{2\cos\theta}$ *and* $-\frac{2m+1}{4}\cos\theta$*, see Fig. 1. Finally,* $y_m(\theta) \leq 0$ *and*

$$y_{m,\min} \geq -\frac{1}{2}\sqrt{m+1/2}. \tag{6.18}$$

*Proof.* It is helpful to refer to Fig. 1 in the following argument. Define $u(\theta) \equiv -\frac{1}{2\cos\theta}$ and $l_m(\theta) \equiv -\frac{2m+1}{4}\cos\theta$. Note that by (2.8) and (2.21) it follows that $y_m(\theta)$ is a smooth even function on $[-\pi/2, \pi/2]$. This implies $y_m'(0) = 0$. Also it is clear that for all $m \geq 2$, $y_m(0) = u(0) > l_m(0)$. Also $u'(0) = 0$ and $u''(0) = -1/2$.

Assume for a moment that

$$y_m''(0) = -\frac{1}{2}\frac{2m-1}{2m-3}. \tag{6.19}$$

Then $y_m''(0) < -1/2$ for $m \geq 2$. Hence for small $\theta > 0$, $y_m$ lies below $u$ and above $l_m$. It follows in particular as $y_m(\pi/2) = 0$ and $u(\theta) \to -\infty$ as $\theta \to \pi/2$, that $y_m$ must intersect $u$ at some point, $\theta_0$, say, in $(0, \pi/2)$.

It the argument that follows the signature table for the vector field

$$V(y_m, \theta) = \frac{4}{\sin\theta}(y_m - l_m(\theta))(y_m - u(\theta))$$

plays a crucial role. We have

- $V(y_m, \theta) > 0$ if $y_m$ lies above $l_m(\theta)$ and $u(\theta)$ or below $l_m(\theta)$ and $u(\theta)$
- $V(y_m, \theta) < 0$ if $y_m$ lies between $l_m(\theta)$ and $u(\theta)$.

Let $\tilde\theta \in (0, \pi/2)$ be the (unique) intersection point for $l_m(\theta)$ and $u(\theta)$, $l_m(\tilde\theta) = u(\tilde\theta) \equiv \tilde y$. From the signature table we see that $V(\tilde y, \theta) < 0$ for all $\theta \neq \tilde\theta$. Moreover if $y_m(\tilde\theta) = \tilde y$ then $y_m'(\tilde\theta) = V(y_m(\tilde\theta), \tilde\theta) = 0$. But $u'(\tilde\theta) < 0$ and $l_m'(\tilde\theta) > 0$, and so $y_m(\theta)$ lies between $l_m(\theta)$ and $u(\theta)$ for $\theta > \tilde\theta$, $\theta$ close to $\tilde\theta$. From the signature table it then follows that $y_m(\theta)$ decreases as $\theta$ crosses $\tilde\theta$. In other words, we see that if $y_m(\theta)$ crosses the level $\tilde y$, then it must decrease. As $y_m(0) = -1/2 > \tilde y$, and as $y_m(\pi/2) = 0$, it follows, therefore, that $y_m(\theta)$ cannot cross $\tilde y$. In particular, $0 < \theta_0 < \tilde\theta$. Now a similar argument shows that for $\theta > \theta_0$, $\theta$ close to $\theta_0$, $y_m(\theta)$ lies in the region above $l_m(\theta)$ and $u(\theta)$ (and increases across $\theta_0$). Suppose that $y_m(\theta)$ exits this region for the first time for some $\theta_1$, $\theta_0 < \theta_1 < \pi/2$. But then, again by the signature table, $y_m'(\theta) = V(y_m(\theta), \theta) > 0$ for $\theta_0 < \theta < \theta_1$, and hence $y_m(\theta_1) > y_m(\theta_0)$. In particular, as $u'(\theta) < 0$, it follows that $y_m(\theta)$ does not exit the region through the curve $u(\theta)$. Thus we must have $y_m(\theta_1) = l_m(\theta_1)$ and $\tilde\theta < \theta_1 < \pi/2$. But arguing as before we see that $y_m(\theta)$ must decrease as it crosses $\theta_1$: as $y_m(\pi/2) = 0$ there must be some point $\theta_2$, $\theta_1 < \theta_2 < \pi/2$ for which $y_m(\theta_2) = y_m(\theta_1)$ and $y_m'(\theta_2) \geq 0$. But $V(y_m(\theta_2), \theta_2) < 0$, by the signature table, which is a contradiction. The above arguments show that $y_m(\theta)$ crosses $u(\theta)$ at a unique point, $\theta_{\min} = \theta_0$, $0 < \theta_0 < \tilde\theta$: for $0 < \theta < \theta_0$, $y_m(\theta)$ lies between $l_m(\theta)$ and $u(\theta)$ and for $\theta_0 < \theta < \pi/2$, $y_m(\theta)$ lies above $l_m(\theta)$ and $u(\theta)$. Thus $y_m(\theta)$ is unimodal, decreasing for $0 < \theta < \theta_0$ and increasing for $\theta_0 < \theta < \pi/2$. Clearly $y_{\min} = y_m(\theta_{\min})$ lies above $\tilde y = u(\tilde\theta) = l_m(\tilde\theta) = -\frac{1}{2}\sqrt{m + 1/2}$. Note finally that for $0 < \theta \leq \theta_{\min}$ we certainly have $y_m(\theta) \leq u(\theta) < 0$. On the other hand as $y_m'(\theta) > 0$ for $\theta_{\min} < \theta < \pi/2$, and as $y_m(\pi/2) = 0$, it follows that we must have $y_m(\theta) < 0$ for all $\theta_{\min} < \theta < \pi/2$. Thus $y_m(\theta) \leq 0$ for all $0 \leq \theta \leq \pi/2$.

It remains to prove (6.19). Divide the ODE (2.22) by $\sin\theta$ to obtain

$$\frac{y_m'(\theta)}{\sin\theta} = 4\left(y_m + \frac{2m+1}{4}\cos\theta\right) \cdot \frac{y_m + \frac{1}{2\cos\theta}}{\sin^2\theta}. \tag{6.20}$$

Now as $y_m(0) = -1/2$,

$$\lim_{\theta\to 0} \frac{y_m(\theta) + \frac{1}{2\cos\theta}}{\sin^2\theta} = \lim_{\theta\to 0} \frac{y_m'(\theta) + \frac{\sin\theta}{2\cos^2\theta}}{2\sin\theta\cos\theta} = \frac{1}{2}\lim_{\theta\to 0}\frac{y_m'(\theta)}{\sin\theta} + \frac{1}{4}.$$

Letting $\theta \to 0$ on both sides of (6.20) we obtain $\lim_{\theta\to 0}\frac{y_m'(\theta)}{\sin\theta} = -\frac{1}{2}\frac{2m-1}{2m-3}$. But $y_m''(0) = \lim_{\theta\to 0}\frac{y_m'(\theta)}{\sin\theta}$, and hence (6.19) follows. This completes the proof of the Proposition. □

Now we integrate by parts to obtain

$$\begin{aligned}|\tilde{I}(q)| &= \Big|\frac{4}{\pi}\int_0^{\pi/2}\cos(q\theta)y_m(\theta)\,d\theta\Big| = \frac{4}{q\pi}\Big|\int_0^{\pi/2}\sin(q\theta)y_m'(\theta)\,d\theta\Big|\\ &\le \frac{4}{q\pi}\int_0^{\pi/2}\big|y_m'(\theta)\big|\,d\theta = \frac{4}{q\pi}\Big(\int_0^{\theta_{\min}}(-y_m'(\theta))\,d\theta + \int_{\theta_{\min}}^{\pi/2}y_m'(\theta)\,d\theta\Big)\end{aligned}$$

where we have used $y_m(\pi/2) = 0$ and also the unimodality of $y_m$ (Proposition 6.5). Hence in view of (6.18) for any $m \ge 2$ and any odd $q \ge 3$ we have

$$|\tilde{I}(q)| \le \frac{8}{q\pi}\big|y_{m,\min}\big| \le \frac{4}{q\pi}\sqrt{m+1/2}. \tag{6.21}$$

In particular, $|\tilde{I}(q)| < 1$ for odd $q$ such that $\big[\frac{4}{\pi}\sqrt{m+1/2}\big] + 1 \le q \le 4m-5$.

Now we can combine the estimates (6.15) and (6.21) at our discretion. We define the estimating Toeplitz matrix $\tilde{X}_{m-1} \equiv (\tilde{X}_{j,k})_{j,k=1}^{m-1}$ in Proposition 6.3 as follows. The diagonals are numbered by odd $q = 3, 5, \cdots, 4m-5$ from the lower left corner. Thus if $j, k = 1, 2, \cdots, m-1$ are the row and column indices starting from the usual upper left corner, then $q = 2(k-j) + (2m-1)$. Therefore, in view of (6.15), (6.21), set

$$\tilde{X}_{j,k} \equiv 1 + \min\Big(L(m), \frac{4}{(2(k-j)+(2m-1))\pi}\sqrt{m+1/2}\Big). \tag{6.22}$$

Also let $Y_{m-1} \equiv (Y_{j,k})_{j,k=1}^{m-1}$ where $Y_{j,k} = \binom{2m-1}{k-j}$ for $1 \le j \le k \le m-1$ and 0 otherwise, be the matrix of binomial coefficients as before. Then one can check by computer (only sums and products are involved) that the RHS in (6.17) is $< 1$ for $m = 2, 3, \cdots, 99$. Thus by Proposition 6.3, $T_{m-1} \equiv I_{m-1} - \gamma_m X_{m-1} Y_{m-1}$ is invertible for $m \le 99$, and hence $\det T_{m-1} \neq 0$ for $m \le 99$.

### 6.5. **Large $m$ asymptotics for $y_m(\theta)$ via an integral representation.**

This is the last and most technical computation. The main idea here is to find for large enough $m$ an approximation to the solution $y_m(\theta)$ of the Riccati equation in $\int_0^{\pi/2}\cos(q\theta)y_m(\theta)d\theta$ such that the integral is well approximated in the range $q = 3, 5, \cdots, O(\sqrt{m})$. We show that the estimate (6.57) below that follows from this approximation, together with (6.21) for $\big[\frac{4}{\pi}\sqrt{m+1/2}\big] + 1 \le q \le 4m-5$, gives the desired result for all $m \ge 38$, so that there is a large overlap with the region $m \le 99$ where the result of the preceding Subsection is valid.

For $\rho \ge 0$ set

$$G(\rho) \equiv \frac{\int_0^1 (1+\rho v)^{-m-1/2}\frac{v\,dv}{\sqrt{1-v}}}{\int_0^1 (1+\rho v)^{-m-1/2}\frac{dv}{\sqrt{1-v}}}. \tag{6.23}$$

The first result we need is the following integral representation for $y_m(\theta)$. Using $x = \sin\theta$,

$$\begin{aligned} y_m(\theta) &\equiv \frac{m}{\sqrt{1-x^2}}\Big(\frac{1}{h(x)} - \frac{1}{4m} - \frac{1-x^2}{2}\Big) \\ &= -\frac{\sqrt{1-x^2}}{2} - \frac{x^2}{2\sqrt{1-x^2}}(m-1)\frac{\int_x^1 \left(\frac{x}{t}\right)^{2m}\left(\left(\frac{t}{x}\right)^2-1\right)\frac{dt}{\sqrt{1-t^2}}}{\int_x^1 \left(\frac{x}{t}\right)^{2m}\frac{dt}{\sqrt{1-t^2}}} \qquad (6.24) \\ &= -\frac{\sqrt{1-x^2}}{2} - \frac{m-1}{2}\sqrt{1-x^2}\, G(\rho) \end{aligned}$$

where

$$\rho \equiv x^{-2} - 1 = \frac{\cos^2\theta}{\sin^2\theta}, \qquad x = \sin\theta. \tag{6.25}$$

We will explicitly indicate the dependence of $h(x)$ on $m$ as $h_m(x)$ when needed. To prove (6.24) we first note that for $m \geq 2$

$$h_m(x) = \frac{2m}{2m-1}(2 + x^2 h_{m-1}(x)), \qquad 0 \leq x \leq 1, \tag{6.26}$$

which follows from the explicit formula (2.8) and in turn leads (after adding and subtracting $\frac{2x^2}{2m-1}(m-1)h_m$) to

$$(1 + (2m-2)(1-x^2))h_m = 4m + 2x^2(mh_{m-1} - (m-1)h_m)$$

from which, dividing by $4mh_m$, we obtain

$$\frac{1}{h_m} - \frac{1}{4m} - \frac{1-x^2}{2} = -\frac{1-x^2}{2m} - \frac{x^2}{2m}\,\frac{mh_{m-1} - (m-1)h_m}{h_m}.$$

Together with (6.10), this implies the second equality in (6.24). The third equality in (6.24) follows after changing the variable $t = x\sqrt{1+\rho v}$ where $\rho$ is as in (6.25).

Now from (6.24) for $q = 3, 5, \cdots$, and (6.4),

$$\tilde{I}(q) = -\frac{m-1}{\pi} 2\sin\frac{q\pi}{2}\int_0^{\pi/2} \big(\cos(q\theta)\cos\theta\big)\, G(\rho(\theta))\, d\theta.$$

Make now a change of variable $\phi = \frac{\pi}{2} - \theta$ giving

$$\rho = \tan^2\phi, \qquad d\rho = \frac{2\sin\phi}{\cos^3\phi}d\phi, \qquad \cos\phi = \frac{1}{\sqrt{1+\rho}}, \qquad \sin\phi = \frac{\sqrt{\rho}}{\sqrt{1+\rho}} \tag{6.27}$$

to find

$$\tilde{I}(q) = -\frac{m-1}{\pi}\int_0^\infty \sin(q\phi(\rho))\, G(\rho)\, \frac{d\rho}{(1+\rho)^{3/2}}.$$

We split the integral into two,

$$\tilde{I}(q) = -\frac{m-1}{\pi}\Big(\int_0^{m^{-1/2}} + \int_{m^{-1/2}}^\infty\Big)\sin(q\phi(\rho))\, G(\rho)\, \frac{d\rho}{(1+\rho)^{3/2}}. \tag{6.28}$$

Assume that $m \geq 3$ (we only need the results below for $m \geq 38$).

**Lemma 6.6.** *We have*

$$-\frac{m-1}{\pi}\int_{m^{-1/2}}^\infty \frac{\sin(q\phi(\rho))\, G(\rho)}{(1+\rho)^{3/2}}\, d\rho = -\frac{m-1}{m-\frac{3}{2}}\frac{1}{\pi}\int_{m^{-1/2}}^\infty \frac{\sin(q\phi(\rho))}{\rho(1+\rho)^{3/2}}d\rho + E_1(m,q)$$

*where the error term* $E_1(m,q)$ *satisfies*

$$|E_1(m,q)| \leq \frac{A(m)+B(m)+C(m)}{Denom(m)}$$

*uniformly for all* $q = 3, 5, \cdots$, *where*

$$\begin{aligned} A(m) &\equiv \frac{7}{\pi\sqrt{2}} \frac{(m-1)\sqrt{m}}{(m-\frac{1}{2})(m-\frac{5}{2})} \\ B(m) &\equiv \frac{3}{\pi\sqrt{2}} \Big( \frac{1}{2} (\log m) \frac{1}{(1+\frac{1}{2}\frac{1}{\sqrt{m}})^{m-3/2}} + \Big(\frac{2}{3}\Big)^{m-\frac{1}{2}} \Big) \\ C(m) &\equiv \frac{3}{\sqrt{2}-1} \frac{1}{\pi} \frac{1}{(1+\frac{\sqrt{2}-1}{\sqrt{2}}\frac{1}{\sqrt{m}})^{m}} \\ Denom(m) &\equiv \frac{m-\frac{3}{2}}{m-\frac{1}{2}} \Big( 1 - e^{-\sqrt{m}\frac{1-\frac{1}{2m}}{1+\frac{1}{\sqrt{m}}}} \Big). \end{aligned} \tag{6.29}$$

*Proof.* We consider $\rho \geq m^{-1/2}$. We need the following elementary inequality:

$$e^{ay/(1+y)} \leq (1+y)^a \leq e^{ay}, \qquad a, y \geq 0. \tag{6.30}$$

Set

$$k \equiv m + 1/2. \tag{6.31}$$

Splitting the interval $[0,1]$ into two and integrating by parts we find

$$\begin{aligned} \int_0^1 (1+\rho v)^{-k} \frac{dv}{\sqrt{1-v}} =& \frac{1}{(k-1)\rho} - \frac{\sqrt{2}}{(k-1)\rho} \frac{1}{(1+\rho/2)^{k-1}} \\ &+ \frac{1}{(k-1)\rho} \int_0^{1/2} (1+\rho v)^{-k+1} \Big( \frac{1}{\sqrt{1-v}} \Big)' dv \\ &+ \int_{1/2}^1 (1+\rho v)^{-k} \frac{dv}{\sqrt{1-v}} \end{aligned} \tag{6.32}$$

and

$$\begin{aligned} \int_0^1 (1+\rho v)^{-k} \frac{v\,dv}{\sqrt{1-v}} =& \frac{1}{(k-1)(k-2)\rho^2} - \frac{1}{(k-1)\rho} \frac{1}{\sqrt{2}} \frac{1}{(1+\rho/2)^{k-1}} \\ &- \frac{1}{(k-1)(k-2)\rho^2} \frac{3}{\sqrt{2}} \frac{1}{(1+\rho/2)^{k-2}} \\ &+ \frac{1}{(k-1)(k-2)\rho^2} \int_0^{1/2} (1+\rho v)^{-k+2} \Big( \frac{v}{\sqrt{1-v}} \Big)'' dv \\ &+ \int_{1/2}^1 (1+\rho v)^{-k} \frac{v\,dv}{\sqrt{1-v}}. \end{aligned} \tag{6.33}$$

This gives the leading behavior of the ratio in $G(\rho)$ for $m$ (and $k$) large and motivates the definition

$$\Delta(\rho) \equiv G(\rho) - \frac{1}{(k-2)\rho} = \frac{(k-2)\rho \cdot (\text{RHS in (6.33)}) - (\text{RHS in (6.32)})}{(k-2)\rho \int_0^1 (1+\rho v)^{-k} \frac{dv}{\sqrt{1-v}}}$$

which yields

$$
\begin{aligned}
&\Big((k-2)\rho\int_0^1 (1+\rho v)^{-k}\,\frac{dv}{\sqrt{1-v}}\Big)\,\Delta(\rho)\\
&\qquad =\frac{1}{(k-1)\rho}-\frac{k-2}{k-1}\,\frac{1}{\sqrt{2}}\,\frac{1}{(1+\rho/2)^{k-1}}\\
&\qquad\quad -\frac{1}{(k-1)\rho}\,\frac{3}{\sqrt{2}}\,\frac{1}{(1+\rho/2)^{k-2}}\\
&\qquad\quad +\frac{1}{(k-1)\rho}\int_0^{1/2}(1+\rho v)^{-k+2}\Big(\frac{v}{\sqrt{1-v}}\Big)''\,dv\\
&\qquad\quad +(k-2)\rho\int_{1/2}^1 (1+\rho v)^{-k}\,\frac{v\,dv}{\sqrt{1-v}}\\
&\qquad\quad -\frac{1}{(k-1)\rho}+\frac{\sqrt{2}}{(k-1)\rho}\,\frac{1}{(1+\rho/2)^{k-1}}\\
&\qquad\quad -\frac{1}{(k-1)\rho}\int_0^{1/2}(1+\rho v)^{-k+1}\Big(\frac{1}{\sqrt{1-v}}\Big)'\,dv\\
&\qquad\quad -\int_{1/2}^1 (1+\rho v)^{-k}\,\frac{dv}{\sqrt{1-v}}.
\end{aligned}
\tag{6.34}
$$

From this we derive an upper and a lower estimate on $\Delta(\rho)$. Discarding the negative terms, we obtain

$$
\begin{aligned}
&\Big((k-2)\rho\int_0^1 (1+\rho v)^{-k}\,\frac{dv}{\sqrt{1-v}}\Big)\,\Delta(\rho)\\
&\qquad \le\frac{1}{(k-1)\rho}\int_0^{1/2}(1+\rho v)^{-k+2}\Big(\frac{v}{\sqrt{1-v}}\Big)''\,dv\\
&\qquad\quad +(k-2)\rho\int_{1/2}^1 (1+\rho v)^{-k}\,\frac{v\,dv}{\sqrt{1-v}}\\
&\qquad\quad +\frac{\sqrt{2}}{(k-1)\rho}\,\frac{1}{(1+\rho/2)^{k-1}}.
\end{aligned}
\tag{6.35}
$$

Note that the second derivative above is indeed nonnegative and that its maximum over $[0,1/2]$ equals $7/\sqrt{2}$. Next changing variables $\sqrt{1-v}=t$ and integrating by parts we obtain

$$
\begin{aligned}
\int_{1/2}^1 (1+\rho v)^{-k}\,\frac{v\,dv}{\sqrt{1-v}} &= 2\int_0^{1/\sqrt{2}}\frac{1-t^2}{(1+\rho(1-t^2))^k}\,dt \le 2\int_0^{1/\sqrt{2}}\frac{1-t^2}{(1+\rho(1-t))^k}\,dt\\
&\le 2\frac{(1+\rho(1-1/\sqrt{2}))^{-k+1}}{(k-1)\rho}\frac{1}{2}-\frac{2}{(k-1)\rho}\int_0^{1/\sqrt{2}}\frac{-2t}{(1+\rho(1-t))^{k-1}}\,dt\\
&\le \frac{1}{(k-1)\rho}\,\frac{2}{(1+\rho(1-1/\sqrt{2}))^{k-1}}\Big(\frac{1}{2}+\int_0^{1/\sqrt{2}}2t\,dt\Big).
\end{aligned}
$$

Thus from (6.35), (6.31)

$$\Big((m-3/2)\rho\int_0^1(1+\rho v)^{-m-1/2}\,\frac{dv}{\sqrt{1-v}}\Big)\,\Delta(\rho)\le\frac{7}{\sqrt{2}}\frac{1}{(m-1/2)(m-5/2)\rho^2}+\frac{2}{(1+(1-1/\sqrt{2})\rho)^{m-1/2}}+\frac{\sqrt{2}}{(m-1/2)\rho}\,\frac{1}{(1+\rho/2)^{m-1/2}}. \tag{6.36}$$

Next discarding the positive terms in (6.34), we find

$$\begin{aligned}&-\Big((k-2)\rho\int_0^1(1+\rho v)^{-k}\,\frac{dv}{\sqrt{1-v}}\Big)\,\Delta(\rho)\\&\quad\le\frac{k-2}{k-1}\,\frac{1}{\sqrt{2}}\,\frac{1}{(1+\rho/2)^{k-1}}+\frac{1}{(k-1)\rho}\,\frac{3}{\sqrt{2}}\,\frac{1}{(1+\rho/2)^{k-2}}\\&\quad+\frac{1}{(k-1)\rho}\int_0^{1/2}(1+\rho v)^{-k+1}\Big(\frac{1}{\sqrt{1-v}}\Big)'\,dv+\int_{1/2}^1(1+\rho v)^{-k}\,\frac{dv}{\sqrt{1-v}}.\end{aligned} \tag{6.37}$$

Note that the first derivative above is indeed nonegative and that its maximum over $[0,1]$ equals $\sqrt{2}$. Note finally that $\int_{1/2}^1(1+\rho v)^{-k}\,\frac{dv}{\sqrt{1-v}}\le\frac{1}{(1+\rho/2)^k}\int_{1/2}^1\frac{dv}{\sqrt{1-v}}=\frac{\sqrt{2}}{(1+\rho/2)^k}$. Thus (6.37), (6.31) give

$$\begin{aligned}&-\Big((m-3/2)\rho\int_0^1(1+\rho v)^{-m-1/2}\,\frac{dv}{\sqrt{1-v}}\Big)\,\Delta(\rho)\\&\quad\le\frac{1}{\sqrt{2}}\,\frac{m-3/2}{m-1/2}\,\frac{1}{(1+\rho/2)^{m-1/2}}+\frac{3}{\sqrt{2}}\,\frac{1}{(m-1/2)\rho}\,\frac{1}{(1+\rho/2)^{m-3/2}}\\&\quad+\frac{\sqrt{2}}{(m-1/2)(m-3/2)\rho^2}+\frac{\sqrt{2}}{(1+\rho/2)^{m+1/2}}.\end{aligned} \tag{6.38}$$

Now using (6.30), we see that the factor in front of $\Delta(\rho)$ in (6.36) satisfies

$$\begin{aligned}\int_0^1(1+\rho v)^{-k}\,\frac{dv}{\sqrt{1-v}}&\ge\int_0^1(1+\rho v)^{-k}\,dv=\frac{1}{(k-1)\rho}\Big(1-\frac{1}{(1+\rho)^{k-1}}\Big)\\&\ge\frac{1}{(k-1)\rho}\Big(1-e^{-(k-1)\frac{\rho}{1+\rho}}\Big)\ge\frac{1}{(k-1)\rho}\Big(1-e^{-(k-1)\frac{m^{-1/2}}{1+m^{-1/2}}}\Big)\\&\ge\frac{1}{(m-1/2)\rho}\Big(1-e^{-(1-\frac{1}{2m})\frac{\sqrt{m}}{1+\frac{1}{\sqrt{m}}}}\Big),\end{aligned} \tag{6.39}$$

by (6.31) since $-\frac{\rho}{1+\rho}$ decreases and $m^{-1/2}\le\rho$.

We now consider the various terms in (6.36), (6.38) to obtain an estimate on $|\Delta(\rho)|$, $\rho\ge m^{1/2}$. Note that:

- the 1st term in (6.36) dominates the 3rd term in (6.38);
- the 2nd term in (6.38) dominates the 3rd term in (6.36);
- both the 2nd term in (6.36) and the sum of the 1st and 4th terms in (6.38) are dominated by $\frac{3/\sqrt{2}}{(1+(1-1/\sqrt{2})\rho)^{m-1/2}}$.

Thus taking into account (6.39) and recalling the notation (6.29) we conclude

(6.40)

$$|\Delta(\rho)| \leq \frac{1}{Denom(m)} \Big( \frac{7}{\sqrt{2}} \frac{1}{(m-1/2)(m-5/2)\rho^2} + \frac{3}{\sqrt{2}} \frac{1}{(m-1/2)\rho} \frac{1}{(1+\rho/2)^{m-3/2}} + \frac{3}{\sqrt{2}} \frac{1}{(1+(1-1/\sqrt{2})\rho)^{m-1/2}} \Big)$$

for all $m^{-1/2} \leq \rho < \infty$.

Now substituting $G(\rho) = \frac{1}{(m-3/2)\rho} + \Delta(\rho)$ into the integral $\int_{m^{-1/2}}^{\infty}$ in (6.28), we obtain

$$-\frac{m-1}{\pi} \int_{m^{-1/2}}^{\infty} \frac{\sin(q\phi(\rho))\, G(\rho)}{(1+\rho)^{3/2}}\, d\rho = -\frac{m-1}{m-3/2} \frac{1}{\pi} \int_{m^{-1/2}}^{\infty} \frac{\sin(q\phi(\rho))}{\rho(1+\rho)^{3/2}} d\rho + E_1(m,q)$$

where

(6.41)

$$|E_1(m;q)| \leq \frac{1}{Denom(m)} \frac{m-1}{\pi} \Big( \frac{7}{\sqrt{2}} \frac{1}{(m-1/2)(m-5/2)} \int_{m^{-1/2}}^{\infty} \frac{d\rho}{\rho^2(1+\rho)^{3/2}} + \frac{3}{\sqrt{2}} \frac{1}{(m-1/2)} \int_{m^{-1/2}}^{\infty} \frac{d\rho}{\rho(1+\rho/2)^{m-3/2}(1+\rho)^{3/2}} + \frac{3}{\sqrt{2}} \int_{m^{-1/2}}^{\infty} \frac{d\rho}{(1+(1-1/\sqrt{2})\rho)^{m-1/2}(1+\rho)^{3/2}} \Big)$$

After elementary manipulations (splitting the integral as below) the first term gives $A(m)$ in (6.29). To estimate the second term, note that

$$\Big( \int_{m^{-1/2}}^{1} + \int_{1}^{\infty} \Big) \frac{d\rho}{\rho(1+\rho/2)^{m-3/2}(1+\rho)^{3/2}} \leq \frac{1}{(1+1/(2\sqrt{m}))^{m-3/2}} \int_{m^{-1/2}}^{1} \frac{d\rho}{\rho} + \frac{1}{(3/2)^{m-3/2}} \int_{1}^{\infty} \frac{d\rho}{\rho^{5/2}}$$

which gives $B(m)$ in (6.29). Finally in the third term in (6.41) we use

$$\int_{m^{-1/2}}^{\infty} \frac{d\rho}{(1+(1-1/\sqrt{2})\rho)^{m-1/2}(1+\rho)^{3/2}} \leq \int_{m^{-1/2}}^{\infty} \frac{d\rho}{(1+(1-1/\sqrt{2})\rho)^{m+1}}$$

and arrive at $C(m)$ in (6.29). This completes the proof of Lemma 6.6. ☐

In the integral $\int_0^{m^{-1/2}}$ in (6.28) we approximate the function $G(\rho)$ by the function

$$G_a(\rho) \equiv \frac{\int_0^1 e^{-\rho(m+1/2)v} \frac{v dv}{\sqrt{1-v}}}{\int_0^1 e^{-\rho(m+1/2)v} \frac{dv}{\sqrt{1-v}}}. \tag{6.42}$$

and use the following result.

**Lemma 6.7.** *For $m \geq 38$, we have*

(6.43)

$$-\frac{m-1}{\pi} \int_0^{m^{-1/2}} \frac{\sin(q\phi(\rho))\, G(\rho)}{(1+\rho)^{3/2}}\, d\rho = -\frac{m-1}{\pi} \int_0^{m^{-1/2}} \frac{\sin(q\phi(\rho))}{(1+\rho)^{3/2}}\, G_a(\rho)\, d\rho + E_2(m,q)$$

*where the error term $E_2(m,q)$ satisfies, uniformly for all $q = 3, 5, \cdots$,*

$$|E_2(m,q)| \leq \frac{1}{\pi} \frac{1}{\sqrt{m}} \cdot 6,$$

*where the number* 6 *provides the following (crude) bound (proved in the Appendix)*

$$\max_{m\geq 38}\max_{s\geq 0} s^2 \frac{\int_0^1 e^{-svR(m)}\,\frac{v^2\,dv}{\sqrt{1-v}}}{\int_0^1 e^{-sv}\,\frac{dv}{\sqrt{1-v}}} \leq 6, \tag{6.44}$$

*where*

$$R(m) \equiv \frac{1}{1+\frac{1}{\sqrt{m}}}. \tag{6.45}$$

*As* $R(m)$ *increases with* $m$*, the maximum on the LHS of* (6.44) *is achieved at* $m = 38$.

*Proof.* We consider $0 \leq \rho \leq m^{-1/2}$. Recall the notation (6.31) and denote

$$\Delta_k(s) \equiv s^2 \frac{\int_0^1 e^{-svR(m)}\,\frac{v^2\,dv}{\sqrt{1-v}}}{\int_0^1 e^{-sv}\,\frac{dv}{\sqrt{1-v}}}.$$

Assume for a moment the estimates

$$0 \leq G(\rho) - G_a(\rho) \leq \frac{\Delta_k(k\rho)}{k}, \qquad 0 \leq \rho \leq m^{-1/2}. \tag{6.46}$$

Then (6.43) holds where $E_2(m,q)$ satisfies

$$\begin{aligned} |E_2(m,q)| &\leq \frac{m-1}{\pi}\frac{1}{k}\int_0^{m^{-1/2}} \Delta_k(k\rho)\,\frac{d\rho}{(1+\rho)^{3/2}} \\ &\leq \Big(\max_{s\geq 0}\Delta_k(s)\Big)\frac{1}{\pi}\int_0^{m^{-1/2}} \frac{d\rho}{(1+\rho)^{3/2}} \leq 6\cdot\frac{1}{\pi}\cdot\frac{1}{\sqrt{m}} \end{aligned}$$

uniformly for all $m \geq 38$ and all $q = 3, 5, \cdots$. This is the desired result.

It remains to prove (6.46). First consider the upper bound. In view of (6.23), (6.42), (6.30)

$$G(\rho) \leq \frac{\int_0^1 e^{-k\rho v/(1+\rho)}\,\frac{vdv}{\sqrt{1-v}}}{\int_0^1 e^{-k\rho v}\,\frac{dv}{\sqrt{1-v}}} = G_a(\rho) + \frac{\int_0^1 \left(e^{-k\rho v/(1+\rho)} - e^{-k\rho v}\right)\frac{vdv}{\sqrt{1-v}}}{\int_0^1 e^{-k\rho v}\,\frac{dv}{\sqrt{1-v}}}.$$

Next, for all $0 \leq v \leq 1$

$$\begin{aligned} e^{-k\rho v/(1+\rho)} - e^{-k\rho v} &= \int_0^1 \frac{d}{dt}\Big(e^{-k\rho v(1-t+t/(1+\rho))}\Big)\,dt = \frac{k\rho^2 v}{1+\rho}\int_0^1 e^{-k\rho v(1-t+t/(1+\rho))}\,dt \\ &\leq \frac{k\rho^2 v}{1+\rho}e^{-k\rho v/(1+\rho)}\int_0^1 dt \leq k\rho^2 v\, e^{-k\rho v R(m)} \end{aligned}$$

where the exponent is estimated by its maximal value at $t = 1$ and we use the fact that $-\frac{1}{1+\rho} \leq -R(m)$ for $0 \leq \rho \leq \frac{1}{\sqrt{m}}$ by the definition (6.45). This proves the upper bound in (6.46).

For the lower bound in (6.46) we note

$$
\begin{aligned}
\big(G(\rho) - G_a(\rho)\big) &\cdot \Big(\int_0^1 (1+\rho v)^{-k}\,\frac{dv}{\sqrt{1-v}}\Big) \cdot \Big(\int_0^1 e^{-k\rho v}\,\frac{dv}{\sqrt{1-v}}\Big) \\
&= \int_0^1\int_0^1 (1+\rho v)^{-k} e^{-k\rho w}\,(v-w)\,\frac{dv}{\sqrt{1-v}}\,\frac{dw}{\sqrt{1-w}} \\
&= \frac{1}{2}\int_0^1\int_0^1 \Big((1+\rho v)^{-k} e^{-k\rho w} - (1+\rho w)^{-k} e^{-k\rho v}\Big)\,(v-w)\,\frac{dv}{\sqrt{1-v}}\,\frac{dw}{\sqrt{1-w}} \\
&= \frac{1}{2}\int_0^1\int_0^1 (v-w)\,(1+\rho v)^{-k}\,e^{-k\rho v}\Big[e^{k\rho(v-w)} - \Big(\frac{1+\rho v}{1+\rho w}\Big)^k\Big]\,\frac{dv}{\sqrt{1-v}}\,\frac{dw}{\sqrt{1-w}}.
\end{aligned}
$$

Consider $f(x) \equiv x^k$, $k \geq 1$. Note that for $a, b > 0$, $f(a) - f(b) = f'(\theta(a,b))(a-b)$ for some $\theta(a,b)$ between $a$ and $b$. As $\theta(a,b) > 0$, we must have $f'(\theta(a,b)) > 0$. Thus

$$
\begin{aligned}
\big(G(\rho) - G_a(\rho)\big) &\cdot \Big(\int_0^1 (1+\rho v)^{-k}\,\frac{dv}{\sqrt{1-v}}\Big) \cdot \Big(\int_0^1 e^{-k\rho v}\,\frac{dv}{\sqrt{1-v}}\Big) \\
&= \frac{1}{2}\int_0^1\int_0^1 (v-w)\Big(e^{\rho(v-w)} - \frac{1+\rho v}{1+\rho w}\Big)\,F_{k,\rho}(v,w)\,\frac{dv}{\sqrt{1-v}}\,\frac{dw}{\sqrt{1-w}}
\end{aligned}
\tag{6.47}
$$

where $F_{k,\rho}(v,w) \geq 0$ everywhere. Consider now $f_w(v) \equiv e^{\rho(v-w)} - \frac{1+\rho v}{1+\rho w}$. Then $f_w(w) = 0$ and $f_w'(v) = \rho\big(e^{\rho(v-w)} - \frac{1}{1+\rho w}\big) > 0$ for $v > w \geq 0$ (note $\rho > 0$). Thus $f_w(v) > 0$ for $v > w$. Also $f_w(v) = e^{\rho(v-w)}\frac{1+\rho v}{1+\rho w}\big(\frac{1+\rho w}{1+\rho v} - e^{\rho(w-v)}\big)$ which is $< 0$ by the same argument for $0 \leq v < w$. We conclude that the indegrand in (6.47) is $\geq 0$ for all $0 \leq v$, $w \leq 1$ which proves the lower bound in (6.46). The proof of Lemma 6.7 is now complete. ☐

The integral that appears on the RHS in (6.43), may be written in the form

$$
\begin{aligned}
-\frac{m-1}{\pi}&\int_0^{m^{-1/2}} \frac{\sin(q\phi(\rho))}{(1+\rho)^{3/2}}\,G_a(\rho)\,d\rho \\
&= -\frac{m-1}{m+\frac{1}{2}}\frac{1}{\pi}\int_0^{m^{-1/2}} \frac{\sin(q\phi(\rho))}{\rho(1+\rho)^{3/2}}\,\Big((m+1/2)\rho G_a(\rho) - 1\Big)\,d\rho \\
&\quad - \frac{m-1}{m+\frac{1}{2}}\frac{1}{\pi}\int_0^{m^{-1/2}} \frac{\sin(q\phi(\rho))}{\rho(1+\rho)^{3/2}}\,d\rho
\end{aligned}
\tag{6.48}
$$

The second integral on the right in (6.48) equals

$$
-\frac{m-1}{m+\frac{1}{2}}\frac{1}{\pi}\int_0^{m^{-1/2}} \frac{\sin(q\phi(\rho))}{\rho(1+\rho)^{3/2}}\,d\rho = -\frac{m-1}{m-\frac{3}{2}}\frac{1}{\pi}\int_0^{m^{-1/2}} \frac{\sin(q\phi(\rho))}{\rho(1+\rho)^{3/2}}\,d\rho + E_3(m;q)
$$

where

$$\begin{aligned}
|E_3(m;q)| \equiv & \frac{2}{\pi}\frac{m-1}{(m+\frac{1}{2})(m-\frac{3}{2})}\int_0^{m^{-1/2}} \frac{|\sin(q\phi(\rho))|}{\rho(1+\rho)^{3/2}}\,d\rho \\
\leq & \frac{2}{\pi}\frac{q}{m+\frac{1}{2}}\frac{m-1}{m-\frac{3}{2}}\int_0^{m^{-1/2}} \frac{|\phi(\rho)|}{\rho(1+\rho)^{3/2}}\,d\rho \\
\leq & \frac{2}{\pi}\frac{q}{m+\frac{1}{2}}\frac{m-1}{m-\frac{3}{2}}\int_0^{m^{-1/2}} \frac{\sqrt{\rho}}{\rho(1+\rho)^{3/2}}\,d\rho \\
\leq & \frac{2}{\pi}\frac{q}{m+\frac{1}{2}}\frac{m-1}{m-\frac{3}{2}}\int_0^{m^{-1/2}} \frac{d\rho}{\sqrt{\rho}} \\
\leq & \frac{4}{\pi}\frac{q}{\sqrt{m+\frac{1}{2}}}\frac{m-1}{m-\frac{3}{2}}\frac{1}{m^{3/4}}
\end{aligned} \tag{6.49}$$

and we have used

$$\phi(\rho) \leq \tan(\phi(\rho)) = \sqrt{\rho} \tag{6.50}$$

by (6.27). Denote the first integral on the right in (6.48) by $E_4(m;q)$ and set

$$F(t) \equiv t\,\frac{\int_0^1 e^{-tv}\,\frac{v\,dv}{\sqrt{1-v}}}{\int_0^1 e^{-tv}\,\frac{dv}{\sqrt{1-v}}} - 1.$$

Then $(m+1/2)\rho G_a(\rho) - 1 = F((m+1/2)\rho)$ and

$$\begin{aligned}
|E_4(m;q)| = & \left| -\frac{m-1}{m+\frac{1}{2}}\frac{1}{\pi}\int_0^{m^{-1/2}} \frac{\sin(q\phi(\rho))}{\rho(1+\rho)^{3/2}}\,F((m+1/2)\rho)\,d\rho \right| \\
\leq & \frac{m-1}{m+\frac{1}{2}}\frac{1}{\pi}\int_0^{m^{-1/2}} \frac{q\phi(\rho)}{\rho(1+\rho)^{3/2}}\,|F((m+1/2)\rho)|\,d\rho \\
\leq & \frac{1}{\pi}q\int_0^{m^{-1/2}} |F((m+1/2)\rho)|\,\frac{d\rho}{\sqrt{\rho}} \\
\leq & \frac{1}{\pi}\frac{q}{\sqrt{m+\frac{1}{2}}}\int_0^{+\infty} |F(w)|\,\frac{dw}{\sqrt{w}} \\
\leq & \frac{1}{\pi}\frac{q}{\sqrt{m+\frac{1}{2}}}\int_0^{+\infty} 2|F(s^2)|\,ds
\end{aligned}$$

where we have again used (6.50). Now we denote

$$H(s) \equiv 2F(s^2) = 2\left(s^2\,\frac{\int_0^1 e^{-s^2 v}\,\frac{v\,dv}{\sqrt{1-v}}}{\int_0^1 e^{-s^2 v}\,\frac{dv}{\sqrt{1-v}}} - 1\right)$$

and refer to the (crude) bound proved in the Appendix

$$\int_0^{+\infty} |H(s)|\,ds \leq 2.8 \tag{6.51}$$

to conclude that

$$|E_4(m;q)| \le \frac{1}{\pi}\frac{q}{\sqrt{m+\frac{1}{2}}}\cdot 2.8. \tag{6.52}$$

Combining Lemma 6.6 and 6.7 we find

$$\tilde{I}(q) = -\frac{m-1}{m-\frac{3}{2}}\frac{1}{\pi}\int_0^\infty \frac{\sin(q\phi(\rho))}{\rho(1+\rho)^{3/2}}d\rho + \sum_{j=1}^4 E_j(m;q). \tag{6.53}$$

Returning to the variable $\theta = \frac{\pi}{2} - \phi(\rho)$ and using (6.27) we see that for any $q = 3, 5, \cdots$

$$\frac{1}{\pi}\int_0^\infty \frac{\sin(q\phi(\rho))}{\rho(1+\rho)^{3/2}}d\rho = \frac{2}{\pi}\int_0^{\pi/2}\frac{\sin(q\theta)}{\sin\theta}\,d\theta = 1.$$

Hence (6.53) becomes

$$\tilde{I}(q) = -1 + \sum_{j=1}^5 E_j(m;q) \tag{6.54}$$

where

$$E_5(m;q) \equiv -\frac{1}{2}\frac{1}{m-\frac{3}{2}}, \qquad |E_5(m;q)| \le \frac{1}{2}\frac{1}{m-\frac{3}{2}}.$$

Recall that we need a bound on $1+\tilde{I}(q)$. Assembling the above estimates for $m \ge 38$, $q = 3, 5, \cdots, 4m-5$,

$$\begin{aligned} |1+\tilde{I}(q)| &= \Big|\sum_{j=1}^5 E_j(m;q)\Big| \le \sum_{j=1}^5 |E_j(m;q)| \\ &\le \frac{A(m)+B(m)+C(m)}{Denom(m)} + \frac{1}{\sqrt{m}}\frac{6}{\pi} + \frac{1}{2}\frac{1}{m-\frac{3}{2}} \\ &\quad + \frac{q}{\sqrt{m+1/2}}\left(\frac{2.8}{\pi} + \frac{4}{\pi}\frac{m-1}{m-\frac{3}{2}}\frac{1}{m^{3/4}}\right). \end{aligned} \tag{6.55}$$

We can now complete the proof of Theorem 2.6 for the case of large $m$. We wish to apply Proposition 6.1. We estimate the elements of the matrix $X_{m-1}$ as follows. Enumerate the diagonals of $X_{m-1}$ by $q = 3, 5, \cdots, 4m-5$ starting from the lower left corner as before. For a given *fixed* $m$, and *large* odd $q$, that is odd $q$ which satisfy the condition $q \ge \left[\frac{4}{\pi}\sqrt{m+1/2}\right] + 1$, we estimate the elements on the corresponding diagonal of $X_{m-1}$ as follows in view of (6.21)

$$|1+\tilde{I}(q)| \le 1 + |\tilde{I}(q)| \le 1 + \frac{\frac{4}{\pi}\sqrt{m+1/2}}{\left[\frac{4}{\pi}\sqrt{m+1/2}\right]+1} \equiv C_1(m) < 2. \tag{6.56}$$

But for the elements in $X_{m-1}$ on the diagonals corresponding to *small* odd $q$, that is for $3 \le q \le \left[\frac{4}{\pi}\sqrt{m+1/2}\right]$, we have $\frac{q}{\sqrt{m+1/2}} \le \frac{4}{\pi}$, and for such $q$ we find from (6.55)

$$\begin{aligned} |1+\tilde{I}(q)| &\le \frac{A(m)+B(m)+C(m)}{Denom(m)} + \frac{1}{\sqrt{m}}\frac{6}{\pi} + \frac{1}{2}\frac{1}{m-\frac{3}{2}} \\ &\quad + \frac{4}{\pi}\left(\frac{2.8}{\pi} + \frac{4}{\pi}\frac{m-1}{m-\frac{3}{2}}\frac{1}{m^{3/4}}\right) \equiv C_2(m). \end{aligned} \tag{6.57}$$

Thus for $m \geq 38$ and for $q = 3, 5, \cdots, 4m-5$ we have

$$|1 + \tilde{I}(q)| \leq C(m)$$

where $C(m) = \max(C_1(m), C_2(m))$. Now it is elementary to show that for $m \geq 58$, all the terms on the RHS of (6.55) decrease monotonically with $m$ and direct computation shows that $C_2(58) < 1.997 < 2$. Together with (6.56) this implies $C(m) < 2$ for all $m \geq 58$ and hence $\det T_m \neq 0$ by Proposition 6.1 for $m$ in this range.

But direct evaluation shows that for $38 \leq m \leq 57$

$$C_2(m) \left( \frac{1}{2} - \frac{(m!)^2}{m(2m)!} \cdot 2^{2m-2} \right) < 0.996 < 1$$

and so, again using (6.56), we have

$$C(m) \left( \frac{1}{2} - \frac{(m!)^2}{m(2m)!} \cdot 2^{2m-2} \right) < 1, \qquad 38 \leq m \leq 57.$$

Hence $\det T_{m-1} \neq 0$ also for $38 \leq m \leq 57$. This completes the proof of Theorem 2.6.

## Appendix

Our goal here is to prove the two (crude) estimates, (6.44) and (6.51) needed in the text. For $s \in [0, +\infty)$, set

$$L(s) \equiv s^2 \frac{\int_0^1 e^{-R_0 s v} \frac{v^2\, dv}{\sqrt{1-v}}}{\int_0^1 e^{-sv} \frac{dv}{\sqrt{1-v}}}$$

where $R_0 \equiv 0.855$. Note that $R(m) > R_0$ for all $m \geq 38$. Set also

$$L_1(s) \equiv s \frac{\int_0^1 e^{-s^2 v} \frac{v\, dv}{\sqrt{1-v}}}{\int_0^1 e^{-s^2 v} \frac{dv}{\sqrt{1-v}}} \tag{A.1}$$

and

$$H(s) \equiv 2\big(sL_1(s) - 1\big).$$

**Proposition A.1.** *For $s \in [0, +\infty)$*

$$0 \leq L(s) \leq 6.$$

**Proposition A.2.** *We have*

$$\int_0^{+\infty} |H(s)|\, ds \leq 2.8.$$

*The function $H(s)$ changes sign.*

The functions $L(s)$ and $H(s)$ are plotted in Figs. 3 and 4 respectively, using Maple. Fig. 3 "proves" Proposition A.1. Integrating by parts we will prove below that for all $s > 0$

$$H(s) \leq \frac{1}{s^2} + \frac{30\sqrt{2}}{s^4} + 2\sqrt{2} s^4 e^{-s^2/2} \tag{A.2}$$

which together with the fact ("evident" from Fig. 4 and proved below) that $H$ is positive on $[6, +\infty)$ implies $\int_6^\infty |H(s)| ds \leq 0.233$. Again, using Maple, direct numerical integration shows that $\int_0^6 |H(s)| ds \leq 2.407$. This "proves" Proposition A.2 with 2.64 on the right. Given the importance of the numerical values of the



FIGURE 3. Graph of $L(s)$ and the line at height 6



FIGURE 4. Graph of $H(s)$

bounds (6.44) and (6.51) in our Proof of Theorem 2.6, we now give a rigorous error analysis (in which we use the computer to the extent of computing the four arithmetic operations and the exponential function only) for these bounds.

In the error analysis both for $L$ and $H$, the integral

$$INT(x) \equiv \int_0^1 e^{-xv}(1-v)^{-1/2}\, dv$$

plays a basic role. Note that for any $0 < a < 1$,

$$\int_a^1 e^{-xv}(1-v)^{-1/2}\, dv \leq e^{-ax} 2\sqrt{1-a}. \tag{A.3}$$

Integrating by parts we find

$$\begin{aligned}\int_0^a e^{-xv}(1-v)^{-1/2}\,dv =&\frac{1}{x}\big[1-e^{-ax}(1-a)^{-1/2}\big]\\ &+\frac{1}{2x^2}\big[1-e^{-ax}(1-a)^{-3/2}\big]\\ &+\frac{3}{4x^3}\big[1-e^{-ax}(1-a)^{-5/2}\big]\\ &+\frac{15}{8x^4}\big[1-e^{-ax}(1-a)^{-7/2}\big]\\ &+\frac{105}{16x^4}\int_0^a e^{-xv}(1-v)^{-9/2}\,dv\\ \le&\frac{1}{x}+\frac{1}{2x^2}+\frac{3}{4x^3}+\frac{15}{8x^4}\,(1-a)^{-7/2}.\end{aligned} \tag{A.4}$$

A.1. **Proof of Proposition A.1.** Clearly $L(s)\ge 0$ for $s\ge 0$. Note that

$$\int_0^1 e^{-xv}(1-v)^{-1/2}\,dv \ge \frac{1-e^{-x}}{x}. \tag{A.5}$$

Expanding the square of $v=1-(1-v)$ and integrating by parts we find

$$\int_0^1 e^{-xv}v^2(1-v)^{-1/2}\,dv = -\frac{1}{x}-\frac{3}{2x^2}+\left(1+\frac{1}{x}+\frac{3}{4x^2}\right)INT(x). \tag{A.6}$$

Substuting here the estimates (A.3), (A.4) for $a=1/2$ and using (A.5) we find (after simplifications) an upper estimate on (nonnegative) $L(s)$ which implies that $L(s)\le 6$ for $s\ge 25$.

Next, making the change of variable $t=\sqrt{1-v}$ in both integrals in $L(s)$

$$L(s)\le s^2\frac{\int_0^1 e^{R_0 s(t^2-1)}\,dt}{\int_0^1 e^{s(t^2-1)}\,dt}\le s^2\,e^{s(1-R_0)}\le 6$$

for $0\le s\le 2$.

Finally consider $s\in[2,25]$. Using (A.5)

$$L(s)\le\frac{1}{1-e^{-s}}\,L_2(s),\qquad L_2(s)\equiv s^3\int_0^1 e^{-R_0 sv}\,\frac{v^2\,dv}{\sqrt{1-v}}.$$

Note that the first factor is $\le 1.157$. Hence it suffices to prove

$$L_2(s)\le 5.185,\qquad 2\le s\le 25.$$

We do this as follows. Assume for a moment that the following (crude) a priori bound holds

$$|L_2'(s)|\le 75,\qquad 2\le s\le 25. \tag{A.7}$$

Consider for some $N_e$ the mesh $s_j=2+jh$, $j=0,1,\cdots,N_e$ where $h=(25-2)/N_e$. By (A.7)

$$|L_2(s)|\le|L_2(s_j)|+75h,\qquad s\in[s_{j-1},s_j],\qquad j=1,2,\cdots,N_e. \tag{A.8}$$

The value of $L_2$ at a point is estimated as follows. $L_2$ is $s^3$ times (A.6) with $x=R_0s$. Making the change of variable $t=\sqrt{1-v}$ in $INT(x)$ we obtain for any $N_i$

$$INT(R_0s)=2\int_0^1 e^{R_0 s(t^2-1)}\,dt\le\frac{2}{N_i}\sum_{k=1}^{N_i}e^{R_0 s((k/N_i)^2-1)} \tag{A.9}$$

where the integral is estimated by the right rectangle sum using the monotonic growth of $e^{R_0 s(t^2-1)}$ for $t \in [0,1]$. Now we can just increase $N_e$, $N_i$ until $\max_{j=1,2,\cdots,N_e}$ of the RHS in (A.8) (where (A.9) has been substituted) becomes $\leq 5.185$. It turns out that $N_e = 8,000$ and $N_i = 12,000$ suffice and for these values the maximum is $\leq 5.162$. We must also check that the error does not accumulate too much when we sum $12,000$ numbers, the precision of each being 10 digits. In the worst scenario, the mantissa after each addition accumulates an error of 0.5 in the 10th digit. After 12,000 additions the error is 6 in the 7th digit. This means that our result has at least 5 correct digits. And in the above explanation we have kept only 4 digits in arriving at the number 5.162.

It remains to prove the bound (A.7). Differentiating $G_2(s)$ we find

$$|G_2'(s)| \leq \max\left\{3s^2\int_0^1 e^{-R_0 sv}v^2\,\frac{dv}{\sqrt{1-v}},\ R_0 s^3\int_0^1 e^{-R_0 sv}v^3\,\frac{dv}{\sqrt{1-v}}\right\}.$$

The first integral is estimated using (A.6), (A.4). For the second integral: start by rewriting the expression in terms of $INT(x)$ (as we have done in (A.6)) and then use (A.4). These estimates imply (A.7) for $s \geq 2$.

A.2. **Proof of Proposition A.2.** Writing $v = 1-(1-v)$ and integrating by parts as above we can rewrite $H(s)$ in terms of $INT(s^2)$

$$H(s) = 2s^2 - 1 - \frac{2}{INT(s^2)} = 2s^2 - 1 - \frac{1}{\int_0^1 e^{s^2(u^2-1)}\,du}. \tag{A.10}$$

Combining (A.10) with (A.4), the estimate (A.2) follows. We need also a *lower* estimate on $H(s)$. As in (A.4), for any $a \in (0,1)$

$$\begin{aligned}\int_0^1 e^{-xv}\,\frac{dv}{\sqrt{1-v}} \geq \int_0^a e^{-xv}\,\frac{dv}{\sqrt{1-v}} &= \frac{1}{x} + \frac{1}{2x^2} + \frac{3}{4x^3} + \frac{15}{8x^4}\\ &- e^{-ax}\left[\frac{1}{x}(1-a)^{-1/2} + \frac{1}{2x^2}(1-a)^{-3/2} + \frac{3}{4x^3}(1-a)^{-5/2} + \frac{15}{8x^4}(1-a)^{-7/2}\right].\end{aligned} \tag{A.11}$$

Together with (A.10), this implies

$$H(\sqrt{x}) \geq \frac{NUMER_a(x)}{DENOM_a(x)}$$

where $DENOM_a(x)$ is the RHS in (A.11) and $NUMER_a(x)$ is a certain, explicit function. It is convenient to take $a = 4/5$. Then it is elementary to check $DENOM_a(x) > 0$ for $x \geq 9$. We find

$$\begin{aligned}x^2 NUMER(x) &\geq 1 + \frac{3}{x} - \frac{15}{8x^2} - 2\sqrt{5}x^2 e^{-4x/5}\left(1 + \frac{5}{2x} + \frac{75}{4x^2} + \frac{1875}{8x^3}\right)\\ &\geq \frac{211}{216} - \frac{27031}{17496}\sqrt{5}x^2 e^{-4x/5}\end{aligned} \tag{A.12}$$

where we have discarded $\frac{3}{x}$ and evaluated the monotonically decreasing terms $-\frac{15}{8x^2}$ and the term in the paranthesis at $x = 9$. It is now elementary to check that the RHS in (A.12) is $> 0$ for $x \geq 9$.

We have proved that $H(s) > 0$ for $s \geq 3$. From this and (A.2) it follows that

$$\int_6^\infty |H(s)|ds \leq 0.233. \tag{A.13}$$

On the interval $[3,6]$ where $H$ does not change sign and so $|H|$ is smooth we use the trapezoidal integration scheme with a rigorous error estimation. However on $[0,3]$ (where $H$ may and indeed does change sign) we use the following argument. Fix $N_e$ and $N_i$ and consider the mesh $s_j = jh$, $j = 0, 1, \cdots, N_e$ where $h = (3-0)/N_e$. Using the left and right rectangles as in (A.9) we find that

$$H_j^- \leq H(s) \leq H_j^+, \qquad s \in [s_j, s_{j+1}], \quad j = 0, 1, \cdots N_e$$

where for any $N_i$

$$\begin{aligned} H_j^- &\equiv 2s_j^2 - \frac{1}{\frac{1}{N_i}\sum_{k=0}^{N_i-1} e^{s_{j+1}^2((k/N_i)^2-1)}} \\ H_j^+ &\equiv 2s_{j+1}^2 - \frac{1}{\frac{1}{N_i}\sum_{k=0}^{N_i-1} e^{s_j^2((k/N_i)^2-1)}}. \end{aligned}$$

This gives

$$|H(s)| \leq \max\left\{|H_j^-|, |H_j^+|\right\}, \qquad s \in [s_j, s_{j+1}], \quad j = 0, 1, \cdots N_e - 1$$

and

$$\int_0^3 |H(s)|\, ds \leq \frac{3}{N_e} \sum_{j=0}^{N_e-1} \max\left\{|H_j^-|, |H_j^+|\right\}$$

for any choice of $N_{i,e}$. Increasing these numbers we improve the estimate. It turns out that for $N_e = N_i = 3,000$ we get $\int_0^3 |H(s)|\, ds \leq 2.242$ (Maple numerical integration suggests the estimate 2.200). Again we only use the computer to evaluate arithmetic operations and exponents. Also the precision is 10 digits so that after $9 \cdot 10^6$ summations a worst possible error of 0.5 in the 10th digit at each step becomes an error of 4.5 in the 4th digit. Thus we write

$$\int_0^3 |H(s)|\, ds \leq 2.247. \tag{A.14}$$

Finally we evaluate the integral over $[3,6]$ (where $H$ is positive) via the trapezoidal scheme to obtain

$$\begin{aligned} \int_3^6 H(s)\, ds &= h \sum_{j=1}^{N_e-1} H(s_j) + \frac{H(3) + H(6)}{2} h + ERR_e + ERR_i \\ |ERR_e| &\leq \frac{(6-3)\, h^2}{12} \max_{s \in [3,6]} |H''(s)| \\ |ERR_i| &\leq (\text{maximal error in computation of } H(s_j) \text{ over all steps}) \times (6-3) \end{aligned}$$

where $s_j = 3 + jh$, $j = 0, 1, \cdots, N_e$, $h = (6-3)/N_e$. At each $s_j$ we will compute $H(s_j)$ also by the trapezoidal scheme with an explicit error estimate. We demand $|ERR_e| \leq 0.05$ and $|ERR_i| \leq 0.05$. Let us first find the required $N_e$. Assume for a moment a (crude) estimate

$$|H''(s)| \leq 24s^4 + 20s^2 + 4, \qquad s \geq 0. \tag{A.15}$$

Then

$$\frac{3}{12}\left(\frac{3}{N_e}\right)^2 (24 \cdot 6^4 + 2 \cdot 6^2 + 4) \leq 0.05 \qquad \text{implies} \qquad N_e = 1,197.$$

Now consider $N_i$. For any $s \in [3,6]$ we compute $\int_0^1 F_s(u)du$ where $F_s(u) \equiv e^{s^2(u^2-1)}$ (see (A.10)) by the trapezoidal scheme. Note

$$\max_{u\in[0,1]} \left|\frac{\partial^2}{\partial u^2} F_s(u)\right| = 4s^4 + 2s^2.$$

Now if $\int_0^1 F_s(u)du = \sum(s) + err_i(s)$, then the error in the computation of $1/\int_0^1 F_s(u)du$ (which enters $H(s)$) is

$$\left|\frac{1}{\int_0^1} - \frac{1}{\sum(s)}\right| \le \frac{|err_i(s)|}{|\int_0^1| \cdot |\sum(s)|}.$$

From $\int_0^1 e^{s^2(u^2-1)}du = \frac{1}{2}\int_0^1 e^{-s^2 t}\frac{dv}{\sqrt{1-v}} \ge \frac{1-e^{-s^2}}{2s^2}$ we get a lower bound

$$\min_{u\in[3,6]} \int_0^1 e^{s^2(u^2-1)}du \ge 0.01388.$$

When computing the sums $\sum(s)$ below we also store the minimal value of $\sum(s_j)$ over $j = 0, 1, \cdots, N_e$ and verify at the end that it is also $\ge 0.01388$. Thus we demand

$$\begin{aligned}|ERR_i| &\le (6-3) \max_{j=0,1,\cdots,N_e} \frac{|err_i(s_j)|}{(0.01388)^2} \\ &\le \frac{1}{12} \max_{j=0,1,\cdots,N_e} \left(\frac{1}{N_i(s_j)}\right)^2 4(s_j^4 + s_j^2/2)\,\frac{3}{(0.01388)^2} \le 0.05\end{aligned}$$

which holds if we choose

$$N_i(s_j) \ge 323\sqrt{s_j^4 + s_j^2/2}, \qquad j = 0, 1, \cdots, N_e$$

where $s_j = 3 + j\frac{6-3}{N_e}$. A computer evaluation (again using only elementary operations) then shows that $\int_3^6 H(s)ds \le 0.208$ with our guaranteed precision $\pm 0.1$. Also we keep 10 digits and the total number of summands is $\le 1,197 \cdot (323\sqrt{6^4 + 6^2/2} + 1) \le 1.5 \cdot 10^8$. Hence if the worst error of 0.5 in the 10th digit is made at each step, the accumulated error does not exceed 7.5 in the 4th digit (in the mantissa of the number $2.08 \cdot 10^{-1}$). Thus

$$\int_3^6 H(s)\,ds \le 0.208 + 0.1 + 0.00075 = 0.30875 \le 0.309. \tag{A.16}$$

(Maple numerical integration gives $\int_3^6 H(s)\,ds \le 0.208$.)

Collecting (A.13), (A.14), (A.16) we conclude

$$\int_0^\infty |H(s)|\,ds \le 0.233 + 2.247 + 0.309 = 2.789 < 2.8$$

which completes the proof up to (A.15). To prove (A.15) denote

$$K(x) \equiv -2 + 2x \cdot \frac{\int v}{\int 1}, \qquad \int v^l \equiv \int_0^1 e^{-xv}\frac{v^l\,dv}{\sqrt{1-v}}, \quad l = 0, 1, 2, \cdots.$$

Then

$$H(s) = K(s^2), \qquad H'(s) = 2s \cdot K'(s^2). \tag{A.17}$$

We find

$$K'(x) = 2 \cdot \frac{\int v}{\int 1} + 2x \cdot \left[ - \frac{\int v^2}{\int 1} + \left( \frac{\int v}{\int 1} \right)^2 \right].$$

If we single out the positive and the negative terms and use the fact that each fraction is bounded from above by 1, we obtain

$$-2x \leq K'(x) \leq 2 + 2x, \qquad x \geq 0.$$

and from (A.17)

$$|H'(s)| \leq 4s^3 + 4s, \qquad s \geq 0.$$

After computing $H''(s)$ in terms of $K'(x)$, $K''(x)$ in the same way, we arrive at (A.15).

## REFERENCES

[AbSt] M. Abramowitz and I. A. Stegun, *Handbook of Mathematical Functions, with Formulas, Graphs and Mathematical Tables.* Dover, New York, 1966.

[AFNvM] M. Adler, P. J. Forrester, T. Nagao and P. van Moerbeke, *Classical skew orthogonal polynomials and random matrices.* J. Statist. Phys. **99** (2000), 141–170.

[AvM] M. Adler and P. van Moerbeke, *Toda versus Pfaff lattice and related polynomials.* Duke Math. J. **112** (2002), 1–58.

[Be] C. W. J. Beenakker, *Universality for Brézin and Zee's spectral correlator.* Nuclear Phys. B **422** (1994), 515–520.

[BI] P. Bleher and A. Its, *Semiclassical asymptotics of orthogonal polynomials, Riemann-Hilbert problem, and universality in the matrix model.* Ann. of Math. (2) **150** (1999), 185–266.

[BrZ] E. Brézin and A. Zee, *Universality of the correlations between eigenvalues of large random matrices.* Nuclear Phys. B **402** (1993), 613–627.

[D] P. Deift, *Orthogonal Polynomials and Random Matrices: a Riemann-Hilbert Approach.* Courant Lecture Notes in Mathematics, 3. New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999.

[DIZ] P. A. Deift, A. R. Its and X. Zhou, *A Riemann-Hilbert approach to asymptotic problems arising in the theory of random matrix models, and also in the theory of integrable statistical mechanics.* Ann. of Math. (2) **146** (1997), 149–235.

[DKMVZ1] P. Deift, T. Kriecherbauer, K. T.-R. McLaughlin, S. Venakides and X. Zhou, *Uniform asymptotics for polynomials orthogonal with respect to varying exponential weights and applications to universality questions in random matrix theory.* Comm. Pure Appl. Math. **52** (1999), 1335–1425.

[DKMVZ2] P. Deift, T. Kriecherbauer, K. T.-R. McLaughlin, S. Venakides and X. Zhou, *Strong asymptotics of orthogonal polynomials with respect to exponential weights.* Comm. Pure Appl. Math. **52** (1999), 1491–1552.

[DVZ] P. Deift, S. Venakides and X. Zhou, *New results in small dispersion KdV by an extension of the steepest descent method for Riemann-Hilbert problems.* Int. Math. Res. Not. **1997** (1997), 286–299.

[DZ] P. Deift and X. Zhou, *A steepest descent method for oscillatory Riemann-Hilbert problems. Asymptotics for the MKdV equation.* Ann. of Math. (2) **137** (1993), 295–368.

[Dy1] F. J. Dyson, *Statistical theory of the energy levels of complex systems. I—III.* J. Mathematical Phys. **3** (1962) 140–156, 157–165 and 166–175.

[Dy2] F. J. Dyson, *A note on correlations between eigenvalues of a random matrix.* Comm. Math. Phys. **19** (1970), 235–250.

[DyM] F. J. Dyson and M. L. Mehta, *Statistical theory of the energy levels of complex systems. IV and V.* J. Mathematical Phys. **4** (1963) 701–712 and 713–719.

[E] B. Eynard, *Asymptotics of skew orthogonal polynomials.* J. Phys. A **34** (2001), 7591–7605.

[FoIKi] A. S. Fokas, A. R. Its and A.V. Kitaev, *Discrete Painleve equations and their appearance in quantum gravity.* Comm. Math. Phys. **142** (1991), 313–344.

[F] P. J. Forrester, *Log–gases and Random Matrices,* book in progress. `www.ms.unimelb.edu.au/~matpjf/matpjf.html`

[FNH] P. J. Forrester, T. Nagao and G. Honner, *Correlations for the orthogonal–unitary and symplectic–unitary transitions at the hard and soft edges.* Nuclear Phys. B **553** (1999), 601–643.

[HWe] G. Hackenbroich and H. A. Weidenmüller, *Universality of Random–Matrix results for non-Gaussian ensembles.* Phys. Rev. Lett. **74** (1995), 4118–4121.

[GM-GW] T. Guhr, A. Müller-Gröling, H. A. Weidenmüller, *Random Matrix Theories in Quantum Physics: Common Concepts,* Phys. Rept. **299** (1998) 189-425. `www.arxiv.org/abs/cond-mat/9707301`

[J] K. Johansson, *On fluctuations of eigenvalues of random Hermitian matrices.* Duke Math. J. **91** (1998), 151–204.

[MaM] G. Mahoux and M. L. Mehta, *A method of integration over matrix variables. IV.* J. Physique I France **1** (1991), 1093–1108.

[M] M. L. Mehta, *Random Matrices,* 2nd Ed., Academic Press, San Diego, 1991.

[M2] M. L. Mehta, *A note on correlations between eigenvalues of a random matrix.* Comm. Math. Phys. **20** (1971), 245–250.

[MGa] M. L. Mehta and M. Gaudin, *On the density of eigenvalues of a random matrix.* Nuclear Phys. **18** (1960), 420–427.

[MhSa] H. N. Mhaskar and E. B. Saff, *Extremal problems for polynomials with exponential weights.* Trans. Amer. Math. Soc. **285** (1984), 203–234.

[NW] T. Nagao and M. Wadati, *Correlation functions of random matrix ensembles related to classical orthogonal polynomials, I—III.* J. Phys. Soc. Japan **60** (1991), 3298–3322, ibid. **61** (1992), 78–88 and 1910–1918.

[PS] L. Pastur and M. Shcherbina, *Universality of the local eigenvalue statistics for a class of unitary invariant random matrix ensembles.* J. Statist. Phys. **86** (1997), 109–147.

[Ra] E. A. Rakhmanov, *Asymptotic properties of orthogonal polynomials on the real axis.* Mat. Sb. (N.S.) **119(161)** (1982), 163–203, 303. (Russian). English transl. in: Math. Sb. (N.S.) **47** (1984), 155–193.

[RSa] Z. Rudnick and P. Sarnak, *Zeros of principal L-functions and random matrix theory.* Duke Math. J. **81** (1996), 269–322.

[SaTo] E. B. Saff and V. Totik, *Logarithmic Potentials with External Fields,* Grundlehren der Mathematischen Wissenschaften, 316. Springer–Verlag, Berlin, 1997.

[Si] B. Simon, *Trace Ideals and Their Applications,* London Mathematical Society Lecture Notes Series, 35. Cambridge University Press, Cambridge–New York, 1979.

[So] A. Soshnikov, *Universality at the edge of the spectrum in Wigner random matrices.* Comm. Math. Phys. **207** (1999), 697–733.

[SeV] M. K. Sener and J. J. M. Verbaarschot, *Universality in chiral random matrix theory at* $\beta = 1$ *and* $\beta = 4$. Phys. Rev. Lett. **81** (1998), 248–251.

[St1] A. Stojanovic, *Des problèmes asymptotiques dans la théorie spectrale des matrices aléatoires.* Ph.D. Thesis, Université Paris 7 Denis Diderot, 2003.

[St2] A. Stojanovic, *Universality in orthogonal and symplectic invariant matrix models with quartic potential.* Math. Phys. Anal. Geom. **3** (2000), 339–373 (2001). *Errata:* ibid., in press, 2004.

[St3] A. Stojanovic, *Universalité pour des modèles matriciels à symétrie orthogonale ou symplectique et à potentiel quartique.* Preprint 02-07-098, revised version of Preprint 00-01-06. `www.physik.uni-bielefeld.de/bibos/preprints`

[Sz] G. Szegö, *Orthogonal Polynomials,* Amer. Math. Soc. Colloq. Publ., v. 23. Amer. Math. Soc., New York, 1939.

[TW1] C. A. Tracy and H. Widom, *Fredholm determinants, differential equations and matrix models.* Comm. Math. Phys. **163** (1994), 33–72.

[TW2] C. A. Tracy and H. Widom, *Correlation functions, cluster functions, and spacing distributions for random matrices.* J. Statist. Phys. **92** (1998), 809–835.

[V] J. Verbaarschot, *Topics in Random Matrix Theory,* lecture notes. `tonic.physics.sunysb.edu/~verbaarschot/lecture/`

[W] H. Widom, *On the relation between orthogonal, symplectic and unitary matrix ensembles.* J. Statist. Phys. **94** (1999), 347–363.

Deift: Department of Mathematics, Courant Institute of Mathematical Sciences, New York University, New York, NY 10012
*E-mail address*: `deift@cims.nyu.edu`

Gioev: Department of Mathematics, University of Pennsylvania, Philadelphia, PA 19104, and Department of Mathematics, Courant Institute of Mathematical Sciences, New York University, New York, NY 10012
*E-mail address*: `gioev@math.upenn.edu, gioev@cims.nyu.edu`